

Jacques Troué

On a theorem on the change of base point in a path connected space

Czechoslovak Mathematical Journal, Vol. 23 (1973), No. 1, 123–124

Persistent URL: <http://dml.cz/dmlcz/101150>

Terms of use:

© Institute of Mathematics AS CR, 1973

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON A THEOREM ON THE CHANGE OF BASE POINT
IN A PATH CONNECTED SPACE

JACQUES TROUÉ, Montreal

(Received January 25, 1972)

In [1], PUPPE has proved by elementary means the following

Theorem 1. *If $A \subset X$, $B \subset Y$ are cofibrations, where A and B are closed subsets of X, Y respectively, then also $(X \times B) \cup (A \times Y) \subset X \times Y$ is a cofibration.*

There is no doubt that this general theorem will soon be part of any introduction to homotopy theory. The object of this note is to remark that it leads to an immediate proof of the following basic.




Fig. 1.

Theorem 2. (cf. [2] pp. 380–382). Let X be a space with a non-degenerate base point x_0 , and assume that $\{x_0\}$ is closed in X . Let w be a path in the space Y . Then there is a bijection depending only on $[w]$, homotopy class of w

$$h_{[w]} : [X, x_0; Y, w(1)] \rightarrow [X, x_0; Y, w(0)],$$

where a representative of $h_{[w]}[f_1]$ is obtained by deforming f_1 so that the image of x_0 moves along w from $w(1)$ to $w(0)$.

$[X, x_0; Y, w(i)]$ is the set of homotopy classes of maps $(X, x_0) \rightarrow (Y, w(i))$.

Proof. Recall (i) x_0 non-degenerate means $\{x_0\} \subset X$ is a cofibration, (ii) $I \subset I$ is a cofibration (obvious) and \dot{I} is closed in I .

Apply Theorem 1: $(x_0 \times I) \cup (X \times \dot{I}) \subset X \times I$ is a cofibration, i.e. each map $\{(x_0 \times I) \cup (X \times \dot{I})\} \times I' \rightarrow Y$ can be extended to a map $(X \times I) \times I' \rightarrow Y$ with a given map $(X \times I) \times 1 \rightarrow Y$ coinciding with the first map where defined; or (cf. fig.) a map to Y from the bottom – top – back faces and the section through x_0 can be extended to the whole cube.

Then the proof follows quickly:

A homotopy $f_1 \simeq f'_1$ in $(X, x_0; Y, w(1))$ is a map to Y of the back face, a map f_0 corresponding to f_1 is obtained by filling in the top face, and similarly for the bottom face; a homotopy of a path in Y rel $(w(0), w(1))$ is a map to Y of the section through x_0 . But then the Theorem of Puppe says that there is map from the front face to Y , i.e. $f_0 \simeq f'_0$ rel x_0 . So $h_{[w]}$ is well-defined. Interchanging the role of the indices 0 and 1, shows that its inverse is also well-defined.

Remark. The condition that $\{x_0\}$ is closed in X is necessary since Theorem 1 is false without it as a simple counter-example in [1] shows. This is not a severe restriction since already all points of a T_1 -space are submitted to it.

References

- [1] D. Puppe, Bemerkungen über die Erweiterung von Homotopien, Arch. der Math. (1967), 81–88.
- [2] E. Spanier, Algebraic Topology, McGraw-Hill, New York, 1966.

Author's address: Department of Mathematics, McGill University, Montreal, Canada.