

Ar. Meenakshi

On the partial ordering of almost definite matrices

*Czechoslovak Mathematical Journal*, Vol. 39 (1989), No. 1, 169–175

Persistent URL: <http://dml.cz/dmlcz/102289>

## Terms of use:

© Institute of Mathematics AS CR, 1989

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ON THE PARTIAL ORDERING OF ALMOST DEFINITE MATRICES

AR. MEENAKSHI, Annamalainagar\*)

(Received October 17, 1986)

## INTRODUCTION

Let  $C_{m \times n}$  be the linear space of complex matrices of order  $m \times n$ . Let  $C_n$  be the linear space of complex  $n$ -tuples. The vectors in  $C_n$  are denoted by lower case letters  $x, y, z$  etc. Matrices are denoted by capital letters such as  $A, B$ , etc. For  $A \in C_{m \times n}$ , let  $A^*, A^+, N(A)$  and  $R(A)$  denote conjugate transpose, Moore Penrose inverse, null space and range space respectively. A matrix  $A \in C_{n \times n}$  is said to be *almost definite* (a.d [4]) if for  $x \in C_n$ ,  $x^*Ax = 0 \Rightarrow Ax = 0$ .  $A$  is *positive semi definite* [4, 7] if  $\text{Re}(x^*Ax) \geq 0$ . A matrix is said to be *almost positive definite* (a.p.d) if it is both a.d and p.s.d. Recently, Mitra and Puri [9] have introduced the notion of *quasi positive definite matrix* (q.p.d).  $A$  is said to be q.p.d if  $A$  is p.s.d and  $\text{Re}(x^*Ax) = 0 \Rightarrow Ax = 0$ . A q.p.d matrix is always a.p.d [9]. These special types of matrices are widely used in the study of electrical networks and in linear electro-mechanical systems [4, 9]. For properties of a.d, a.p.d and q.p.d matrices one may refer [4, 9, 10]. It was pointed out by Duffin and Morley [4] that the unique transfer impedance in a general linear electromechanical system exists for every structure operator if and only if the constitutive operator is a.d. For terminology and representation of a general linear electromechanical system by a pair of equations, one may refer [4].

We are concerned with the hermitian positive semi definite partial ordering on complex matrices. For  $A, B \in C_{n \times n}$ ,  $A \geq B \Leftrightarrow A - B \geq 0 \Leftrightarrow (A - B)$  is *hermitian positive semi definite* (h.p.s.d). It is well known [2, p. 59] that for non-singular matrices  $A, B$  if  $A \geq B \geq 0$  then  $B^{-1} \geq A^{-1} \geq 0$ . This was extended to generalized inverses of certain types of pairs of singular matrices  $A \geq B \geq 0$  by Hans J. Werner [5] and independently by Hartwig [6]. Here we have extended their results for a wider class of a.p.d matrices. Further, some well known [3] matrix inequalities on a pair of h.p.s.d matrices have been extended to a pair of a.p.d matrices. As an application, it is shown that under certain conditions the monotonicity of the constitutive operators in linear electromechanical systems having the same structure operator is preserved for the corresponding transfer impedances.

\*) This work was done when the author visited Indian Statistical Institute, New Delhi, India.

## RESULTS

First we shall prove certain lemmas which will simplify the proof of the main result.

**Lemma 1.** *Let  $A, B \in C_{n \times n}$  such that  $A \geq B$ . If  $B$  is a.p.d (q.p.d) then  $A$  is a.p.d (q.p.d).*

*Proof.* Since  $A \geq B$ ,  $A - B = P$  is h.p.s.d.  $\Rightarrow x^*Px$  is real and  $x^*Px \geq 0$  for all  $x \in C_n \Rightarrow \operatorname{Re}(x^*Ax) \geq \operatorname{Re}(x^*Bx)$  and  $\operatorname{Im}(x^*Ax) = \operatorname{Im}(x^*Bx)$ . For  $x \in C_n$ ,  $x^*Ax = 0 \Rightarrow \operatorname{Im}(x^*Ax) = 0$  and  $\operatorname{Re}(x^*Ax) = 0 \Rightarrow \operatorname{Im}(x^*Bx) = 0$  and  $\operatorname{Re}(x^*Bx) + x^*Px = 0$ .

Since  $B$  and  $P$  are p.s.d,  $\operatorname{Re}(x^*Bx) \geq 0$  and  $x^*Px \geq 0$ . Hence  $x^*Ax = 0 \Rightarrow \operatorname{Im}(x^*Bx) = 0$ ,  $\operatorname{Re}(x^*Bx) = 0$  and  $x^*Px = 0 \Rightarrow x^*Bx = 0$  and  $x^*Px = 0$ .

On account of  $B$  a.d and  $P$  h.p.s.d,  $x^*Bx = 0 \Rightarrow Bx = 0$  and  $x^*Px = 0 \Rightarrow Px = 0$ . Therefore,  $Ax = Bx + Px = 0$ . Thus for  $x \in C_n$  if  $x^*Ax = 0$  then  $Ax = 0$ , and hence  $A$  is a.d. Since  $B$  and  $P$  are p.s.d.  $A$  is also p.s.d. Thus  $A$  is a.p.d. Similarly we can prove that  $A$  is q.p.d whenever  $B$  is q.p.d. Hence the Lemma.

**Lemma 2.** *Let  $A, B \in C_{n \times n}$  such that  $A \geq B$ . If  $B$  is a.p.d then  $R(B) \subseteq R(A)$  or equivalently  $N(A) \subseteq N(B)$ .*

*Proof.* Since  $B$  is a.p.d, by Lemma 1,  $A$  is a.p.d. On account of Lemma (2.1) of [9] both  $A$  and  $B$  are EP matrices. Therefore  $R(B) \subseteq R(A)$  is equivalent to  $N(A) \subseteq N(B)$ . Now we shall prove that  $N(A) \subseteq N(B)$  as follows:

$x \in N(A) \Rightarrow Ax = 0 \Rightarrow x^*Ax = 0 \Rightarrow x^*Bx + x^*Px = 0$ . Since  $B$  is p.s.d and  $P$  is h.p.s.d. as in the proof of Lemma 1, we get,  $x^*Bx = 0$  and  $x^*Px = 0$ . Since  $B$  is a.d,  $x^*Bx = 0 \Rightarrow Bx = 0$ .  $\Rightarrow x \in N(B)$ . Thus  $N(A) \subseteq N(B)$ . Hence the Lemma.

**Remark 1.** We note that the condition on  $B$  cannot be relaxed. For example, let

$$A = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} -1 & 2 \\ 0 & -1 \end{bmatrix}.$$

Both  $A$  and  $B$  are neither a.d nor p.s.d. Hence  $A$  and  $B$  are not a.p.d.

$$A - B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \geq 0$$

but  $R(B) \not\subseteq R(A)$ .

Thus Lemma 2 fails.

**Lemma 3.** *Let  $A, B \in C_{n \times n}$  such that  $A \geq B$ . If  $B$  is a.p.d then  $R(A) = R(B) \Leftrightarrow \operatorname{rank} A = \operatorname{rank} B$ .*

*Proof.*  $(\Rightarrow)$  is trivial and  $(\Leftarrow)$  is a direct consequence of Lemma 2.

**Theorem 1.** *Let  $A, B \in C_{n \times n}$  such that  $A \geq B$  and  $B$  be a.p.d. Then  $B^+ \geq A^+ \Rightarrow R(A) = R(B)$ .*

*Proof.* Since  $A \geq B$  and  $B$  is a.p.d, by Lemma 1,  $A$  is a.p.d and by Lemma 2,  $R(B) \subseteq R(A)$ . Since  $A$  and  $B$  are a.p.d, on account of Lemma 2.5 of [9],  $A^+$  and  $B^+$

are a.p.d with  $B^+ \geq A^+$ ; another application of Lemma 2, yields  $R(A^+) \subseteq R(B^+) \Rightarrow R(A^*) \subseteq R(B^*) \Rightarrow R(A) \subseteq R(B)$ . Thus  $R(A) = R(B)$ . Hence the Theorem.

For a complex matrix  $V$ , let  $\frac{1}{2}(V + V^*)$  be denoted by  $\text{Sym } V$ . It is known that  $V$  is q.p.d  $\Leftrightarrow \text{Sym } V$  is h.p.s.d and  $\text{rank } (\text{Sym } V) = \text{rank } (V)$ . [refer Lemma (2.8) in 9]

**Theorem 2.** Let  $A, B \in C_{n \times n}$  such that  $A \geq B$  and  $B$  be q.p.d. Then the following are equivalent.

- i)  $R(A) = R(B)$ ;
- ii)  $\text{rank } (A) = \text{rank } (B)$ ;
- iii)  $(\text{Sym } B)^+ \geq (\text{Sym } A)^+$ .

**Proof.** (i)  $\Leftrightarrow$  (ii). Since  $B$  is q.p.d,  $B$  is also a.p.d. Hence by Lemma 3, this equivalence is clear.

(ii)  $\Rightarrow$  (iii): Since  $A$  and  $B$  are q.p.d, by Lemma (2.8) of [9],  $\text{Sym } A \geq 0$ ;  $\text{Sym } B \geq 0$  and  $\text{rank } (\text{Sym } A) = \text{rank } (A)$ ;  $\text{rank } (\text{Sym } B) = \text{rank } (B)$ . Further  $A \geq B \Rightarrow A^* \geq B^* \Rightarrow A + A^* \geq B + B^* \Rightarrow \text{Sym } A \geq \text{Sym } B$ . Thus  $(\text{Sym } A) \geq (\text{Sym } B) \geq 0$  and by (ii)  $\text{rank } (\text{Sym } A) = \text{rank } (\text{Sym } B)$ . Now by Theorem 1 of [5],  $(\text{Sym } B)^+ \geq (\text{Sym } A)^+$ . Thus (iii) holds.

Conversely,  $A$  and  $B$  are q.p.d with  $A \geq B \Rightarrow \text{Sym } A \geq \text{Sym } B \geq 0$  and by (iii)  $(\text{Sym } B)^+ \geq (\text{Sym } A)^+$ . Now by Theorem 1 of [5],  $\text{rank } (\text{Sym } A) = \text{rank } (\text{Sym } B)$  which implies  $\text{rank } (A) = \text{rank } (B)$ . Thus (ii) holds. Hence the Theorem.

**Remark 2.** We note that the condition on  $B$  cannot be relaxed. This is illustrated in the following.

$$A = \begin{bmatrix} 2 + i & 1 + i \\ 1 + i & 1 + i \end{bmatrix} \text{ is q.p.d by Lemma (2.8) of [9],}$$

$$B = \begin{bmatrix} 1 + i & i \\ i & i \end{bmatrix} \text{ is a.p.d but not q.p.d.}$$

$A$  and  $B$  are nonsingular, hence  $R(A) = R(B)$  and  $\text{rank } A = \text{rank } B$ .

$$A - B = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \geq 0,$$

$$\text{Sym } B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} = (\text{Sym } B)^+,$$

$$\text{Sym } A = \begin{bmatrix} 2 & 1 \\ 1 & 1 \end{bmatrix}, \quad (\text{Sym } A)^+ = (\text{Sym } A)^{-1} = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix},$$

$$(\text{Sym } B)^+ - (\text{Sym } A)^+ = \begin{bmatrix} 0 & 1 \\ 1 & -2 \end{bmatrix} \not\geq 0.$$

Thus the Theorem fails.

**Remark 3.** In particular if  $A \geq B \geq 0$ , then  $\text{Sym } A = A$  and  $\text{Sym } B = B$ ; and the above Theorem reduces to the following known results.

**Corollary 1** (Theorem 1 in [6]). Let  $A, B \in C_{n \times n}$  such that  $A \geq B \geq 0$  then  $B^+ \geq A^+ \Leftrightarrow R(A) = R(B)$ .

**Corollary 2.** (Theorem 1 in [5]): For  $A, B \in C_{n \times n}$ , any two of the following three conditions imply the other one.

- 1)  $A \geq B \geq 0$ ;
- 2)  $\text{rank}(A) = \text{rank}(B)$ ;
- 3)  $B^+ \geq A^+ \geq 0$ .

*Proof.* (1) and (2)  $\Rightarrow$  (3), (1) and (3)  $\Rightarrow$  (2) follow from Theorem 2, using  $\text{Sym } A = A$ ; and  $\text{Sym } B = B$ .

The proof for (2) and (3)  $\Rightarrow$  (1) runs as follows: Since  $\text{rank } A = \text{rank } A^+$  and  $\text{rank } B = \text{rank } B^+$ ;  $B^+ \geq A^+ \geq 0$  and  $\text{rank } A^+ = \text{rank } B^+ \Rightarrow (A^+)^+ \geq (B^+)^+ \geq 0 \Rightarrow A \geq B \geq 0$ . Thus (1) holds. Hence the corollary.

**Corollary 3.** Let  $A, B \in C_{n \times n}$  such that  $A \geq B$  and  $B$  be q.p.d. Then  $B^+ \geq A^+$  imply the following equivalent statements.

- i)  $R(A) = R(B)$ ;
- ii)  $\text{rank}(A) = \text{rank}(B)$ ;
- iii)  $(\text{Sym } B)^+ \geq (\text{Sym } A)^+$ .

*Proof.* The equivalence of these statements follows from Theorem 2. Now by Theorem 1,  $A \geq B$  and  $B^+ \geq A^+ \Rightarrow R(A) = R(B)$  holds for any a.p.d matrix  $B$ . Since  $B$  is q.p.d,  $B$  is also a.p.d. Thus,  $A \geq B$  and  $B^+ \geq A^+ \Rightarrow R(A) = R(B)$ . Hence the corollary.

**Remark 4.** In particular for  $A \geq B \geq 0$ , the converse holds. However in general the converse need not be true can be seen from the following example.

Let

$$A = \begin{bmatrix} 2 & i \\ i & 2 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & i \\ i & 1 \end{bmatrix}$$

both  $A$  and  $B$  are q.p.d.

$$A - B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \geq 0.$$

$$\text{Sym } A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix} \quad \text{and} \quad \text{Sym } B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$(\text{Sym } A)^+ = \begin{bmatrix} 1/2 & 0 \\ 0 & 1/2 \end{bmatrix} \quad \text{and} \quad (\text{Sym } B)^+ = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$(\text{Sym } B)^+ \geq (\text{Sym } A)^+.$$

Since  $A$  and  $B$  are nonsingular,  $R(A) = R(B)$ .  $\text{rank } A = \text{rank } B$ . Thus statement (i), (ii) and (iii) hold.

$$B^+ = B^{-1} = \frac{1}{2} \begin{bmatrix} 1 & -i \\ -i & 1 \end{bmatrix} \quad \text{and} \quad A^+ = A^{-1} = \frac{1}{5} \begin{bmatrix} 2 & -i \\ -i & 2 \end{bmatrix},$$

$$10(B^+ - A^+) = \begin{bmatrix} 1 & -3i \\ -3i & 1 \end{bmatrix} \not\geq 0 \Rightarrow B^+ \not\geq A^+ .$$

Thus the converse is not true.

For a partitioned matrix

$$M = \begin{bmatrix} A & B \\ C & D \end{bmatrix},$$

the matrix  $D - CA^+B$  is called *generalized Schur complement of A in M* and is denoted by  $M/A$ . Now we shall generalize the results found in [3] in the following way.

**Lemma 4.** Let  $H$  and  $K$  be a.p.d matrices of order  $n$  such that  $H \geq -K$ . Let  $X$  and  $Y$  be  $n \times m$  matrices satisfying

$$(1.1) \quad N(H) \subseteq N(X^*); \quad N(K) \subseteq N(Y^*)$$

and

$$(1.2) \quad X^*H^+ = (H^+X)^*; \quad Y^*K^+ = (K^+Y)^* .$$

Then the  $m \times m$  Hermitian matrix

$$Q = X^*H^+X + Y^*K^+Y - (X + Y)^*(H + K)^+(X + Y) \geq 0 .$$

*Proof.* Let us consider

$$L = \begin{bmatrix} H & X \\ X^* & X^*H^+X \end{bmatrix} \quad \text{and} \quad M = \begin{bmatrix} K & Y \\ Y^* & Y^*K^+Y \end{bmatrix}$$

Since  $H$  is a.p.d, by Lemma (2.1) of [9],  $H$  is EP. Hence  $N(H^*) = N(H) \subseteq N(X^*)$ . Further the generalized Schur complement of  $H$  in  $L$ ,  $L/H = 0$ . Hence by corollary under Theorem 1 of [3],  $\text{rank } L = \text{rank } H$ . By applying Theorem 3 of [8] and using  $X^*H^+ = (H^+X)^*$ ;  $H$  is a.p.d. implies  $L$  is a.p.d. Hence  $L$  is p.s.d. Similarly we can see that  $M$  is p.s.d. Hence  $L + M$  is p.s.d. Since  $H \geq -K$ ,  $H + K \geq 0$ , which implies  $H + K$  is hermitian. By (1.2)  $X^*H^+X + Y^*K^+Y$  is hermitian. Hence,

$$L + M = \begin{bmatrix} H + K & X + Y \\ (X + Y)^* & X^*H^+X + Y^*K^+Y \end{bmatrix} \text{ is hermitian .}$$

Thus  $L + M \geq 0$ . Now, by a result of Albert [1],

$$Q = X^*H^+X + Y^*K^+Y - (X + Y)^*(H + K)^+(X + Y) = L + M/H + K \geq 0 .$$

Hence the Lemma.

**Remark 5.** In particular if  $H$  and  $K$  are h.p.s.d then the conditions  $H \geq -K$  and (1.2) hold automatically and the above Lemma 4, reduces to Theorem 5 of [3].

**Theorem 3.** Let  $H$  and  $K$  be a.p.d. matrices of order  $n$  such that  $H \geq -K$ ; partitioned in the form.

$$(1.3) \quad H = \begin{bmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{bmatrix} \quad \text{and} \quad K = \begin{bmatrix} K_{11} & K_{12} \\ K_{21} & K_{22} \end{bmatrix}$$

with

$$(1.4) \quad H_{21}H_{11}^+ = (H_{11}^+H_{21}^*)^* = (H_{11}^+H_{12})^*$$

and

$$(1.5) \quad K_{21}K_{11}^+ = (K_{11}^+K_{21}^*)^* = (K_{11}^+K_{12})^*.$$

Then,  $H + K/H_{11} + K_{11} \geq H/H_{11} + K/K_{11} \geq 0$ .

Proof. Since  $H$  and  $K$  are a.p.d. by Lemmas (2.3) and (2.4) of [9],  $H_{11}$  and  $K_{11}$  are a.p.d. and (1.1) hold for the matrices  $H$  and  $K$ . Now, by definition of generalized Schur complement [3], we have

$$H + K/H_{11} + K_{11} = H_{22} + K_{22} - (H_{21} + K_{21})(H_{11} + K_{11})^+ (H_{12} + K_{12}).$$

By using (1.4), (1.5) and applying Lemma 4, we get

$$\begin{aligned} H + K/H_{11} + K_{11} &\geq H_{22} + K_{22} - (H_{21}H_{11}^+H_{21}^* + K_{21}K_{11}^+K_{21}^*) = \\ &= H_{22} + K_{22} - H_{21}H_{11}^+H_{12} - K_{21}K_{11}^+K_{12} = \\ &= (H_{22} - H_{21}H_{11}^+H_{12}) + (K_{22} - K_{21}K_{11}^+K_{12}) = H/H_{11} + K/K_{11}. \end{aligned}$$

Thus,  $H + K/H_{11} + K_{11} \geq H/H_{11} + K/K_{11}$ .

Since  $H_{21}H_{11}^+ = (H_{11}^+H_{12})^*$  and  $K_{21}K_{11}^+ = (K_{11}^+K_{12})^*$  by applying Theorem 1 of [8] for the a.p.d. matrices  $H$  and  $K$ , we see that  $H/H_{11}$  and  $K/K_{11}$  are both a.p.d., hence p.s.d. Since  $H \geq -K$ ,  $H + K$  is hermitian  $\Rightarrow H_{22} + K_{22}$  is hermitian. By using (1.4),  $H_{21}H_{11}^+H_{12}$  is hermitian and using (1.5),  $K_{21}K_{11}^+K_{12}$  is hermitian. Hence  $H/H_{11} + K/K_{11} = H_{22} + K_{22} - H_{21}H_{11}^+H_{12} - K_{21}K_{11}^+K_{12}$  is hermitian. Since  $H/H_{11}$  and  $K/K_{11}$  are p.s.d.  $H/H_{11} + K/K_{11}$  is also p.s.d. Thus  $H/H_{11} + K/K_{11} \geq 0$ . Hence the theorem.

Remark 6. In a special case if  $H$  and  $K$  are h.p.s.d. matrices of the form (1.3) then  $H \geq -K$ , the conditions (1.4) and (1.5) automatically hold, and Theorem 3 reduces to the following known result.

**Corollary.** (Theorem 6 in [3]): *Let  $H$  and  $K$  be h.p.s.d. matrices partitioned in the form (1.3) then  $H + K/H_{11} + K_{11} \geq H/H_{11} + K/K_{11}$ .*

#### APPLICATION TO LINEAR ELECTROMECHANICAL SYSTEMS

Let us consider two linear electromechanical systems with constitutive operators  $H$  and  $K$  and having the same structure operator  $A$ . For terminology and notation the reader may refer Duffin and Morley [4]. If  $H$  and  $K$  are a.d operators then their transfer impedances  $\psi(H)$  and  $\psi(K)$  exist and both are a.d. If  $H$  and  $K$  are partitioned in the form (1.3) then by Theorem 7 of [4],

$$\psi(H) = (A^+)^* (H_{22} - H_{21}H_{11}^+H_{12}) A^+ = (A^+)^* (H/H_{11}) A^+$$

$$\psi(K) = (A^+)^* (K_{22} - K_{21}K_{11}^+K_{12}) A^+ = (A^+)^* (K/K_{11}) A^+.$$

If we assume that the constitutive operators  $H$  and  $K$  satisfies (1.4) and (1.5)

respectively and  $H \geq -K$ , then by Theorem 3,

$$\begin{aligned} H/H_{11} \geq -K/K_{11} &\Rightarrow (A^+)^*(H/H_{11})A^+ \geq -(A^+)^*(K/K_{11})A^+ \Rightarrow \\ &\Rightarrow \psi(H) \geq -\psi(K). \end{aligned}$$

Thus the monotonicity of the constitutive operators is preserved for the corresponding transfer impedances.

**Acknowledgement.** The author wishes to express her sincere thanks to Professor S. K. Mitra for valuable discussions.

#### References

- [1] *A. Albert*: Conditions for positive and nonnegative definiteness in terms of pseudo inverse, SIAM. J. Appl. Math. 17 (1969) 434—440.
- [2] *R. Bellman*: Introduction to Matrix Analysis, McGraw Hill, New York (1960).
- [3] *D. Carlson, E. Haynsworth and Th. Markham*: Generalization of Schur complement by means of Moore Penrose inverse, SIAM. J. Appl. Math. 26 (1974) 169—175.
- [4] *R. J. Duffin and T. D. Morley*: Almost definite operators and electromechanical systems, SIAM. J. Appl. Math. 35 (1978) 21—30.
- [5] *Hans J. Werner*: On the Matrix Monotonicity of generalized inversion, J. Linear Algebra and Appl. 27 (1979) 141—145.
- [6] *R. E. Hartwig*: A note on the partial orderings on positive semi definite matrices, J. Linear and Multilinear Alg. 6 (1978) 223—226.
- [7] *T. O. Lewis and T. G. Newman*: Pseudoinverses of positive semidefinite matrices, SIAM. J. Appl. Math. 10 (1968) 701—703.
- [8] *AR. Meenakshi*: On Almost Positive definite matrices, Tech. Report # 8616, Indian Statistical Institute, New Delhi, August 1986.
- [9] *S. K. Mitra and M. L. Puri*: The Fundamental Bordered Matrix of Linear Estimation and the Duffin-Morley General Linear Electromechanical Systems, Applicable Analysis, 14 (1983) 241—258.
- [10] *S. K. Mitra and P. L. Odell*: On parallel Summability of matrices, J. Linear Algebra. Appl. 74 (1986) 239—255.

*Author's address*: Department of Mathematics, Annamalai University, Annamalainagar, 608 002, India.