

Josef Daneš

On norms and subsets of linear spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 12 (1971), No. 4, 835--844

Persistent URL: <http://dml.cz/dmlcz/105387>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1971

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON NORMS AND SUBSETS OF LINEAR SPACES

Josef DANĚŠ, Praha

J. Zemánek has given [10] an example of a non-empty finitely open and nowhere dense convex subset of a normed linear space. Some general theorems concerning the existence of comparable non-equivalent norms in infinite-dimensional spaces give a possibility to construct simpler examples of that type (see Proposition 1 and Examples 1 - 3 below).

Throughout this paper, X denotes a real linear space. Let G be a subset of X . G is said to be: (1) finitely open (see [6], Definition 1.10.2) if each finite-dimensional affine subspace L of X intersects G in a set open in L (in the unique linear topology on L), (2) linearly bounded if its intersection with any line is bounded (as a subset of the line). The convex hull of G is denoted by $\text{conv } G$, $\text{diam}_{\|\cdot\|} G$ denotes the diameter of G in $(X, \|\cdot\|)$, where $\|\cdot\|$ is a norm on X , " $\xrightarrow{\|\cdot\|}$ " denotes the convergence in the topology given by $\|\cdot\|$. G is said to be $\|\cdot\|$ - P if G is P in $(X, \|\cdot\|)$ where P is a property of subsets of X (we shall use P = weak, bounded, open). G is a convex body if it is convex and has a

non-empty interior in $(X, \|\cdot\|)$.

We begin with

Proposition 1. Let $\|\cdot\|_0$ and $\|\cdot\|_1$ be two non-equivalent norms on a linear space X such that $\|\cdot\|_0 \leq K \|\cdot\|_1$ (for some $K > 0$). Then $C = \{x \in X : \|x\|_1 < 1\}$ is a finitely open nowhere dense absolutely convex (non-empty) subset of $(X, \|\cdot\|_0)$. Clearly, X must be infinite-dimensional.

Proof. Clearly, C is absolutely convex and non-empty. Since C is open in $(X, \|\cdot\|_1)$ it is finitely open. Let C_0 denote the closure of C in $(X, \|\cdot\|_0)$. For each $y \in C_0$ there is $x_y \in C$ such that $\|y - x_y\|_0 < 1$. Then $\|y\|_0 \leq \|y - x_y\|_0 + \|x_y\|_0 \leq \|y - x_y\|_0 + K \|x_y\|_1 < 1 + K$. Hence $C_0 \subset (K+1)C$. Suppose that C_0 has a non-empty interior in $(X, \|\cdot\|_0)$. Then the absolute convexity of C_0 implies the existence of some $\epsilon > 0$ such that $\{x \in X : \|x\|_0 < \epsilon\} \subset C_0$. This and $C_0 \subset (K+1)C$ imply that $\|\cdot\|_1 \leq \epsilon^{-1}(K+1)\|\cdot\|_0$, a contradiction to the non-equivalence of both norms.

Proposition 2. Let $\|\cdot\|_0$ and $\|\cdot\|_1$ be two norms on a linear space X such that $\|\cdot\|_0 \leq K \|\cdot\|_1$ ($K > 0$). Define $\|\cdot\|_t = (1-t)\|\cdot\|_0 + t\|\cdot\|_1$ for $0 \leq t \leq 1$. Then $1^\circ \|\cdot\|_t, t \in [0, 1]$ are the norms on X , $2^\circ \|\cdot\|_{t_1} \leq K(t_1, t_2)\|\cdot\|_{t_2}$ for $0 \leq t_1 \leq t_2 \leq 1$, where $K(t_1, t_2) = [t_1 + K(1-t_1)][t_2 + K(1-t_2)]^{-1}$, $3^\circ \|\cdot\|_{t_2} \leq t_2 t_1^{-1} \|\cdot\|_{t_1}$ for $0 < t_1 \leq t_2 \leq 1$, and hence the norms $\|\cdot\|_{t_1}$ and $\|\cdot\|_{t_2}$ are equivalent, 4° if the norms $\|\cdot\|_0$ and $\|\cdot\|_1$ are non-equivalent, then $\|\cdot\|_0$ and $\|\cdot\|_t, t \in (0, 1]$, are non-equivalent.

The proof goes by a direct computation.

Proposition 2 says that two comparable norms can be joined by a "continuum" of pairwise equivalent norms.

The following two theorems were first proved in our thesis [3] and published without proof in [4].

Theorem 1. Let $(X, \|\cdot\|)$ be a normed linear space such that its dual space X^* is separable. Then there exists a norm $\|\cdot\|_w$ on X such that the $\|\cdot\|$ -weak topology and the $\|\cdot\|_w$ -topology coincide on the $\|\cdot\|$ -bounded subsets of X , and $\|\cdot\|_w \leq \|\cdot\|$. If X has an infinite dimension, then the norms $\|\cdot\|_w$ and $\|\cdot\|$ are non-equivalent.

Proof. Let $\{\mu_n\}$ be a dense sequence in the unit ball of X^* and $\|x\|_w = \sum_{n=1}^{\infty} 2^{-n} |\mu_n(x)|$ for x in X . It is easy to see that $\|\cdot\|_w$ is a norm and $\|\cdot\|_w \leq \|\cdot\|$. Let M be a $\|\cdot\|$ -bounded subset of X , x_0 a point of M . If W is a weak neighbourhood of x_0 in M then there exist $\epsilon > 0$ and $f_1, \dots, f_m \in X^*$, $\|f_j\| = 1$ ($j = 1, \dots, m$) such that $W_1 = \{x \in M : |f_j(x - x_0)| < \epsilon \text{ for } j = 1, \dots, m\} \subset W$. Clearly, W_1 is a weak neighbourhood of x_0 in M . Without loss of generality we may suppose that M contains at least two points. There are integers n_1, \dots, n_m such that


$$\|\mu_{n_j} - f_j\| < \epsilon (4 \operatorname{diam}_{\|\cdot\|} M)^{-1} \text{ for } j = 1, \dots, m. \text{ Let } N = 1 + \max\{n_1, \dots, n_m\} \text{ and } V = \{x \in M : \|x - x_0\|_w < \epsilon 2^{-N}\}.$$

We shall show that $W_1 \supset V$. Let $x \in V$. Then

$$2^{-n_j} |(\mu_{n_j} - f_j)(x - x_0) + f_j(x - x_0)| \leq \|x - x_0\|_w < \epsilon 2^{-N} \leq \frac{\epsilon}{2} 2^{-n_j}$$

for $j = 1, \dots, m$. Since $|(\mu_{n_j} - f_j)(x - x_0)| \leq \|\mu_{n_j} - f_j\| \|x - x_0\| < \epsilon/4$, there is $|f_j(x - x_0)| < \frac{\epsilon}{2} + \frac{\epsilon}{4} < \epsilon$ for $j = 1, \dots, m$. Hence $x \in W_1$ and $V \subset W_1 \subset W$. Conversely, let $V = \{x \in M :$

$\{x \in M : \|x - x_0\|_{w'} < \varepsilon\}$ ($\varepsilon > 0$) be a $\|\cdot\|_{w'}$ -neighbourhood of x_0 in M . A direct calculation shows that V contains $W = \{x \in M : \sum_{n=1}^m 2^{-n} |\mu_n(x - x_0)| < \varepsilon/2\}$ where m is so large that $\sum_{n=m+1}^{\infty} 2^{-n+1} \text{diam } M < \varepsilon$. Clearly, W is a $\|\cdot\|$ -weak neighbourhood of x_0 in M . Suppose that X is infinite-dimensional and the norms $\|\cdot\|_{w'}$ and $\|\cdot\|$ are equivalent. Let us denote $X^* = (X, \|\cdot\|)$ and $B = \{x \in X : \|x\| \leq 1\}$. Then



is a commutative diagram of topological spaces and continuous mappings; (B, τ) denotes the set B with the topology induced by the τ -topology of X . Thus, the three topologies $\|\cdot\|$, $\|\cdot\|_{w'}$, and $\sigma(X, X^*)$ coincide on B , a contradiction to the infinite dimensionality of X (see [5], Chapt.V, Exerc. 7.9). Hence the norms $\|\cdot\|_{w'}$ and $\|\cdot\|$ are non-equivalent. The proof is complete.

Theorem 2. Let $(X, \|\cdot\|)$ be a separable normed linear space. Then there exists a norm $\|\cdot\|_{w'}$ on X such that the $\|\cdot\|$ -weak topology is on $\|\cdot\|$ -bounded subsets of X stronger than the $\|\cdot\|_{w'}$ -topology, and $\|\cdot\|_{w'} \leq \|\cdot\|$. If X has infinite dimension, then the norms $\|\cdot\|_{w'}$ and $\|\cdot\|$ are non-equivalent.

Proof. By [1], Chapt.III, Theorem 9.16 the unit ball of X^* contains a sequentially $\sigma(X^*, X)$ -dense sequence

$(X^* = (X, \|\cdot\|)^*)$; let $S = \{\mu_n\}$ be such a sequence, and set $\|x\|_{uv} = \sum_{n=1}^{\infty} 2^{-n} |\mu_n(x)|$ for x in X . Let $0 \neq x \in X$. Then there is $f \in X^*$ such that $|f(x)| = \varepsilon > 0$. Since $RS = \{\kappa \mu_m : \kappa \in \mathbb{R}, m = 1, 2, \dots\}$ is $\sigma(X^*, X)$ -dense in X^* , there exist $\kappa \in \mathbb{R}$ and μ_m such that $\kappa \mu_m$ lies in the $\sigma(X^*, X)$ -neighbourhood $\{x^* \in X^* : |(x^* - f)(x)| < \varepsilon\}$ of f . Then $|\kappa \mu_m(x)| \geq |f(x)| - |\kappa \mu_m - f(x)| > 0$. Hence $\|x\|_{uv} > 0$, and $\|\cdot\|_{uv}$ is a norm on X . The proof of the other assertions of the theorem is the same as that of the corresponding assertions of Theorem 1.

Theorem 3 below is the precise statement of the results of the proof of Proposition 1.1 in [2]. That proof relies on a paper of V. Klee [7]. We repeat their proof making use of Theorem 2 instead of [7].

Theorem 3. Let $(X, \|\cdot\|)$ be an infinite-dimensional normed linear space. Then there are two norms $|\cdot|$ and $\|\cdot\|$ on X such that $|\cdot| \leq \|\cdot\| \leq \|\cdot\|$ and none of them is equivalent to $\|\cdot\|$. If $\|\cdot\|$ is complete (that is, $(X, \|\cdot\|)$ is complete), the norms $|\cdot|$ and $\|\cdot\|$ are not.

Proof. Let B be a Hamel basis for X such that $\|b\| \leq 1$ for all $b \in B$ and $\inf\{\|b\| : b \in B\} = 0$. It is easy to verify that $\|\cdot\|$ defined as the Minkowski functional of the absolutely convex hull of B , satisfies our requirements.

Let L be a separable infinite-dimensional subspace of $(X, \|\cdot\|)$, $\|\cdot\|_{uv}$ the norm of Theorem 2 corresponding

to $(L, \|\cdot\|)$, and $V = \{x \in L: \|x\|_{uv} \leq 1\}$. By Theorem 2, the norms $\|\cdot\|_{uv}$ and $\|\cdot\|$ on L are non-equivalent and $\|\cdot\|_{uv} \leq \|\cdot\|$. This implies that the set V is unbounded in $(L, \|\cdot\|)$; V is linearly bounded since it is bounded in $(L, \|\cdot\|_{uv})$. Hence V is an absolutely convex, linearly bounded, unbounded closed body in $(L, \|\cdot\|)$. Let $U = \{x \in X: \|x\| \leq 1\}$. Then $C = \text{con}(U \cup V)$ is an absolutely convex body in $(X, \|\cdot\|)$. Suppose that C is not linearly bounded. Then C contains a line J through 0 . Let $x \in J$. For each integer n , $nx \in J$ and hence $nx = \lambda_n x_n + (1 - \lambda_n) y_n$ for some $\lambda_n \in [0, 1]$, $x_n \in U$, $y_n \in V$. Since $n^{-1} \lambda_n x_n \xrightarrow{\|\cdot\|} 0$, we have $V \ni n^{-1} (1 - \lambda_n) y_n \xrightarrow{\|\cdot\|} x$. V is $\|\cdot\|$ -closed and hence $x \in V$. This implies that $J \subset V$, a contradiction to the linear boundedness of V . We have proved that the set C must be linearly bounded. Hence its Minkowski functional $|\cdot|$ defines a norm for X . The inclusion $U \subset C$ implies $|\cdot| \leq \|\cdot\|$. Since C is unbounded in $(X, \|\cdot\|)$, the norms $|\cdot|$ and $\|\cdot\|$ are non-equivalent.

The part of the theorem concerning the completeness follows from the open mapping theorem.

Theorem 4. Let X be an infinite-dimensional linear space and C a non-empty absolutely convex, linearly bounded, finitely open subset of X . Then there are two norms $|\cdot|$ and $\|\cdot\|$ on X such that C is open in $(X, \|\cdot\|)$ and nowhere dense in $(X, |\cdot|)$ and $|\cdot| \leq \|\cdot\|$.

Proof. Let $\|\cdot\|$ be the Minkowski functional of C . It is a norm on X . It is sufficient to use Theorem 3 and

then Proposition 1.

Theorem 5. Let $(X, \|\cdot\|)$ be a normed linear space of infinite dimension. Then there is a non-empty absolutely convex finitely open bounded and nowhere dense subset C of $(X, \|\cdot\|)$.

Proof. Let $\|\cdot\|$ be as in Theorem 3. It is sufficient to set $C = \{x \in X : \|x\| < 1\}$ and apply Proposition 1.

Corollary. Let X be an infinite-dimensional linear space. Then:

1. there is neither a minimal nor maximal norm on X (a norm $\|\cdot\|$ on X is said to be minimal [maximal] if for any norm $\|\cdot\|$ on X there exists $K > 0$ such that $\|\cdot\| \leq K \|\cdot\|$ [$K \|\cdot\| \leq \|\cdot\|$]);

2. the strongest locally convex topology on X is not normable;

3. if (X, τ) is a locally convex space of minimal type (see [9], Chapt. IV, Exerc. 6), it is non-normable.

Remark. Any finitely open convex subset of X is open in the strongest locally convex topology on X . Hence there is no finitely open non-empty convex subset of X which is nowhere dense in the strongest locally convex topology. The second part of our corollary is not the best possible result; see [9], Chapt. II, Exerc. 7.

Examples. 1. Let G be a compact subset of \mathbb{R}^n ($n \geq 1$) with a positive Lebesgue measure, $mes G > 0$, X the linear space of all continuous real-valued functions on G , $\|\cdot\|$ the sup norm on X , $|\cdot| = \|\cdot\|_{L_p(G)}$ ($p \geq 1$). Then

$|\cdot| \leq \|\cdot\|$ and these norms are non-equivalent on X .
 (Hint: For any $\varepsilon > 0$, there exist disjoint closed sub-
 sets $M_\varepsilon, N_\varepsilon$ of G such that $0 < \text{mes } M_\varepsilon < \varepsilon$,
 $\text{mes } N_\varepsilon > \text{mes } G - 2\varepsilon$. Let $\mu_\varepsilon \in X$ be such that
 $\mu_\varepsilon|_{M_\varepsilon} = (2\varepsilon)^{-1/p}$, $\mu_\varepsilon|_{N_\varepsilon} = 0$, $0 \leq \mu_\varepsilon \leq (2\varepsilon)^{-1/p}$.
 Then $\|\mu_\varepsilon\| = (2\varepsilon)^{-1/p}$, $|\mu_\varepsilon| = (\int_{G \setminus N_\varepsilon} |\mu_\varepsilon(x)|^q dx)^{1/q} \leq$
 $\leq (2\varepsilon \cdot (2\varepsilon)^{-1})^{1/q} = 1$.

Another hint: If both norms are equivalent on $X = C(G)$, then
 $C(G)$ is a closed and dense subspace of $L_p(G)$. This
 leads to a contradiction.) By Proposition 1, $C = \{x \in X :$
 $\|x\| < 1\}$ is a finitely open, absolutely convex, nowhe-
 re dense, bounded (non-empty) subset of $(X, |\cdot|)$.

2. Let G be as above and $1 \leq p'' < p < p' \leq \infty$.
 Set
 $X = L_p(G)$, $\|\cdot\| = \|\cdot\|_{L_p(G)}$, $\|\cdot\| = (\text{mes } G)^{1/p - 1/p'} \|\cdot\|_{L_{p'}(G)}$,
 and $|\cdot| = (\text{mes } G)^{1/p - 1/p''} \|\cdot\|_{L_{p''}(G)}$. Then $|\cdot| \leq \|\cdot\| \leq$
 $\leq \|\cdot\|$. Any two of these norms are non-equivalent on
 X . (Hint: By [8], § 12, Sect. 1, we may restrict oursel-
 ves to the easy case $G = [0, 1]$.)

3. Let
 $1 \leq p'' < p < p' \leq \infty$, $X = L_p$, $\|\cdot\| = \|\cdot\|_{L_p}$, $|\cdot| = \|\cdot\|_{L_{p'}}$,
 and $\|\cdot\| = \|\cdot\|_{L_{p''}}$. Then $|\cdot| \leq \|\cdot\| \leq \|\cdot\|$ and any two
 of these norms are non-equivalent.

Remark. Does Theorem 4 hold with "absolutely convex"
 replaced by "convex"? This leads to another question. Is
 the absolute convex hull of a convex linearly bounded finitely
 open set linearly bounded? We conjecture that the answer
 is (generally) no.

If $\|\cdot\|_0$ and $\|\cdot\|_1$ in Proposition 2 are non-equivalent, does there exist a "monotone continuum" of pairwise non-equivalent comparable norms? The answer is yes, when $\|\cdot\|_i$ ($i = 0, 1$) are the L_{p_i} -norms on $X = L_{p_1}$ ($p_0 < p_1$) or the l_{p_i} -norms on $X = l_{p_1}$ ($p_0 > p_1$).

R e f e r e n c e s

- [1] ALEXIEWICZ A.: Functional Analysis (in Polish), PWN, Warsaw, 1969.
- [2] BESSAGA C. and KLEE V.: Two topological properties of linear spaces, Israel J. Math. 2(1964), 211-220.
- [3] DANÉŠ J.: Nonlinear operators and functionals, Thesis, Charles University, Prague, 1968 (in Czech).
- [4] DANÉŠ J.: Continuity properties of nonlinear mappings, Comment. Math. Univ. Carolinae 9(1968), 353-364.
- [5] DUNFORD N., SCHWARTZ J.T.: Linear Operators. Vol. I., Interscience Publ., New York, 1958.
- [6] HILLE E., PHILLIPS R.S.: Functional Analysis and Semigroups, Amer. Math. Soc. Colloquium Publ., vol. XXXI, Providence, R.I., 1957.
- [7] KLEE V.: A note on topological properties of normed linear spaces, Proc. Amer. Math. Soc. 7(1956), 735-737.
- [8] KRASNOSELSKII M.A., RUTICKII Ia.B.: Convex Functions and Orlicz's Spaces (in Russian), Gostehizdat, Moscow, 1956.
- [9] SCHAEFER H.H.: Topological Vector Spaces, The Macmillan Comp., New York, 1966.

[10] ZEMÁNEK J.: Nowhere dense set which is finitely open,
Comment.Math.Univ.Carolinae 11(1970),83-89.

Matematický ústav
Karlova universita
Sokolovská 83
Praha 8, Československo

(Oblatum 7.6.1971)