

Robert Frič; Darrell C. Kent

Completion of sequential Cauchy spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 18 (1977), No. 2, 351--361

Persistent URL: <http://dml.cz/dmlcz/105779>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1977

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

COMPLETION OF SEQUENTIAL CAUCHY SPACES

R. FRIČ, Žilina and D.C. KENT, Pullman

Abstract: We study two types of sequential Cauchy spaces projectively generated by classes of functions, their completions, and their mutual relations.

Key words: Sequential Cauchy space, completion, convergence space, sequential envelope.

AMS: 54D55

Ref. Ž.: 3.961.1

1. Introduction. For the reader's convenience we recall in this section some basics about (sequential) Cauchy spaces.

Notation 1.1. If $\langle x_n \rangle, \langle y_n \rangle$ are two sequences, then $\langle x_n \rangle \wedge \langle y_n \rangle$ denotes a sequence $\langle z_n \rangle$ defined as follows: $z_1 = x_1, z_2 = y_1, z_3 = x_2, z_4 = y_2, \dots$, i.e. $x_n = z_{2n-1}, y_n = z_{2n}$.

Definition 1.2. A Cauchy space is a pair (X, L) , where X is a set and L a collection of sequences ranging in X such that

- (1) $\langle x \rangle \in L$ for each $x \in X$;
- (2) $\langle x_n \rangle \in L$ implies $\langle x'_n \rangle \in L$ for each subsequence $\langle x'_n \rangle$ of $\langle x_n \rangle$;
- (3) if $\langle x_n \rangle, \langle y_n \rangle \in L$ and there are subsequences $\langle x'_n \rangle$ of $\langle x_n \rangle$ and $\langle y'_n \rangle$ of $\langle y_n \rangle$ such that $x'_n = y'_n, n \in \mathbb{N}$, then $\langle x_n \rangle \wedge \langle y_n \rangle \in L$; and

(4) if $\langle x_n \rangle \wedge \langle x \rangle \in L$ and $\langle x_n \rangle \wedge \langle y \rangle \in L$, then $x = y$.

If (X, L) is a Cauchy space, then L is called a Cauchy structure for X . If L satisfies the additional condition

(5) $\langle x_n \rangle \in L$ whenever

(a) each subsequence $\langle x'_n \rangle$ of $\langle x_n \rangle$ contains a subsequence $\langle x''_n \rangle$ of $\langle x'_n \rangle$ such that $\langle x''_n \rangle \in L$; and

(b) if $\langle x'_n \rangle$ and $\langle x''_n \rangle$ are subsequences of $\langle x_n \rangle$ such that $\langle x'_n \rangle, \langle x''_n \rangle \in L$, then $\langle x'_n \rangle \wedge \langle x''_n \rangle \in L$;

then (X, L) is said to be a *Cauchy space.

The effect of condition (5) can be brought out by considering the real line with its usual metric. Every bounded sequence of real numbers has a Cauchy subsequence. Hence, every bounded sequence of real numbers satisfies condition (a). Yet every bounded sequence of real numbers is not Cauchy in the usual sense because (b) is lacking; e.g. consider the sequence 0, 1, 0, 1, 0, 1,

A Cauchy space (X, L) induces a convergence space $(X, \mathcal{L}, \lambda)$ in the following natural way: $x = \mathcal{L}\text{-lim } x_n$ iff $\langle x_n \rangle \wedge \langle x \rangle \in L$. Moreover, if (X, L) is a *Cauchy space, then $\mathcal{L} = \mathcal{L}^*$. The topological modification λ^{ω_1} of λ will be called a topological closure for X . A subspace Y of X is topologically dense in X if $\lambda^{\omega_1} Y = X$. A Cauchy space is said to be complete if each Cauchy sequence converges in the induced convergence space. A mapping $f: (X_1, L_1) \rightarrow (X_2, L_2)$ is said to be Cauchy-continuous if $\langle x_n \rangle \in L_1$ implies $\langle f(x_n) \rangle \in L_2$. The set of all Cauchy-continuous functions on (X, L) is denoted by $\hat{C}(X, L)$. The set

$M = \{ \langle f_m \rangle \in (\hat{C}(X, L))^N; \lim_{m, n \rightarrow \infty} f_m(x_n) \text{ exists for each } \langle x_n \rangle \in L \}$
is a Cauchy structure for $\hat{C}(X, L)$ and is said to be the continuous Cauchy structure. The space $(\hat{C}(\hat{C}(X, L), M), M)$ is denoted by $(\hat{C}^2(X, L), M^2)$. The evaluation mapping $ev_x: (X, L) \rightarrow (\hat{C}^2(X, L), M^2)$ is defined by $ev_x(x) = \Phi_x$, where for $f \in \hat{C}(X, L)$ we define $\Phi_x(f) = f(x)$; it is always Cauchy-continuous. If it is a Cauchy-embedding (i.e. a Cauchy-homeomorphism into), then (X, L) is said to be \hat{C} -embedded.

2. Projective generations of Cauchy structures.

Proposition and definition 2.1. Let (X, L) be a Cauchy space and $D \subset \hat{C}(X, L)$, D separates points of X . Let $L_D = \{ \langle x_n \rangle \in X^N; \lim f(x_n) \text{ exists whenever } f \in D \}$ and $L_d = \{ \langle x_n \rangle \in X^N; \lim_{m, n \rightarrow \infty} f_m(x_n) \text{ exists whenever } \langle f_m \rangle, f_m \in D \text{ is a Cauchy sequence in } (\hat{C}(X, L), M) \}$. Then L_D and L_d are $*$ Cauchy structures for X and $L \subset L_d \subset L_D$. If $L = L_D$, then L , resp. (X, L) , is said to be projectively generated by D. If $L = L_d$, then L , resp. (X, L) , is said to be c-projectively generated by D.

It follows immediately that if a space is projectively generated by D , then it is also c-projectively generated by D . The converse statement is not true in general as it will be shown by a counterexample (see Proposition 4.7). In

[I - K] it was proved that for $D = C(X, L)$ the following are equivalent:

- (a) (X, L) is \hat{C} -embedded; (b) $L = L_D$; (c) $L = L_d$ (the original notation is $L_D = L_P, L_d = L_M$).

3. d-completion.

Definition 3.1. Let (X,L) be a Cauchy space c -projectively generated by $D \subset \hat{C}(X,L)$. A complete Cauchy space (X_1, L_1) is said to be a d-completion of (X,L) if

- (a) (X,L) is a topologically dense subspace of (X_1, L_1) ;
- (b) for each $f \in D$ there is $\bar{f} \in \hat{C}(X_1, L_1)$ such that $f = \bar{f} \upharpoonright X$, i.e. $D \subset \hat{C}(X_1, L_1) \upharpoonright X$;
- (c) (X_1, L_1) is c -projectively generated by $\bar{D} = \{\bar{f} \in \hat{C}(X_1, L_1); \bar{f} \upharpoonright X \in D\}$; and
- (d) \bar{D} and D endowed with the corresponding continuous Cauchy structures are Cauchy-homeomorphic under the natural correspondence, i.e. the correspondence $\bar{f} \rightarrow \bar{f} \upharpoonright X = f$ is one-to-one and $\langle \bar{f}_n \rangle, \bar{f}_n \in \bar{D}$, is a Cauchy sequence in $(\hat{C}(X_1, L_1), M)$ iff $\langle f_n \rangle, f_n = \bar{f}_n \upharpoonright X$, is a Cauchy sequence in $(\hat{C}(X,L), M)$.


Lemma 3.2. Let (X,L) be a Cauchy space c -projectively generated by $D \subset \hat{C}(X,L)$, $(D, M \upharpoonright D)$ the subspace of $(\hat{C}(X,L), M)$, and e a mapping of (X,L) into $(\hat{C}(D, M \upharpoonright D), M)$ defined as follows: $e(x) = \Phi_x$, where for $f \in D$ we define $\Phi_x(f) = f(x)$. Then e is a Cauchy embedding.

Lemma 3.2 was proved in [I - K] in the special case of $D = \hat{C}(X,L)$. The proof of the general case is similar.

Theorem 3.3. Let (X,L) be a Cauchy space c -projectively generated by $D \subset \hat{C}(X,L)$. Then there exists a d -completion of (X,L) .

Proof. It follows from Lemma 3.2 that identifying x with $e(x)$ we can consider (X,L) as a subspace of $(\hat{C}(D, M \upharpoonright D), M)$. We shall prove that the subspace (X_1, L_1) of $(\hat{C}(D, M \upharpoonright D), M)$, where

X_1 is the topological closure of X in $(\hat{C}(D, M | D), M)$ and $L_1 = M | X_1$, is a d -completion of (X, L) . It was proved in [I - K] that $(\hat{C}(D, M | D), M)$ is a complete space. Thus the closed subspace (X_1, L_1) of $(\hat{C}(D, M | D), M)$ is complete. We are to prove that (X_1, L_1) satisfies conditions (a) - (d) of Definition 3.1. Condition (a) follows from the construction of (X_1, L_1) . It was proved in [F] that the space $(\hat{C}(X, L), M)$ is \hat{C} -embedded. Thus the subspace $(D, M | D)$ is also \hat{C} -embedded, and hence the evaluation mapping $ev_D: (D, M | D) \rightarrow (\hat{C}^2(D, M | D), M^2)$ is a Cauchy embedding. Consequently, for each $f \in D$ the image $ev_D(f) = \hat{f}$ is a Cauchy-continuous function on $(\hat{C}(D, M | D), M)$. Since $\hat{f}(\Phi) = \Phi(f)$ for each $\Phi \in \hat{C}(D, M | D)$, we have $\hat{f}(x) = f(x)$ for each $\Phi_x = x \in X$. Hence $\bar{f} = \hat{f} | X_1$ is a Cauchy-continuous extension of f onto (X_1, L_1) and condition (b) is satisfied. The construction of \bar{f} is shown on the following diagram:



Now, we shall prove condition (d). It follows by a standard topological argument that the extension \bar{f} of f is uniquely determined. Hence the natural correspondence $\bar{f} \rightarrow \bar{f} | X = f$ is one-to-one. Clearly, if $\langle \bar{f}_n \rangle$, $\bar{f}_n | X \in D$, is a Cauchy sequence in $(\hat{C}(X_1, L_1), M)$, then $\langle f_n \rangle$, $f_n = \bar{f}_n | X$, is a Cauchy sequence in $(\hat{C}(X, L), M)$. Conversely, let $\langle f_n \rangle$ be a Cauchy sequence in $(D, M | D)$. Since ev_D is a Cauchy embedding, the se-

quence $\langle \hat{f}_n \rangle$, $\hat{f}_n = \text{ev}_D(f_n)$, is a Cauchy sequence in $(\hat{C}^2(D, M | D), M^2)$. Hence $\langle \bar{f}_n \rangle$, $\bar{f}_n | X = f_n$, is a Cauchy sequence in $(\hat{C}(X_1, L_1), M)$.

It remains to prove condition (c). Let $\langle \Phi_n \rangle$ be a sequence in $X_1 \subset \hat{C}(D, M | D)$ such that

(1) $\lim_{m, n \rightarrow \infty} \bar{f}_m(\Phi_n)$ exists whenever $\langle \bar{f}_m \rangle$, $\bar{f}_m \in \bar{D}$, is a Cauchy sequence in $(\hat{C}(X_1, L_1), M)$.

Since $\bar{f}_m(\Phi_n) = \Phi_n(f_m)$, $f_m = \bar{f}_m | X$, it follows from (d) that

(1) is equivalent to

(2) $\lim_{m, n \rightarrow \infty} \Phi_n(f_m)$ exists whenever $\langle f_m \rangle$ is a Cauchy sequence in $(D, M | D)$.

Thus $\langle \Phi_n \rangle \in L_1$ and the proof is complete.

Theorem 3.4. Let (X, L) be a Cauchy space c -projectively generated by $D \subset \hat{C}(X, L)$. If (X_1, L_1) and (X_2, L_2) are two d -completions of (X, L) , then there is a Cauchy homeomorphism $h: (X_1, L_1) \rightarrow (X_2, L_2)$ such that $h(x) = x$ for each $x \in X$.

Proof. For $i = 1, 2$, denote by $D_i = \{f \in \hat{C}(X_i, L_i); f | X \in D\}$, by $(D_i, M | D_i)$ the subspace of $(\hat{C}(X_i, L_i), M)$, and by $(D, M | D)$ the subspace of $(\hat{C}(X, L), M)$. It follows from (d) in Definition 3.2 that $(D_i, M | D_i)$ and $(D, M | D)$ are Cauchy-homeomorphic under the natural correspondence. Consequently, $\varphi: (D_2, M | D_2) \rightarrow (D_1, M | D_1)$, where for $f \in D_2$ its image $\varphi(f)$ is determined by $\varphi(f) | X = f | X$, and hence also its first conjugate $\varphi^*: (\hat{C}(D_1, M | D_1), M) \rightarrow (\hat{C}(D_2, M | D_2), M)$, $\varphi^*(\Phi) = \varphi \circ \Phi$, are Cauchy homeomorphisms. It follows from Lemma 3.2 that identifying x with $e_1(x)$, where for $f \in D_1$ we define $(e_1(x))(f) = f(x)$, we can consider the complete space (X_1, L_1) as a closed subspace of $(\hat{C}(D_1, M | D_1), M)$.

Now, an easy computation shows that for each $x \in X$ we have $\varphi^*(x) = x$.

$$\begin{array}{ccccc}
 X_1 & \xleftarrow{id} & X & \xrightarrow{\quad} & X_2 \\
 \downarrow id = e & & & & \downarrow id = e_2 \\
 C(D_1, M|D_1) & \xrightarrow{\quad} & & \xrightarrow{\varphi^*} & C(D_2, M|D_2)
 \end{array}$$

Since X is topologically dense in (X_1, L_1) , it follows by a standard topological argument that $h = \varphi^*|X_1$ is the desired Cauchy homeomorphism.

4. D-completion.

Definition 4.1. Let (X, L) be a Cauchy space projectively generated by $D \subset \hat{C}(X, L)$. A complete Cauchy space (X_1, L_1) is said to be a D-completion of (X, L) if

- (a) (X, L) is a topologically dense subspace of (X_1, L_1) ;
- (b) for each $f \in D$ there is $\bar{f} \in \hat{C}(X_1, L_1)$ such that $f = \bar{f}|X$, i.e. $D \subset \hat{C}(X_1, L_1)|X$; and
- (c) (X_1, L_1) is projectively generated by $D = \{\bar{f} \in \hat{C}(X_1, L_1); \bar{f}|X \in D\}$.

Proposition 4.2. Let (X, L) be a Cauchy space projectively generated by $D \subset \hat{C}(X, L)$ and $(X, \mathcal{L}^*, \lambda)$ the associated convergence space. Then:

- (a) $D \subset C(X)$ and $(X, \mathcal{L}^*, \lambda)$ is D-sequentially regular.
- (b) L is the set of all D-fundamental sequences in $(X, \mathcal{L}^*, \lambda)$.
- (c) $(X, \mathcal{L}^*, \lambda)$ is D-sequentially complete iff (X, L)

is complete.

The straightforward proof is omitted.

Proposition 4.3. Let $(X, \mathcal{L}^*, \lambda)$ be a D-sequentially regular convergence space and L the set of all D-fundamental sequences. Then:

- (a) L is a $*$ Cauchy structure for X.
- (b) $D \subset \hat{C}(X, L)$ and (X, L) is projectively generated by D.
- (c) $(X, \mathcal{L}^*, \lambda)$ is associated with (X, L) .
- (d) (X, L) is complete iff $(X, \mathcal{L}^*, \lambda)$ is D-sequentially complete.

The straightforward proof is omitted.

Theorem 4.4. Let (X, L) be a Cauchy space projectively generated by $D \subset \hat{C}(X, L)$. Then there exists a D-completion of (X, L) .

Proof. Let $(X, \mathcal{L}, \lambda)$ be the convergence space associated with (X, L) . It follows from (a) in Proposition 4.2 that $(X, \mathcal{L}, \lambda)$ is D-sequentially regular. Let $(X_1, \mathcal{L}_1, \lambda_1)$ be a D-sequential envelope of $(X, \mathcal{L}, \lambda)$, $\bar{D} = \{\bar{f} \in \mathcal{C}(X_1); \bar{f} \upharpoonright X \in D\}$, and L_1 the set of all \bar{D} -fundamental sequences in X_1 . It follows from Proposition 4.2 and Proposition 4.3 that (X_1, L_1) is a D-completion of (X, L) .

Note 4.5. Let $(X, \mathcal{L}^*, \lambda)$ be a D-sequentially regular convergence space. Let L be the set of all D-fundamental sequences in X. It follows from Proposition 4.3 that (X, L) is a $*$ Cauchy space projectively generated by $D \subset \hat{C}(X, L)$. Let (X_1, L_1) be a D-completion of (X, L) . Using Proposition 4.2 and Proposition 4.3 it is easy to see that the convergence space $(X_1, \mathcal{L}_1, \lambda_1)$ associated with (X_1, L_1) is a D-sequential enve-

lopes of $(X, \mathcal{C}^*, \lambda)$.

Theorem 4.6. Let (X, L) be a Cauchy space projectively generated by $D \subset \hat{C}(X, L)$. If (X_1, L_1) and (X_2, L_2) are two D-completions of (X, L) , then there is a Cauchy homeomorphism $h: (X_1, L_1) \rightarrow (X_2, L_2)$ such that for each $x \in X$ we have $h(x) = x$.

Proof. Let $(X, \mathcal{C}, \lambda)$ be the convergence space associated with (X, L) and $(X_i, \mathcal{C}_i, \lambda_i)$ the convergence space associated with (X_i, L_i) , $i = 1, 2$. It follows from Note 4.5 that $(X_i, \mathcal{C}_i, \lambda_i)$ is a D-sequential envelope of $(X, \mathcal{C}, \lambda)$. Hence there is a homeomorphism $h: (X_1, \mathcal{C}_1, \lambda_1) \rightarrow (X_2, \mathcal{C}_2, \lambda_2)$ such that for each $x \in X$ we have $h(x) = x$ (cf. Theorem 5 in [N]). Since (X_i, L_i) are complete space, $h: (X_1, L_1) \rightarrow (X_2, L_2)$ is a Cauchy homeomorphism.

5. Example.

Definition 5.1. Let $X \neq \emptyset$ and $\langle x_n \rangle, \langle y_n \rangle \in X^{\mathbb{N}}$. We say that $\langle y_n \rangle$ is derived from $\langle x_n \rangle$, in symbols $\langle y_n \rangle \rightarrow \langle x_n \rangle$, if $F(\langle y_n \rangle) \supset F(\langle x_n \rangle)$, where $F(\langle x_n \rangle)$ denotes the filter generated by sections of a sequence $\langle x_n \rangle$.

Example 5.2. Let $X_2 = (\bigcup_{m \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} (x_{mn})) \cup (\bigcup_{m \in \mathbb{N}} (x_m)) \cup (x_0)$. Let $\alpha \in \mathbb{N}^{\mathbb{N}}$, $m_0 \in \mathbb{N}$, $A \subset (\bigcup_{m \in \mathbb{N}} (x_{m\alpha(m)}))$, and $f \in \{0, 1\}^{X_2}$ a function on X_2 defined as follows:

$$f(x) = 1 \text{ for } x \in (A \cup (\bigcup_{n \in \mathbb{N}} (x_{m_0 n})) \cup (x_{m_0})),$$

$$f(x) = 0 \text{ otherwise.}$$

Let \bar{D} be the set of all such functions and (X_2, L_2) the Cauchy space projectively generated by \bar{D} . Let $X = \bigcup_{m \in \mathbb{N}} \bigcup_{n \in \mathbb{N}} (x_{mn})$,

$X_1 = X \cup \left(\bigcup_{m \in \mathbb{N}} \langle x_m \rangle \right)$, $L = L_2 | X$, $L_1 = \{ \langle z_n \rangle \in X_1^{\mathbb{N}}; \langle z_n \rangle \rightarrow \langle x \rangle, x \in X_1, \text{ or } \langle z_n \rangle \rightarrow (\langle x_{mn} \rangle \wedge \langle x_m \rangle), m \in \mathbb{N} \}$, and $D = \bar{D} | X$.

Since (X, L) is clearly projectively generated by D it is also c -projectively generated by D and hence (X, L) possesses both a D -completion and a d -completion.

Proposition 5.3. For $\hat{D} = \bar{D} | X_1$ the space (X_1, L_1) is c -projectively generated by \hat{D} , but not projectively generated by \hat{D} .

Hint. $L_2 = \{ \langle z_n \rangle \in X_2^{\mathbb{N}}; \langle z_n \rangle \rightarrow \langle x \rangle, x \in X_2, \text{ or } \langle z_n \rangle \rightarrow (\langle x_{mn} \rangle \wedge \langle x_m \rangle), \text{ or } \langle z_n \rangle \rightarrow (\langle x_m \rangle \wedge \langle x_0 \rangle) \}$ and $\langle x_m \rangle \in (L_2 | X_1 - L_1)$.

Proposition 5.4. (X_1, L_1) is a d -completion of (X, L) .

Hint. $L = \{ \langle z_n \rangle \in X^{\mathbb{N}}; \langle z_n \rangle \rightarrow \langle x \rangle, x \in X, \text{ or } \langle z_n \rangle \rightarrow \langle x_{mn} \rangle, m \in \mathbb{N} \}$.

Proposition 5.5. (X_2, L_2) is a D -completion of (X, L) .

Proposition 5.6. \bar{D} and D endowed with the corresponding continuous Cauchy structures are not Cauchy-homeomorphic under the natural correspondence:

Proof. For otherwise (X_2, L_2) would be also a d -completion of (X, L) , which would imply the existence of a Cauchy homeomorphism $h: (X_1, L_1) \rightarrow (X_2, L_2)$ such that for each $x \in X$ we have $h(x) = x$.

Note 5.7. This shows that the condition (d) in Definition 3.1 is necessary and sufficient for the uniqueness of the d -completion up to a commuting Cauchy homeomorphism (cf. Theorem 3.4).

Note 5.8. Let (X, L) be a \hat{C} -embedded Cauchy space. Since for $D = \hat{C}(X, L)$ we have $L = L_D = L_D$, it follows immediately that a d -completion of (X, L) is also a D -completion of (X, L) . Consequently, the two completions are equivalent. It might be of some interest to characterize classes $D \subset \hat{C}$ for which the two completions are equivalent.

R e f e r e n c e s

- [I - K] J. IRWIN and D.C. KENT: Sequential Cauchy spaces, to appear.
- [F] R. FRIČ: On sequential Cauchy spaces, to appear.
- [N] J. NOVÁK: On sequential envelopes defined by means of certain classes of continuous functions, Czechoslovak Math. J. 18(93)(1968), 450-456.

Vysoká škola dopravná	Department of Pure and Applied
Katedra matematiky F SEF	Mathematics,
Marxa-Engelsa 25	Washington State University
010 88 Žilina	Pullman, Washington 99163
Československo	U. S. A.

(Obitum 16.12. 1976)