

Karel Čuda; Blanka Vojtášková  
Minimal monads

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 28 (1987), No. 4, 691--706

Persistent URL: <http://dml.cz/dmlcz/106583>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

MINIMAL MONADS

K. ČUDA, B. VOJTÁŠKOVÁ

**Abstract:** In the paper we present an analogue of some properties of classes of indiscernibles for minimal monads. Further we prove the existence of a minimal monad of a special character and using it we show six equivalents of the property of natural numbers "to be very far one from the other".

**Key words:** Alternative set theory, minimal monad, infinite natural number, Rudin-Keisler's ordering, definability with a parameter.

**Classification:** Primary 03E70  
Secondary 54J05

In this paper we are interested in a relation between relative definability and distance of natural numbers. This relationship is important especially when one supposes the existence of infinitely large definable natural numbers. For our investigations there are very convenient minimal monads which have some properties analogous to the properties of classes of indiscernibles.

We introduce and examine the notion  $\alpha \ll_{FN} \beta$  of great distance between  $\alpha, \beta$  which expresses the fact that  $\beta$  cannot be reached from  $\alpha$  by any definable function transforming FN into FN. Six equivalents of this feature are given. The most interesting are the following ones:

There are an endomorphic universe A with standard extension and  $\gamma \ll \beta$  such that  $\alpha, \gamma \in \text{Ex}_A(\text{FN})$  and  $\alpha \in \text{Ex}_{A(\gamma)}(\text{FN})$ .

There is an endomorphic universe A such that for each function  $f \in A$  we have

$$f''\text{FN} \subseteq \text{FN} \Rightarrow f(\alpha) < \beta.$$

Let us remind now several facts from [Č-K].

Let  $\mu_1, \mu_2$  be monads in  $\sum_{i \in \mathbb{N}} \mathcal{C}_i$  (i.e. classes of decomposition of V according to  $\sum_{i \in \mathbb{N}} \mathcal{C}_i$ ). We say that  $\mu_1 \ll_{\sum_{i \in \mathbb{N}} \mathcal{C}_i} \mu_2$  iff there is a function  $F \in \text{Sd}_{\sum_{i \in \mathbb{N}} \mathcal{C}_i}$

such that  $F^{\mu_2} = \mu_1$ .

The ordering  $\{ \mathcal{C} \}$  on monads is similar to Rudin-Keisler's ordering on ultrafilters (monads correspond with ultrafilters on the ring of  $Sd_{\mathcal{C}}$  classes by a one-one correspondence).

A monad  $\mu$  (in  $\{ \mathcal{C} \}$ ) is minimal in  $\{ \mathcal{C} \}$  iff each function  $F \in Sd_{\mathcal{C}}$  is either constant or one-one mapping on  $\mu$ .

In further considerations we shall limit ourselves to infinite monads. Moreover, we shall assume that they are subclasses of  $N$ , where we have natural ordering " $\leq$ ". This restriction is not substantial since there exists  $F \in Sd_0$  such that  $F$  is a one-one mapping of  $N$  onto  $V$  (see [V]).

For an easier typing we shall, through the whole paper, use for elements of  $N$  also small Latin letters, e.g.  $x \in \text{Def}$  is an abbreviation for  $x \in \text{Def} \cap N$ . This convention does not refer to the notation of set-definable functions and subsets of  $N$ .

When writing  $F: A \rightarrow B$  we bear in mind that  $F$  is a function such that  $F^*A \subseteq B$  (i.e. we do not ask for  $\text{dom}(F)=A$ ).

§ 1. At first we prove that monads cannot be divided into two parts by any definable element, i.e. that all elements of a monad have "the same position" with respect to the elements of  $\text{Def}$ .

**Definition.** Let  $X$  be a class. We shall call the class  $\{ \alpha; (\exists \beta, \gamma \in X) \beta \leq \alpha \leq \gamma \}$  a convex hull of  $X$  and denote  $\text{Cnh}(X)$ .

**Lemma 1.** Let  $\mu \subseteq N$  be a monad in  $\{ \mathcal{C} \}$ . Then  $\text{Cnh}(\mu) \cap \text{Def}_{\mathcal{C}} = \emptyset$ .

**Proof.** Let  $x \in \text{Cnh}(\mu) \cap \text{Def}_{\mathcal{C}}$ . Then there are  $t, u \in \mu$  such that  $t < x < u$  - a contradiction with the fact that the elements of a monad fulfil the same formulas.

**Theorem 1.** Let  $\mu$  be a monad in  $\{ \mathcal{C} \}$ . Then

(a)  $(\forall x) [x < \text{Cnh}(\mu) \Rightarrow (\exists a \in \text{Def}_{\mathcal{C}}) x < a < \text{Cnh}(\mu)]$ ,

(b)  $(\forall x) [x > \text{Cnh}(\mu) \Rightarrow (\exists a \in \text{Def}_{\mathcal{C}}) \text{Cnh}(\mu) < a < x]$ .

**Proof:** We shall prove (a), the assertion (b) can be proved analogously.

Take  $x < \text{Cnh}(\mu)$ .

Let  $\mu = \bigcap \{ X_n; n \in \mathbb{N} \}$ , where  $\{ X_n; n \in \mathbb{N} \}$  is a descending sequence (in  $\mathcal{C}$ ) of classes from  $Sd_0$  (see [V]). For  $X_n$  we shall ask moreover:  $\min(X_{n+1}) > \min(X_n)$ . (If  $\min(X_{n+1}) = \min(X_n)$ , we put  $\bar{X}_{n+1} = X_{n+1} - \{ \min(X_{n+1}) \}$  and

$\mu = \bigcap \{X_n; n \in \text{FN}\}$ .) We would like to prove:

$$(1) \quad (\exists n \in \text{FN}) \min(X_n) > x;$$

i.e. we show that  $\{X_n; n \in \text{FN}\}$  is cofinal with  $\mu$ . Denote  $a_n = \min(X_n)$  for every  $n \in \text{FN}$ . Let us prolong the sequence  $\{a_n; n \in \text{FN}\}$ . Then we can easily prove that

$$(2) \quad (\exists \beta)(\forall \gamma) [ \text{FN} < \gamma < \beta \Rightarrow a_\gamma \in \mu ].$$

To this end we shall construct a descending sequence  $\{\beta_n; n \in \text{FN}\}$  such that  $\beta_i > \text{FN}$  for every  $i \in \text{FN}$  and

$$(\forall \delta)(i < \delta < \beta_i \Rightarrow a_\delta \in X_i).$$

Obviously, it is enough to choose  $\beta_i$  like this: Let  $\beta_i$  be the maximal element such that for each  $\gamma$

$$i < \gamma < \beta_i \Rightarrow a_\gamma \in X_i.$$

But then there is such  $\beta^*$  that  $\text{FN} < \beta^* < \beta_i$  for all  $i \in \text{FN}$ . Hence (2) is valid.

Thus, for each  $\gamma$ , it is true that  $a_\gamma \in \mu$ . Let  $a_\xi$  be the smallest element of the prolonged sequence for which  $a_\xi > x$  holds. For proving (1) it is sufficient to show now that there exists  $k \in \text{FN}$  such that  $\xi = k$  (i.e. that the index of  $a_\xi$  is finite). From the construction of  $a_\xi$  we know, however, that  $a_{\xi-1} < x$  which implies  $\xi-1 \in \text{Fin}$  (in the opposite case we obtain  $a_{\xi-1} \in \mu$  which contradicts (2)); therefore  $\xi \in \text{Fin}$ . Thus  $a_k$  is the required element from  $\text{Def}_{\{c\}}$  fulfilling (a).

For further considerations it is useful to take the following notation:

$$\text{Def } N = \text{Def} \cap N, \quad \infty \text{Def } N = \text{Def } N - \text{FN}$$

and analogously for  $\text{Def}_{\{c\}}$ .

If we suppose that infinitely large definable numbers exist, then they can separate monads. It is therefore useful to examine elements which are very far (in natural ordering) one from the other. A characterization of "great distance" gives the following definition.

**Definition.** Let  $a, b \in \infty \text{Def } N$ . We say that  $b$  is much greater than  $a$  with respect to  $\text{FN}$  (notation  $a \ll_{\text{FN}} b$ ) iff

$$(3) \quad (\forall F \in \text{Sd}_0)(F: \text{FN} \rightarrow \text{FN} \Rightarrow F(a) < b).$$

Notice that, for  $\infty \text{Def } N = \emptyset$ ,  $a \ll_{\text{FN}} b$

describes the distance determined by all definable functions.

Very often, when working with functions from FN into FN, it is convenient to suppose that they are non-descending. Such a "trick" justifies the next lemma.

**Lemma 2.** There is a sequence of non-descending  $Sd_0$  functions  $\{F_i; i \in \mathbb{N}\}$  such that

- 1)  $(\forall i \in \mathbb{N})(\text{dom}(F_i) = \mathbb{N} \& F_i : \mathbb{N} \rightarrow \mathbb{N})$ .
- 2) For each  $Sd_0$  function  $G : \mathbb{N} \rightarrow \mathbb{N}$  there is  $F_k \in \{F_i; i \in \mathbb{N}\}$  such that  $G(\alpha) \leq F_k(\alpha)$  for every  $\alpha \in \text{dom}(G)$ .

**Proof.** Let us enumerate all  $Sd_0$  functions from FN into FN by  $\{G_i; i \in \mathbb{N}\}$ . Put, for every  $\alpha \in \mathbb{N}$ ,

$$(4) F_i(\alpha) = \max_{j \leq i} G_j(\alpha + 1).$$

Then  $\{F_i; i \in \mathbb{N}\}$  is the required sequence.

The next theorem asserts that each monad which comes into consideration intersects segments determined by the points which are very far one from the other.

**Theorem 2.** Let  $a, b \in {}^\omega \text{Def } \mathbb{N}$ ,  $a \ll_{\text{FN}} b$  and let  $\mu$  be such a monad that  $\mu \cap (\bigcap {}^\omega \text{Def } \mathbb{N}) \neq \emptyset$ . Then  $\mu \cap [a, b] \neq \emptyset$  (where  $[a, b] = \{x; a \leq x \leq b\}$ ).

**Proof.** Let  $A$  be a set-definable class,  $\mu \subseteq A \subseteq \mathbb{N}$ . We shall define a function  $G$  on  $\mathbb{N}$  as follows:

$$(5) (\forall t \in \mathbb{N}) G(t) = \min \{A - [ [\min(A - (t+1))] + 1 ]\}.$$

Realize that for setting  $G(t)$  we find, roughly speaking, the "first" element of  $A$  which is over  $t$  and then  $G(t)$  is the "second one with the same property".

Obviously  $G : \mathbb{N} \rightarrow \mathbb{N}$ ,  $G \in Sd_0$ . From Lemma 2 it follows that there is  $i \in \mathbb{N}$  such that  $F_i(t) > G(t)$  for every  $t \in \mathbb{N}$ . Furthermore, for all  $n \in \mathbb{N}$ , we have

$$(6) [n, F_i(n)] \cap A \neq \emptyset.$$

Let  $\gamma_0$  be the largest element such that every less element fulfils (6); i.e.

$$(\forall \gamma \leq \gamma_0) [\gamma, F_i(\gamma)] \cap A \neq \emptyset.$$

Then  $\gamma_0 \in \text{Def}$  and it is infinite. Therefore  $a \ll \gamma_0$  (since  $a \ll {}^\omega \text{Def } \mathbb{N}$ ) and (6) for  $a$  holds. Thus  $[a, F_i(a)] \cap A \neq \emptyset$ . Suppose  $\mu = \bigcap \{A_n; n \in \mathbb{N}\}$ , where  $\{A_n; n \in \mathbb{N}\}$  is a descending sequence of  $Sd_0$  classes. Then (note that  $\mu \subseteq A$ )

for each  $i \in FN$  we have  $[a, b] \cap A_i \neq \emptyset$ , which - due to the prolongation (see [LV]) - completes the proof.

**Remarks.** Definition of  $\ll_{FN}$ , Lemma 2 and the previous theorem can be reformulated into "parametric version" and proved analogously.

It follows from compactness of equivalences of indiscernibility that there are monads whose intersection with segments determined by points which are not very distant, is non-empty; hence the large distance is not a necessary condition.

In the next paragraph we show that each two different points of minimal monads are very far from each other.

Theorem 2 can be generalized also in another direction, as will be shown later. Before this we introduce a new notion and prove several assertions.

**Definition.** Let  $x \notin \text{Def}_{\{c\}}$ . Then we shall denote the class  $N - [Cnh(\text{Def}_{\{c\}} \cap x) \cup Cnh(\text{Def}_{\{c\}} \cap (N-x))]$   $\text{Int}_{\{c\}}(x)$  and call it an interval of  $x$  (with respect to the parameter  $c$ ).

Notice that  $\text{Int}_{\{c\}}(x)$  is the class of all natural numbers which have "the same position" w.r.t.  $\text{Def}_{\{c\}}$  as  $x$  has.

From the definition it follows immediately:

**Lemma 3.** Let  $x, y \notin \text{Def}_{\{c\}}$ , then the following holds:

- a)  $x \in \text{Int}_{\{c\}}(x)$ ,
- b)  $\text{Int}_{\{c\}}(x) \neq \text{Int}_{\{c\}}(y) \Rightarrow x \neq y$ ,
- c)  $x \frac{c}{c} y \Rightarrow \text{Int}_{\{c\}}(x) = \text{Int}_{\{c\}}(y)$ ,
- d)  $\text{Int}_{\{c\}}(x) = Cnh(\text{Int}_{\{c\}}(x)) = Cnh(\mu_{\{c\}}(x))$ .

Further we shall deal with a generalization of the minimal cut (i.e. FN).

**Definition.** Let  $X \subseteq \text{Def}_{\{c\}}$  such that  $(\forall t \in \text{Def}_{\{c\}}) t \in X \Leftrightarrow (\exists u \in X) u \geq t$ . Then  $X$  is called a cut.

We shall work only with cuts which have not the largest element - we denote them  $\wp$ .

**Lemma 4.** For each cut  $\wp$  there is  $\alpha \in N$  such that  $\wp = \text{Def}_{\{c\}} \cap \alpha$ ; such a cut we denote  $\wp(\alpha)$ .

**Proof.** Since  $\wp \subseteq \text{Def}_{\{c\}}$ ,  $\wp$  is a countable class and therefore there is an increasing sequence  $\{x_i \in \text{Def}_{\{c\}}; i \in FN\}$  which is cofinal with  $\wp$ .

We prolong this sequence in such a way that the monotony will be preserved. Let  $\{y_i; i \in \mathbb{N}\}$  be an enumeration of all elements from  $(\text{Def}_{\{c\}} - \rho)$ . Let  $\beta_i$  be the largest index of the prolonged sequence such that  $x_{\beta_i} < y_i$  (for all  $i \in \mathbb{N}$ ). Then  $\{\beta_i; i \in \mathbb{N}\}$  is a non-increasing sequence. Because  $\{\beta_i; i \in \mathbb{N}\}$  is countable, there exists  $\beta \notin \mathbb{N}$  such that  $\beta < \beta_i$  for every  $i \in \mathbb{N}$ . Put  $\alpha = x_\beta$ , obviously  $\rho(\alpha) = \text{Def}_{\{c\}} \cap \alpha$ .

Further we shall take the following notation: Let  $x \notin \text{Def}_{\{c\}}$ , then we put

$$\rho_{\{c\}}(x) = \text{Def}_{\{c\}} \cap x,$$

$$\bar{\rho}_{\{c\}}(x) = \text{Def}_{\{c\}} \cap (N-x).$$

Note that now

$$\text{Int}_{\{c\}}(x) = N - [\text{Cnh}(\rho_{\{c\}}(x)) \cup \text{Cnh}(\bar{\rho}_{\{c\}}(x))].$$

For an easier typing we shall use  $\rho$  when there is no danger of misunderstanding - only  $\rho$  instead of  $\rho_{\{c\}}(x)$  and analogously  $\bar{\rho}$  instead of  $\bar{\rho}_{\{c\}}(x)$ .

**Lemma 5.** Let  $x \notin \text{Def}_{\{c\}}$ ,  $X \in \text{Sd}_{\{c\}}$ . Then the following properties are equivalent:

- 1)  $X$  is cofinal with  $\rho_{\{c\}}(x)$ ,
- 2)  $X$  is coincial with  $\bar{\rho}_{\{c\}}(x)$ ,
- 3)  $X \cap \text{Int}_{\{c\}}(x) \neq \emptyset$ .

**Proof.** We prove only 1)  $\Leftrightarrow$  3); for 2)  $\Leftrightarrow$  3) it suffices to modify this proof for the inverse ordering.

Suppose 1) holds. Since  $\rho, \bar{\rho}$  are countable, we can enumerate them. Let  $\rho = \{a_i; i \in \mathbb{N}\}$ ,  $\bar{\rho} = \{b_i; i \in \mathbb{N}\}$ . From 1) it follows that

$$(\forall j \in \mathbb{N})(\exists i \in \mathbb{N}) [a_i, b_j] \cap X \neq \emptyset.$$

But  $\bigcap_{i \in \mathbb{N}} [a_i, b_i] = \text{Int}_{\{c\}}(x)$ ; hence 3) is valid.

For proving 3)  $\Rightarrow$  1) assume that  $a \in X \cap \text{Int}_{\{c\}}(x)$  and  $X$  is not cofinal with  $\rho$ . Then there is  $\bar{a} \in \rho$  such that there is no  $b \in (X \cap \rho) - \bar{a}$ . Put  $d = \min(X - (\bar{a} + 1))$ . Then  $d \in X$  and  $d \in \rho$  (since  $d \in \text{Def}_{\{c\}}$ ). Moreover,  $\bar{a} < d \leq a < \bar{\rho}$  - which contradicts 3).

The notion of the property of points "to be very distant" introduced above will be established now for more general cuts, namely  $\rho$ , and an analogue to Theorem 2 will be proved.

**Definition.** Let  $x \notin \text{Def}_{\mathcal{A}}(x)$ ,  $a, b \in \text{Int}_{\mathcal{A}}(x)$ . We say that  $b$  is much greater than  $a$  with respect to  $\mathcal{A}$  iff

$$(7) (\forall F \in \text{Sd}_{\mathcal{A}})(F: \mathcal{A} \rightarrow \mathcal{A} \Rightarrow F(a) \ll b);$$

notation  $a \ll \mathcal{A} b$  or only  $a \ll b$ , when there is no danger of confusion.

For proving the next theorem, we shall modify firstly Lemma 2.

**Lemma 6.** Let  $\mathcal{A}$  be a non-trivial cut on  $\text{Def}_{\mathcal{A}}$ . Then there is a sequence of non-descending  $\text{Sd}_{\mathcal{A}}$  functions  $\{F_i; i \in \mathbb{N}\}$  such that

- 1)  $(\forall i \in \mathbb{N}) (\text{dom}(F_i) = \mathbb{N} \ \& \ F_i: \mathcal{A} \rightarrow \mathcal{A})$ .
- 2) For each  $\text{Sd}_{\mathcal{A}}$  function  $G: \mathcal{A} \rightarrow \mathcal{A}$  there is  $F_k \in \{F_i; i \in \mathbb{N}\}$  such that  $G(\alpha) \leq F_k(\alpha)$  for every  $\alpha \in \text{dom}(G)$ .

**Proof.** Let  $\{G_i; i \in \mathbb{N}\}$  be an enumeration of all  $\text{Sd}_{\mathcal{A}}$  functions which transform  $\mathcal{A}$  into  $\mathcal{A}$ . Put

$$F_i(t) = \max(\bigcup_{j \leq i} G_j(t)).$$

When we prove that  $F_i: \mathcal{A} \rightarrow \mathcal{A}$  for each  $i \in \mathbb{N}$ , then  $\{F_i; i \in \mathbb{N}\}$  will be the required sequence. Suppose, by contradiction, that there exists  $F \in \{F_i; i \in \mathbb{N}\}$  such that  $F \not\subseteq \mathcal{A}$ . Let  $a \in \mathcal{A}$  and  $F(a) \notin \mathcal{A}$ . Then (note that  $F(a) \in \text{Def}_{\mathcal{A}}$ ) there is  $b \in \mathcal{A}$  such that  $F(a) \gg b$ . From our definition of  $\{F_i; i \in \mathbb{N}\}$  it follows that there are  $j \in \mathbb{N}$  and  $t \in \mathcal{A}$  such that  $G_j(t) \gg b$ . Denote  $\bar{t}$  the smallest of elements for which  $G_j(t) \gg b$ . Then  $\bar{t} \in \mathcal{A}$  (since  $\bar{t} \in \text{Def}_{\mathcal{A}}$ ), but  $G_j(\bar{t}) \notin \mathcal{A}$  (because  $G_j(\bar{t}) \gg b$  and  $b \in \mathcal{A}$ ) - a contradiction with  $G_j: \mathcal{A} \rightarrow \mathcal{A}$ .

**Theorem 3.** Let  $a, b \in \text{Int}(x)$ ,  $a \ll (x) b$  and let  $\mathcal{A}$  be such a monad that  $\mathcal{A} \cap \text{Int}(x) \neq \emptyset$ . Then  $\mathcal{A} \cap [a, b] \neq \emptyset$ .

**Proof.** For proving this assertion it suffices to use Lemma 6 and modify the proof of Theorem 2. Analogously to (5) we choose the class  $A$  and define the function  $G$ .

We have, however, to show that  $G: \mathcal{A} \rightarrow \mathcal{A}$  (an analogous fact for  $G$  in the proof of Theorem 2 was obvious). Let  $u \in \mathcal{A}$ ; then there exists  $v \gg u$  such that  $v \in A$ . Take the smallest of such elements - denote it  $v_1$ . Then  $v_1 \in \mathcal{A}$  ( $A \cap \text{Int}(x) \neq \emptyset$  since  $\mathcal{A} \subseteq A$ ; but this implies that  $A$  is cofinal with  $\mathcal{A}$  - see Lemma 5). Repeat now this consideration for "starting point"  $v_1$ ; it brings us to  $w_1 \in \mathcal{A}$ . But  $G(u) = w_1$  and consequently  $G: \mathcal{A} \rightarrow \mathcal{A}$ .

When "checking" furthermore the proof of Theorem 2, we obtain here ins-



stead of (6)

$$(8) (\forall p \in \mathcal{P})(\forall t \in \mathcal{P}) [t; F_i(t)] \cap A \neq \emptyset.$$

If  $\gamma_0$  is the greatest element for which (8) holds, we have that  $\gamma_0 \in \text{Def-}\mathcal{P}$ . Hence  $a < \gamma_0$  (since  $a < \bar{\mathcal{P}}$ ) and thus  $[a; F_i(a)] \cap A \neq \emptyset$ . For completing this proof it is enough to use the same arguments as in the proof of Theorem 2.

**Remark.** We know that if  ${}^\infty\text{Def } N = \emptyset$ , then  $FN = \text{Def} \cap N$ . When, however,  ${}^\infty\text{Def } N \neq \emptyset$ , a natural question arises: How do all elements under  ${}^\infty\text{Def } N$  and reasonably definable look like? Such sort of considerations has brought us to the following definition.

**Definition.** Let  $a \notin \text{Def}_{\{c\}}$ . We put

$$\text{Def}_{\{c\}}^{\mathcal{P} \rightarrow \mathcal{P}}(\{a\}) = \{x; (\exists F)(F \in \text{Sd}_{\{c\}} \ \& \ F: \mathcal{P} \rightarrow \mathcal{P} \ \& \ x = F(a))\};$$

we write here briefly only  $\mathcal{P}$  instead of  $\mathcal{P}_{\{c\}}(a)$ .

When we compare this definition with the one of  $a <<_{\mathcal{P}} b$ , we obtain immediately that

$$a <<_{\mathcal{P}} b \equiv \text{Def}_{\{c\}}^{\mathcal{P} \rightarrow \mathcal{P}}(\{a\}) \subseteq b.$$

The next theorem asserts that under the same assumptions as in Theorem 3 there exists even an infinite set inside  $\mu \cap [a, b]$ . At first we prove, however, an auxiliary result.

**Lemma 7.** Let  $a <<_{\mathcal{P}} b$ ,  $d \in [a, b]$ . Then either  $a <<_{\mathcal{P}} d$  or  $d <<_{\mathcal{P}} b$ .

**Proof.** Evidently it suffices to verify that

$$d < \text{Def}_{\{c\}}^{\mathcal{P} \rightarrow \mathcal{P}}(\{a\}) \Rightarrow \text{Def}_{\{c\}}^{\mathcal{P} \rightarrow \mathcal{P}}(\{d\}) \subseteq \text{Def}_{\{c\}}^{\mathcal{P} \rightarrow \mathcal{P}}(\{a\}).$$

Let  $F \in \text{Sd}_{\{c\}}$ ,  $F: \mathcal{P} \rightarrow \mathcal{P}$ . We have to show that there exists  $G \in \text{Sd}_{\{c\}}$ ,  $G: \mathcal{P} \rightarrow \mathcal{P}$  such that  $G(a) \geq F(d)$ . It follows from Lemma 6 that there are non-descending  $\text{Sd}_{\{c\}}$  functions  $G_1, G_2$  such that  $d \leq G_1(a)$  and  $F(d) \leq G_2(d)$ . Then, however,  $G_2(d) \leq G_2(G_1(a))$ . Put now  $G = G_2 \circ G_1$ .

**Theorem 4.** Let  $a, b \in \text{Int}(x)$ ,  $a <<_{\mathcal{P}(x)} b$  and  $\mu$  be such a monad that  $\mu \cap \text{Int}(x) \neq \emptyset$  (hence  $\mu \subseteq \text{Int}(x)$ ). Then there is  $m \in \text{Fin}$  such that  $m \subseteq \mu \cap [a, b]$ .

**Proof.** The monad  $\mu$  is a  $\pi$ -class and therefore it is revealed. Since  $[a, b]$  is a set,  $\mu \cap [a, b]$  is revealed, too. Hence it is sufficient to prove that there is an infinite countable class  $C$  such that  $C \subseteq \mu \cap [a, b]$ . For making the proof shorter we show only that  $C \subseteq \mu \cap \{x; a < x < b\}$ . We shall construct the class  $C$  by means of Lemma 7 and Theorem 3.

Let  $a_1 \in \mu \cap \{x; a < x < b\}$ ; we have that  $a < \rho(x)a_1$  or  $a_1 < \rho(x)b$ . Suppose e.g. that  $a < \rho(x)a_1$  and apply again the above mentioned assertion, etc. Thus we obtain a countable sequence  $\{a_i; i \in \mathbb{N}\}$  of elements from  $\mu \cap \{x; a < x < b\}$ . Put  $C = \{a_i; i \in \mathbb{N}\}$ .

**Remarks.** When substituting  $\rho(x)$  by FN in Theorem 4, we have an analogous generalization of Theorem 2.

Theorems 3 and 4 can be similarly proved also for their parametrical version.

The following theorem speaks about monads generally, but its main significance will not become evident before studying monads of indiscernibles. In the forthcoming paper we show that if Ind is a monad of indiscernibles for the language L and  $t \in \text{Ind}$  then  $t \cap \text{Ind}$  is a monad of indiscernibles for  $L_{\{t\}}$ . Theorem 5 implies that the assumption  $t \in \text{Ind}$  is a substantial one.

**Theorem 5.** Let  $\mu$  be a monad. Then there are  $c, x$  such that  $\mu \cap \text{Int}_{\{c\}}(x)$  is not a monad in  $\frac{\mu}{\{c\}}$ .

**Proof.** Since  $\mu \neq \text{Fin}$ , there is  $c \in \mu$  which is infinite, too. Then  $(\frac{\mu}{\{c\}})$  is compact) there are  $x, y \in c, x \neq y$  such that  $x \frac{\mu}{\{c\}} y$ . Suppose  $x < y$ . For proving Theorem 5 it suffices to find  $t \in \mu \cap \text{Int}_{\{c\}}(x)$  for which  $\neg(t \frac{\mu}{\{c\}} x)$  holds. Denote the smallest element of  $c$  which is greater (in  $<$ ) than  $x$  by  $z$ . Obviously  $z \in \mu$  (since  $z \in c$ ). Moreover,  $z \in \text{Int}_{\{c\}}(x)$ ; for this realize that  $x, y \in \text{Int}_{\{c\}}(x) = \text{Cnh}(\text{Int}_{\{c\}}(x))$  - see Lemma 3. Hence  $z \in \mu \cap \text{Int}_{\{c\}}(x)$ .

We show that  $\neg(z \frac{\mu}{\{c\}} x)$ . Suppose that  $z \frac{\mu}{\{c\}} x$ . From the construction of  $z$  it follows that  $z$  is definable from  $x$  and  $c$ . This fact implies the existence of  $\text{Sd}_{\{c\}}$  function  $F$  such that  $F(x)=z$ . Then, however,  $F(x) \frac{\mu}{\{c\}} x$  which resulted in  $F(x)=x$  (see Theorem 1 from [Č-K]) and hence  $z=x$  - a contradiction with the definition of  $z$ . For completing the proof it is enough to put  $z=t$ .

**§ 2.** In the first theorem of this paragraph we prove that there exists a minimal monad close over FN which, moreover, possesses the following property: each set-definable function (with parameter  $c$ ) either maps FN into FN or transfers this monad over an infinitely large definable element.

At first we shall formulate an assertion which is an immediate consequence of Lemma 5.

**Lemma 8.** Let  $x \in \text{Def}_{\{c\}}$  be not cofinal with FN. Then there is  $k \in \text{FN}$  such

that  $(x-k) \cap FN = \emptyset$ .

**Theorem 6.** There exists a minimal monad  $\mu$  in the class  $X = \{\alpha; FN \prec \alpha \prec \langle {}^{\infty}\text{Def}_{\{c\}}N \rangle\}$  such that for every  $F \in \text{Sd}_{\{c\}}$ ,  $F: N \rightarrow N$ , there is  $Z \in \text{Sd}_{\{c\}}$ ,  $Z \geq \mu$ , such that either (i) or (ii) takes place, where  
 (i)  $F \circ Z : FN \rightarrow FN$ ,  
 (ii)  $F \circ Z \cap FN = \emptyset$ .

**Proof.** For  ${}^{\infty}\text{Def}_{\{c\}}N = \emptyset$  we have  $X = N \text{--} FN$  and  $c \in \text{Def}$ . Then the assertion is valid - see [Č-K], Theorem 11, and realize that (i) is true for each  $F \in \text{Sd}_0$ .

Assume further  ${}^{\infty}\text{Def}_{\{c\}}N \neq \emptyset$ . We shall construct a descending sequence  $\{x_n; n \in FN\}$  with the following properties:

- 1)  $(\forall n \in FN) x_n \in \text{Def}_{\{c\}}$ ;
- 2)  $(\forall n \in FN) x_n$  is cofinal with  $FN$ ;
- 3)  $(\forall Y \in \text{Sd}_{\{c\}}) [Y \subseteq N \Rightarrow (\exists n \in FN) (x_n \subseteq Y \vee x_n \subseteq N \text{--} Y)]$ ;
- 4) Let  $\{F_n; n \in FN\}$  be an enumeration of all  $\text{Sd}_{\{c\}}$  functions. Then  $F_n$  is either constant or a one-one mapping on  $x_{n+1}$  (for all  $n \in FN$ );
- 5) If  $\{y_n; n \in FN\}$  is an enumeration of  ${}^{\infty}\text{Def}_{\{c\}}N$ , then  $x_{n+1} \subseteq y_n$  (for all  $n \in FN$ );
- 6) For each  $F_k \in \{F_n; n \in FN\}$  either (i) or (ii) is true when we put  $Z = x_{k+1}$ .

The sets  $x_n$  will be constructed by induction based on  $n$ .

At first take, arbitrarily,  $x_1 \in {}^{\infty}\text{Def}_{\{c\}}N$ ; denote  $\hat{x}_1 = x_1 \cap y_1$ , where  $y_1$  is the first element of our enumeration of  ${}^{\infty}\text{Def}_{\{c\}}N$ . We shall investigate  $\hat{x}_1 = \hat{x}_1 \cap \text{dom}(F_1)$ , where  $F_1$  is the first element of  $\{F_n; n \in FN\}$ .

If  $\hat{x}_1$  is not cofinal with  $FN$ , put  $x_2 = \hat{x}_1 \text{--} \text{dom}(F_1)$ . Obviously  $x_2$  possesses each required property except 3), which will be examined later for all possible cases all at once.

For  $\hat{x}_1$  being cofinal with  $FN$  we shall examine the system of classes of decomposition of  $\hat{x}_1$  according to the equivalence  $a \sim b \Leftrightarrow F_1(a) = F_1(b)$ . There are two possibilities:

- a)  $(\exists t \in \text{rng}(F_1)) F_1^{-1} \{t\} \cap \hat{x}_1$  is cofinal with  $FN$ ,
- b) the negation of a) holds.

In the case a) take  $t \in \text{rng}(F_1)$  such that  $F_1^{-1} \{t\} \cap \hat{x}_1$  is cofinal with  $FN$ . Then evidently  $t \in \text{Def}_{\{c\}}$  (realize for it that  $F_1 \in \text{Sd}_{\{c\}}$  and  $F_1 \circ FN \subseteq \text{Def}_{\{c\}} \neq \emptyset$ ). Put  $\tilde{x}_1 = F_1^{-1} \{t\} \cap \hat{x}_1$ ; except 3) and 6) all conditions are valid.

Assume further b). Then we can take  $\tilde{x}_1$  as the set of all smallest elements from classes of decomposition of  $\hat{x}_1$  (according to  $F_1$ ). Obviously 1), 4) and 5) hold. We prove now 2) for  $\tilde{x}_1$ . Suppose the contrary. Then there is  $k \in FN$

such that  $(\tilde{x}_1 - k) \cap FN = \emptyset$  - see Lemma 8. Put  $\alpha = \min(\tilde{x}_1 - k)$ ; then  $\alpha \notin FN$ . Let further  $w$  contain all classes of decomposition of  $\tilde{x}_1$  (over  $F_1$ ) such that their smallest elements belong to  $\tilde{x}_1 - k$ . Then for each  $z \in w$  we have  $z \geq \alpha$  and hence  $w$  is not cofinal with  $FN$ . The class  $\hat{x}_1 - w$  has, however, only a finite number of elements from the decomposition. Thus at least one of them must be cofinal with  $FN$  - a contradiction (we suppose that none is).

Now we shall construct  $x_2$ . If  $F$  is constant on  $\tilde{x}_1$ , it suffices to put  $x_2 = \tilde{x}_1$  and all conditions except 3) are valid for  $x_2$ . Assume further that  $F_1$  is one-one on  $\tilde{x}_1$ . If  $F_1'' \tilde{x}_1$  is not cofinal with  $FN$ , there is  $k \in FN$  such that  $(F_1'' \tilde{x}_1 - k) \cap FN = \emptyset$  - see Lemma 8. Put  $x_2 = \tilde{x}_1 - F_1^{-1}k$ ; it is evident that 1), 2), 4), 5) are true. For 6) realize that now (ii) takes place.

Suppose  $F_1'' \tilde{x}_1$  is cofinal with  $FN$ . Without loss of generality we may think that  $F_1(t) > t$  (otherwise we shall take  $F_1 + 1$ ). The set  $x_2$  will be constructed by induction; we demand, at the same time,  $x_2 \subseteq \tilde{x}_1$ . Let  $t_1$  be such an element that  $F_1(t_1)$  is minimal in  $F_1'' \tilde{x}_1$ . Obviously  $t_1 \in FN$  (cofinality condition) and  $F_1''(\tilde{x}_1 - (t_1 + 1))$  is cofinal with  $FN$ . Let  $t_2$  be such an element that  $F_1(t_2)$  is minimal in  $F_1''(\tilde{x}_1 - (t_1 + 1))$ , etc. This set-definable construction will go until a certain number  $\eta \in N$ . When we put now  $x_2 = \{t_\gamma; \gamma \in \eta\}$ , we have that 1), 2), 4) and 5) are fulfilled. Moreover, from our construction of  $x_2$ , it follows that (i) comes (if  $t_1 \in x_2 \cap FN$ , then  $i \in FN$  and therefore  $F_1(t_1) \in FN$ ). Hence 6) is also true for  $x_2$ .

Analogously we can construct  $x_3$  from  $x_2$ ,  $x_4$  from  $x_3$ , etc. The obtained sequence  $\{x_n; n \in FN\}$  possesses all required properties except 3), which we prove now. Let  $Y \in Sd_{\{FC\}}$ ,  $Y \subseteq N$  and let  $\chi_Y$  be its characteristic function. Then  $\chi_Y = F_k$ , where  $F_k$  is the function from our enumeration which is on  $x_{k+1}$  either one-one or constant. Since  $\text{dom}(\chi_Y) \neq \text{Fin}$  and  $\text{rng}(\chi_Y) = \{0, 1\}$ ,  $\chi_Y$  has to be constant on  $x_{k+1}$ . If for each  $t \in x_{k+1}$  we have  $\chi_Y(t) = 0$ , then  $x_{k+1} \subseteq N - Y$ ; in the case  $\chi_Y(t) = 1$  we obtain  $x_{k+1} \subseteq Y$ . The sequence  $\{x_n; n \in FN\}$  fulfils therefore 3) and hence  $\bigcap \{x_n; n \in FN\}$  is a monad - let us denote it  $\mu$ . Owing to 4)  $\mu$  is a minimal monad.

For proving  $\mu \subseteq X$  it suffices to show that  $X \cap \mu \neq \emptyset$ . Assume the contrary; let  $u \in X$ ,  $t \in \mu$ . Then  $\text{Int}_{\{FC\}}(u) \cap \text{Int}_{\{FC\}}(t) = \emptyset$ . Construct  $\rho_{\{FC\}}(u)$  and  $\rho_{\{FC\}}(t)$ . Evidently  $\rho_{\{FC\}}(u) = FN$  and  $\rho_{\{FC\}}(t) \neq FN$ , which implies the existence of  $\alpha \in {}^\omega \text{Def}_{\{FC\}} N$  such that  $\alpha \in \rho_{\{FC\}}(t)$ . Then  $\alpha < t$ , but  $\alpha < \mu$ , therefore there is  $k \in FN$  for which  $\alpha < x_k$  - a contradiction with 2).

For completing the proof it remains to show that for each  $F \in Sd_{\{FC\}}$ ,  $F: N \rightarrow N$ , there is  $\mathcal{X} \in Sd_{\{FC\}}$ ,  $\mathcal{X} \supseteq \mu$ , such that either (i) or (ii) takes place.

ce. Obviously it is enough to find  $F_k \in \{F_n; n \in \mathbb{N}\}$  for which  $F = F_k$  and put  $\mathcal{X} = \mathcal{X}_{k+1}$ .

**Remark.** In a forthcoming paper we prove the consistency of the existence of a minimal monad  $\mu$  in  $X$  such that for a certain  $Sd_0$  function there is no  $\mathcal{X} \in Sd_0$ ,  $\mathcal{X} \supseteq \mu$ , with properties (i) or (ii).

There is a question: For what  $t$  there exists a minimal monad inside  $Int_{\{C\}}(t)$ . A partial solution gives the following theorem, which is a simple consequence of Theorem 6.

**Theorem 7.** Let  $t \notin Def_{\{C\}}$ . Let there exist  $\alpha \in X = \{\gamma; \mathbb{N} < \gamma < \infty, Def_{\{C\}} \mathbb{N}\}$  and a non-descending  $Sd_{\{C\}}$  function  $F: \mathbb{N} \rightarrow \mathbb{N}$  such that  $F(\alpha) \in Int_{\{C\}}(t)$ . Then there is a minimal monad  $\nu \subseteq Int_{\{C\}}(t)$ .

**Proof.** Firstly we show that we can assume  $dom(F) = \mathbb{N}$ . If  $dom(F) = m$ , denote  $\sigma = \max(m)$  and for all  $t > \sigma$  put  $F(t) = F(\sigma)$ . Let further  $dom(F)$  be a proper class. If  $F$  is not defined in some  $\sigma$ , put  $F(\sigma) = F(\beta)$ , where  $\beta$  is the smallest of all elements from  $dom(F)$  which are larger than  $\sigma$ .

Suppose now  $dom(F) = \mathbb{N}$ . Evidently,  $F$  remains to be non-descending. This, however, implies that  $F'' Int_{\{C\}}(\alpha) = Int_{\{C\}}(F(d))$ . Let  $\mu$  be a minimal monad in  $X$  - it exists due to Theorem 6. Denote  $\nu = F''\mu$ . Then  $\nu \subseteq Int_{\{C\}}(t)$  (if  $\beta \in \mu$  then  $Int_{\{C\}}(\beta) = Int_{\{C\}}(\alpha) = X$  and  $F(\beta) \in Int_{\{C\}}(t)$ ). The monad  $\nu$  cannot be a trivial one, since  $t \notin Def_{\{C\}}$  and hence  $Int_{\{C\}}(t)$  cannot be a singleton. As  $\mu$  is minimal,  $F$  has to be one-one on  $\mu$  and hence  $\nu$  is minimal, too.

Next four theorems show an analogue of one property of classes of indiscernibles for minimal monads.

**Theorem 8.** Let  $\mu$  be a minimal monad in  $\frac{\mathbb{Q}}{\{C\}}$  which is a semiset. Let  $F \in Sd_{\{C\}}$  be such a function that  $F(t) < t$  for some (and hence every)  $t \in \mu$ . Then either 1) or 2) is valid, where

- 1)  $(\forall u \in \mu)(u < t \Rightarrow u < F(t))$ ;
- 2)  $(\exists d \in Def_{\{C\}})(\forall t \in \mu) F(d) < t < d$ .

**Proof.** Since  $F$  is defined on the minimal monad  $\mu$ ,  $F$  is either constant or one-one on  $\mu$ .

Suppose, at first, that  $F$  is constant on  $\mu$ . Denote  $F(t) = e$  for  $t \in \mu$ ; evidently  $e \in Def_{\{C\}}$ . Construct  $F^{-1}''\{e\}$ . Obviously  $F^{-1}''\{e\} \in Sd_{\{C\}}$  and  $\mu \subseteq F^{-1}''\{e\}$ . Because  $\mu$  is a semiset, there exists  $\bar{e} \in Def_{\{C\}}$  such that  $\bar{e} > \mu$ . Put  $d = \max((F^{-1}''\{e\}) \cap \bar{e})$ . Since  $e, \bar{e} \in Def_{\{C\}}$ ,  $F \in Sd_{\{C\}}$  and  $F(d) = e$ ,

we have  $d \in \text{Def}_{\{C\}}$ . Moreover,  $\mu \in \bar{e}$  (since  $\bar{e} > \mu$ ),  $\mu \in F^{-1}\{e\}$ , therefore  $d > \mu$ . But  $F(d) = e = F(t) < t$  for each  $t \in \mu$ . Hence  $F(d) < t < d$  and 2) is fulfilled.

Let further  $F$  be one-one on  $\mu$ . We shall assume  $\gamma 1) \& \gamma 2)$ . It follows from  $\gamma 2)$  that there exists  $t \in \mu$  such that  $F^{-1}\rho_{\{C\}}(t) \subseteq \rho_{\{C\}}(t)$ . Let us fix such  $t$ . Put  $k = \min(\text{dom}(F^{-1}))$  and define for each  $u \geq k$  a function  $H$  as follows:

$$H(u) = \max(F^{-1}(u+1) \cup \{u+1\}).$$

Obviously  $H \in \text{Sd}_{\{C\}}$  is a non-descending function and  $H(u) > u$  for each  $u \geq k$ . Define further a function  $G$  as follows:

$$G(0) = k \& G(\gamma+1) = H(G(\gamma)),$$

i.e.  $G(\gamma)$  is the  $\gamma$ -th iteration of  $H$  when starting from  $k$ . Finally, put for each  $u \geq k$

$$\bar{F}(u) = \max(\{\gamma; G(\gamma) < u\}).$$

Then  $\bar{F} \in \text{Sd}_{\{C\}}$  and  $\bar{F}$  is also a non-descending function. Denote  $\alpha = \bar{F}(t)$ . We would like to prove that  $\alpha \in \text{Def}_{\{C\}}$ . In accordance with our assumption  $\gamma 1)$  we have that there is  $v \in \mu$  such that  $v < t$  and  $F(t) \in v$ . If we prove that  $\bar{F}(t) = \bar{F}(v)$ , we shall know that also  $\bar{F}(v) = \alpha$  and since  $\mu$  is a minimal monad, this will imply that  $\bar{F}$  is constant on  $\mu$  and therefore  $\bar{F}(t) = \alpha \in \text{Def}_{\{C\}}$ .

Let us prove  $\bar{F}(t) = \bar{F}(v)$ . Since  $\bar{F}$  is a non-descending function, the inequality  $v < t$  implies  $\bar{F}(v) \leq \bar{F}(t) = \alpha$ . For proving the converse inequality we shall show firstly that  $G(\alpha) < t$ . Suppose  $G(\alpha) \geq v$ . From the construction of  $\alpha$  it follows that  $G(\alpha) < t$  and  $G(\alpha+1) \geq t$ . Denote  $\bar{F}(v) = \beta$ . Then  $G(\alpha+1) \geq t$  implies  $G(\beta+1) \geq v$ . But  $G(\alpha-1) < v$  (since  $G(\alpha-1) < H^{-1}(t)$ ). Thus  $\alpha > \beta \geq \alpha-1$  and hence  $\beta = \alpha-1$ . This is, however, a contradiction, since  $t, v$  belong to the same monad. Therefore  $G(\alpha) < v$ , which implies  $\bar{F}(v) \geq \alpha = \bar{F}(t)$ . Hence  $\bar{F}(v) = \bar{F}(t) = \alpha$ .

Since  $\alpha \in \text{Def}_{\{C\}}$ , also  $G(\alpha) \in \text{Def}_{\{C\}}$  and at the same time  $G(\alpha) < t$ . Obviously  $G(\alpha+1) \in \text{Def}_{\{C\}}$ , too, and from the definition of  $\bar{F}$  we have  $G(\alpha+1) > t$ . Remember now that  $G(\alpha+1) = H(G(\alpha))$  and the fact that for  $H$  it is true  $H\rho_{\{C\}}(t) \subseteq \rho_{\{C\}}(t)$  - see Lemma 6. We have proved here, however, that  $H$  does not fulfil this inclusion - a contradiction.

**Remark.** Note that for the classes of indiscernibles 2) from the previous theorem may be strengthened to the assertion "F is constant". In the forthcoming paper we prove that it is consistent with AST that there exists a

minimal monad  $\mu$  for which 2) cannot be substituted by "F is constant on  $\mu$ ".

Now we shall present several modifications of Theorem 8.

**Theorem 9.** Let  $\mu$  be a minimal monad in  $\frac{\mathbb{O}}{\mathbb{C}}$  which is a proper class. Let  $F \in \text{Sd}_{\mathbb{C}}$  be such a function that  $F(t) < t$  for every  $t \in \mu$ . Then either 1) or 2) is valid, where

- 1)  $(\forall u \in \mu)(u < t \Rightarrow u < F(t))$ ;
- 2) F is constant on  $\mu$ .

**Proof.** Suppose that F is one-one on  $\mu$ . In this case realize that the condition 2) from Theorem 8 cannot come (since  $\text{Def}_{\mathbb{C}} < \mu$  and thus for each  $F \in \text{Sd}_{\mathbb{C}}$  and each  $t \in \mu$   $F^{-1} \circ \text{Def}_{\mathbb{C}}(t) \subseteq \text{Def}_{\mathbb{C}}(t)$  holds - see Lemma 6) and proceed analogously as in the proof of this theorem.

**Theorem 10.** Let  $\mu$  be a minimal monad in  $\frac{\mathbb{O}}{\mathbb{C}}$  which is a semiset. Let  $F \in \text{Sd}_{\mathbb{C}}$  be such a function that  $F(t) > t$  for one (and hence every)  $t \in \mu$ . Then either 1) or 2) is valid, where

- 1)  $(\forall v \in \mu)(v > t \Rightarrow v > F(t))$ ;
- 2)  $(\exists d \in \text{Def}_{\mathbb{C}})(\forall t \in \mu) d < t < F(d)$ .

**Proof.** Let F be constant on  $\mu$ . Denote  $F(t) = e$  for each  $t \in \mu$ . Construct  $F^{-1}\{e\}$ ; then this class is a non-empty  $\text{Sd}_{\mathbb{C}}$  class (since  $e \in \text{Def}_{\mathbb{C}}$  and  $F \in \text{Sd}_{\mathbb{C}}$ ). Put  $d = \min(F^{-1}\{e\})$ . Then  $d \in \text{Def}_{\mathbb{C}}$ ,  $d < t$  and  $F(d) = F(t) > t$  - 2) is valid.

Suppose further that F is one-one on  $\mu$ . Recall that also  $F^{-1}$  is one-one on  $\mu$  and  $\nu = F^{-1}\mu$  is a minimal monad in  $\frac{\mathbb{O}}{\mathbb{C}}$ . Moreover,  $\text{Cnh}(\mu) = \text{Cnh}(\nu)$ , since F(t) and t are "likewise" situated with respect to  $\text{Def}_{\mathbb{C}}$ ; if  $\text{Int}_{\mathbb{C}}(F(t)) \neq \text{Int}_{\mathbb{C}}(t)$ , then 2) holds. Now apply Theorem 8 on  $\nu$ ,  $F^{-1}$  and F(t).

**Theorem 11.** Let  $\mu$  be a minimal monad in  $\frac{\mathbb{O}}{\mathbb{C}}$  which is a proper class. Let  $F \in \text{Sd}_{\mathbb{C}}$  be such a function that  $F(t) > t$  for every  $t \in \mu$ . Then

$$(\forall v \in \mu)(v > t \Rightarrow v > F(t)).$$

**Proof.** Note, at first, that F cannot be constant on  $\mu$  (in the opposite case we have  $F(t) \in \text{Def}_{\mathbb{C}}$  for  $t \in \mu$ , which is in contradiction with  $t > \text{Def}_{\mathbb{C}}$  and  $F(t) > t$ ).

Hence F is one-one on  $\mu$ . Put  $G = F^{-1}$ . Then G is also one-one on  $\mu$  and  $G(t) < t$  for each  $t \in \mu$ . Denote  $\nu = F^{-1}\mu$ ; then  $\nu$  is a minimal monad which is a proper class. Now apply Theorem 9 on G and  $\nu$ .

Especially, let us stress one consequence of the previous theorems for minimal monads lying close behind FN.

**Theorem 12.** Let  $\mu \in {}^{\infty}\text{Def } N$  be a minimal monad,  $x, y \in \mu$  and  $x < y$ . Then  $\text{Def}_{\{c\}}^{\text{FN} \rightarrow \text{FN}}(\{x\}) < y$ .

**Proof.** Suppose, at first, that  ${}^{\infty}\text{Def}_{\{c\}} N = \emptyset$ . Then  $c \in \text{Def}$  and therefore  $\mu$  is a proper class. In this case it is sufficient to use Theorem 11.

If  ${}^{\infty}\text{Def}_{\{c\}} N \neq \emptyset$ , we shall apply Theorem 10. Since we work with a function which maps FN into FN, the case 2) from this theorem cannot set in, the case 1) is exactly what we want to prove here.

The last theorem which can be also reformulated into a parametric version, and its corollary describe several equivalent expressions of the property "to be very far one from the other".

**Theorem 13.** Let  $\alpha \in X = \{ \eta, \text{FN} < \eta < {}^{\infty}\text{Def } N \}$ . Then for each  $\gamma$  the following are equivalent:

- (i)  $\alpha <<_{\text{FN}} \gamma$ ;
- (ii)  $(\exists \beta \in X)(\beta < \gamma \ \& \ \alpha < {}^{\infty}\text{Def}_{\{\beta\}} N)$ .

**Proof.** (i)  $\Rightarrow$  (ii). Let  $\mu$  be the monad from Theorem 6. In accordance with Theorem 4 there are  $x, y \in \mu$  such that  $\alpha < x < y < \gamma$  (take  $x, y \in m$ ). We would like to prove that for each  $F \in \text{Sd}_0$ ,  $F: N \rightarrow N$ , we obtain  $F(y) > x$ . From the construction of  $\mu$  we know that there is  $\mathcal{Z} \in \text{Sd}_0$  such that  $\mu \subseteq \mathcal{Z}$  and  $F''\mathcal{Z} \cap \text{FN} = \emptyset$  or  $F \upharpoonright \mathcal{Z}: \text{FN} \rightarrow \text{FN}$ . In the first case we have that  $\min(F''\mathcal{Z}) > \text{FN}$  and  $\min(F''\mathcal{Z}) \in \text{Def}$ . This implies  $\min(F''\mathcal{Z}) > X$  and therefore  $F''\mathcal{Z} > X$ . Since  $y \in \mu \subseteq X$  and  $F(y) > X$ , we obtain  $F(y) > x$ . If  $F \upharpoonright \mathcal{Z}: \text{FN} \rightarrow \text{FN}$ , we apply Theorem 8. The assertion 2) from this theorem is now excluded, hence 1) has to take place. Put now  $\beta = y$ ; we have then  ${}^{\infty}\text{Def}_{\{\beta\}} > x > \alpha$ .

(ii)  $\Rightarrow$  (i). We have to prove now that for each  $F \in \text{Sd}_0$ ,  $F: \text{FN} \rightarrow \text{FN}$  we have  $F(\alpha) < \gamma$ . In accordance with Lemma 6 we can suppose that  $F$  is a non-descending function. Put  $G(t) = \min(F^{-1}\{t\})$ , where  $u$  is the largest element from  $\text{rng}(F)$  which is smaller or equal to  $t$ . Then  $G$  is also non-descending. We have  $G(\beta) \in \text{Def}_{\{\beta\}}$  and, since  $\beta \in X$ , at the same time  $G(\beta) \notin \text{FN}$  (otherwise  $F$  is constant). Hence  $G(\beta) > \alpha$  and  $F(\alpha) \in \beta < \gamma$ .

**Corollary.** For  $\alpha < \beta$ ,  $\alpha \in X = \{ \eta, \text{FN} < \eta < {}^{\infty}\text{Def } N \}$ , the following are equivalent:

- 1)  $\alpha <<_{\text{FN}} \beta$ ;



- 2)  $(\exists \text{e.u.}A)(\alpha \in E_A(FN) < \beta)$ , where e.u.A means an endomorphic universe A;
- 3)  $(\exists \text{e.u.s.}A)(\alpha \in \text{Ex}_A(FN) < \beta)$ , where e.u.s.A means e.u. with standard extension;
- 4)  $(\exists \text{e.u.}A)(\exists \gamma \leq \beta)(\alpha, \gamma \in E_A(FN) \& \alpha \in E_{A[\gamma]}(FN))$ ;
- 5)  $(\exists \text{e.u.s.}A)(\exists \gamma \leq \beta)(\alpha, \gamma \in \text{Ex}_A(FN) \& \alpha \in E_{A[\gamma]}(FN))$ ;
- 6)  $(\exists \gamma)(\alpha < \gamma \leq \beta \& \alpha <^{\infty} \text{Def}_{\{\gamma\}} N)$ ;
- 7)  $(\exists \text{e.u.}A)(\forall f \in A)(f \text{ " } FN \subseteq FN \Rightarrow f(\alpha) < \beta)$ .

**Proof** - see [Č-I].

#### References

- [V] P. VOPĚNKA: Mathematics in the Alternative Set Theory, Teubner-Texte, Leipzig 1979.
- [Č-K] K. ČUDA, B. KUSSOVÁ: Basic equivalences in the alternative set theory, Comment. Math. Univ. Carolinae 23(1982), 629-644.
- [Č-I] K. ČUDA, A. TZOUVARAS: Endomorphic cuts and tails, Comment. Math. Univ. Carolinae 28(1987), 241-249.

Matematický ústav UK, Sokolovská 83, 18600 Praha 8, Czechoslovakia

(Oblatum 3.9. 1987)