

Mouffak Benchohra; Johnny Henderson; Sotiris K. Ntouyas  
On first order impulsive semilinear functional differential inclusions

*Archivum Mathematicum*, Vol. 39 (2003), No. 2, 129--139

Persistent URL: <http://dml.cz/dmlcz/107859>

## Terms of use:

© Masaryk University, 2003

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

**ON FIRST ORDER IMPULSIVE SEMILINEAR  
FUNCTIONAL DIFFERENTIAL INCLUSIONS**

M. BENCHOHRA, J. HENDERSON AND S. K. NTOUYAS

ABSTRACT. In this paper the Leray-Schauder nonlinear alternative for multivalued maps combined with the semigroup theory is used to investigate the existence of mild solutions for first order impulsive semilinear functional differential inclusions in Banach spaces.

1. INTRODUCTION

This paper is concerned with the existence of mild solutions for the impulsive semilinear functional differential inclusion of the form:

$$(1.1) \quad y' - Ay \in F(t, y_t), \quad t \in J = [0, b], \quad t \neq t_k, \quad k = 1, \dots, m,$$

$$(1.2) \quad y(t_k^+) = I_k(y(t_k^-)), \quad k = 1, \dots, m,$$

$$(1.3) \quad y_0 = \phi,$$

where  $F : J \times D \rightarrow 2^E$  is a closed, bounded and convex valued multivalued map  $D = \{\psi : [-r, 0] \rightarrow E \mid \psi \text{ is continuous everywhere except for a finite number of points } s \text{ at which } \psi(s^-) \text{ and } \psi(s^+) \text{ exist and } \psi(s^-) = \psi(s)\}$ ,  $\phi \in C([-r, 0], E)$ ,  $A$  is the infinitesimal generator of a strongly continuous semigroup  $T(t)$ ,  $t \geq 0$  and  $E$  a real Banach space with the norm  $|\cdot|$ ,  $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = b$ ,  $I_k \in C(E, E)$  ( $k = 1, 2, \dots, m$ ),  $y(t_k^-)$  and  $y(t_k^+)$  represent the left and right limits of  $y(t)$  at  $t = t_k$ , respectively.

For any continuous function  $y$  defined on the interval  $[-r, b] - \{t_1, \dots, t_m\}$  and any  $t \in J$ , we denote by  $y_t$  the element of  $C([-r, 0], E)$  defined by

$$y_t(\theta) = y(t + \theta), \quad \theta \in [-r, 0].$$

Here  $y_t(\cdot)$  represents the history of the state from time  $t - r$ , up to the present time  $t$ .

---

2000 *Mathematics Subject Classification*: 34A37, 34A60, 34G20, 35R10.

*Key words and phrases*: initial value problem, impulsive functional differential inclusions, convex multivalued map, fixed point, mild solution.

Received April 19, 2001.

Impulsive differential equations have become more important in recent years in some mathematical models of real world phenomena, especially in the biological or medical domain (see the monographs of Lakshmikantham et al. [10], and Samoilenko and Perestyuk [16], and the papers of Agur et al. [1], Erbe et al. [6], Goldbeter et al. [7], Kirane and Rogovchenko [9], Liu et al. [12] and Liu and Zhang [13]).

This paper will be divided into three sections. In Section 2 we will recall briefly some basic definitions and preliminary facts from multivalued analysis which will be used throughout Section 3. In Section 3 we establish an existence theorem for (1.1)-(1.3). Our approach is based on the nonlinear alternative of Leray-Schauder type for multivalued maps combined with the semigroup theory [15].

In our results we do not assume any type of monotonicity condition on  $I_k, k = 1, \dots, m$  which is usually the situation in the literature, see for instance, [6], [9] and [12].

## 2. PRELIMINARIES

We will briefly recall some basic definitions and facts from multivalued analysis that we will use in the sequel. Let the Banach space  $E$ .  $B(E)$  denotes the Banach space of bounded linear operators from  $E$  into  $E$ .

A measurable function  $y : J \rightarrow E$  is Bochner integrable if and only if  $|y|$  is Lebesgue integrable. For properties of the Bochner integral, we refer to Yosida [17].

$L^1(J, E)$  denotes the Banach space of functions  $y : J \rightarrow E$  which are Bochner integrable normed by

$$\|y\|_{L^1} = \int_0^b |y(t)| dt \quad \text{for all } y \in L^1(J, E).$$

Let  $(X, \|\cdot\|)$  be a Banach space. A multivalued map  $G : X \rightarrow 2^X$  is convex (closed) valued if  $G(x)$  is convex (closed) for all  $x \in X$ .  $G$  is bounded on bounded sets if  $G(B) = \cup_{x \in B} G(x)$  is bounded in  $X$  for any bounded set  $B$  of  $X$  (i.e.  $\sup_{x \in B} \{\sup\{\|y\| : y \in G(x)\}\} < \infty$ ).

$G$  is called upper semicontinuous (u.s.c.) on  $X$  if for each  $x_* \in X$  the set  $G(x_*)$  is a nonempty, closed subset of  $X$ , and if for each open set  $B$  of  $X$  containing  $G(x_*)$ , there exists an open neighbourhood  $V$  of  $x_*$  such that  $G(V) \subseteq B$ .

$G$  is said to be completely continuous if  $G(B)$  is relatively compact for every bounded subset  $B \subseteq X$ .

If the multivalued map  $G$  is completely continuous with nonempty compact values, then  $G$  is u.s.c. if and only if  $G$  has a closed graph (i.e.  $x_n \rightarrow x_*, y_n \rightarrow y_*, y_n \in G(x_n)$  imply  $y_* \in G(x_*)$ ).  $G$  has a fixed point if there is  $x \in X$  such that  $x \in G(x)$ .

In the following  $BCC(X)$  denotes the set of all nonempty bounded, closed and convex subsets of  $X$ .

A multivalued map  $G : J \rightarrow BCC(E)$  is said to be measurable if for each  $x \in E$  the function  $Y : J \rightarrow \mathbb{R}$  defined by

$$Y(t) = d(x, G(t)) = \inf\{|x - z| : z \in G(t)\}$$

is measurable. For more details on multivalued maps see the books of Deimling [4] and Hu and Papageorgiou [8].

**Definition 2.1.** A multivalued map  $F : J \times D \rightarrow 2^E$  is said to be  $L^1$ -Carathéodory if

- (i)  $t \mapsto F(t, u)$  is measurable for each  $u \in D$ ;
- (ii)  $u \mapsto F(t, u)$  is upper semicontinuous for almost all  $t \in J$ ;
- (iii) For each  $\rho > 0$ , there exists  $\phi_\rho \in L^1(J, \mathbb{R}_+)$  such that  $\|F(t, u)\| = \sup\{|v| : v \in F(t, u)\} \leq \phi_\rho(t)$  for all  $\|u\| \leq \rho$  and for almost all  $t \in J$ .

In order to define the mild solution of (1.1)-(1.3) we shall consider the following space

$$\Omega = \{y : [-r, b] \rightarrow E : y_k \in C(J_k, E), k = 0, \dots, m \text{ and there exist } y(t_k^-) \text{ and } y(t_k^+), k = 1, \dots, m \text{ with } y(t_k^-) = y(t_k) \text{ with } y(t) = \phi(t), \forall t \in [-r, 0]\}$$

which is a Banach space with the norm

$$\|y\|_\Omega = \max\{\|y_k\|_\infty, k = 0, \dots, m\},$$

where  $y_k$  is the restriction of  $y$  to  $J_k = [t_k, t_{k+1}]$ ,  $k = 0, \dots, m$ .

So let us start by defining what we mean by a mild solution of problem (1.1)-(1.3).

**Definition 2.2.** A function  $y \in \Omega$  is said to be a mild solution of (1.1)-(1.3) (see [15]) if there exists a function  $v \in L^1(J, E)$  such that  $v(t) \in F(t, y_t)$  a.e. on  $J_k$ , and

$$y(t) = \begin{cases} \phi(t), & t \in [-r, 0] \\ T(t)\phi(0) + \int_0^t T(t-s)v(s) ds, & t \in [0, t_1], \\ I_k(y(t_k^-)) + \int_{t_k}^t T(t-s)v(s) ds, & t \in J_k, k = 1, \dots, m. \end{cases}$$

For the multivalued map  $F$  and for each  $y \in C([-r, b], E)$  we define  $S_{F,y}^1$  by

$$S_{F,y}^1 = \{v \in L^1(J, E) : v(t) \in F(t, y_t) \text{ for a.e. } t \in J\}.$$

Our main result is based on the following:

**Lemma 2.3** [11]. *Let  $I$  be a compact real interval and  $X$  be a Banach space. Let  $F : I \times X \rightarrow BCC(X); (t, y) \rightarrow F(t, y)$  measurable with respect to  $t$  for any  $y \in X$  and u.s.c. with respect to  $y$  for almost each  $t \in I$  and  $S_{F,y}^1 \neq \emptyset$  for any  $y \in C(I, X)$  and let  $\Gamma$  be a linear continuous mapping from  $L^1(I, X)$  to  $C(I, X)$ . Then the operator*

$$\Gamma \circ S_F^1 : C(I, X) \rightarrow BCC(C(I, X)), \quad y \mapsto (\Gamma \circ S_F^1)(y) := \Gamma(S_{F,y}^1)$$

*is a closed graph operator in  $C(I, X) \times C(I, X)$ .*

**Lemma 2.4** (Nonlinear Alternative [5]). *Let  $X$  be a Banach space with  $C \subset X$  convex. Assume  $U$  is a relatively open subset of  $C$  with  $0 \in U$  and  $G : \bar{U} \rightarrow 2^C$  is a compact multivalued map, u.s.c. with convex closed values. Then either,*

- (i)  $G$  has a fixed point in  $\bar{U}$ ; or
- (ii) there is a point  $u \in \partial U$  and  $\lambda \in (0, 1)$  with  $u \in \lambda G(u)$ .

**Remark 2.5.** By  $\bar{U}$  and  $\partial U$  we denote the closure of  $U$  and the boundary of  $U$  respectively.

### 3. MAIN RESULT

We are now in a position to state and prove our existence result for the IVP (1.1)-(1.3).

**Theorem 3.1.** *Let  $t_0 = 0, t_{m+1} = b$ . Suppose:*

- (H1)  $A$  is the infinitesimal generator of a linear bounded compact semigroup  $T(t), t \geq 0$  and there exists  $M \geq 1$  such that  $|T(t)|_{B(E)} \leq M$ ;
- (H2)  $F : J \times D \rightarrow BCC(E)$  is an  $L^1$ -Carathéodory multivalued map and for each fixed  $y \in C([-r, b], E)$  the set

$$S_{F,y}^1 = \{v \in L^1(J, E) : v(t) \in F(t, y_t) \quad \text{for a.e. } t \in J\}$$

*is nonempty;*

- (H3) there exists a continuous nondecreasing function  $\psi : [0, \infty) \rightarrow (0, \infty)$  and  $p \in L^1(J, \mathbb{R}_+)$  such that

$$\|F(t, u)\| := \sup\{|v| : v \in F(t, u)\} \leq p(t)\psi(\|u\|)$$

*for a.e.  $t \in J$  and each  $u \in D$  with*

$$\int_{t_{k-1}}^{t_k} p(s) ds < \int_{N_{k-1}}^{\infty} \frac{d\tau}{\psi(\tau)}, \quad k = 1, \dots, m + 1,$$

*where  $N_0 = M\|\phi\|$ , and for  $k = 2, \dots, m + 1$ ,*

$$N_{k-1} = \sup_{y \in [-M_{k-2}, M_{k-2}]} |I_{k-1}(y)|, \quad M_{k-2} = \Gamma_{k-1}^{-1} \left( M \int_{t_{k-2}}^{t_{k-1}} p(s) ds \right)$$

*with*

$$\Gamma_l(z) = \int_{N_{l-1}}^z \frac{d\tau}{\psi(\tau)}, \quad z \geq N_{l-1}, \quad l \in \{1, \dots, m + 1\}.$$

Then the problem (1.1)-(1.3) has at least one mild solution  $y \in \Omega$ .

**Remark 3.2.** (i) If  $\dim E < \infty$ , then for each  $y \in C([-r, b], E)$ ,  $S_{F,y}^1 \neq \emptyset$  (see Lasota and Opial [11]).

(ii) If  $\dim E = \infty$  and  $y \in C([-r, b], E)$  the set  $S_{F,y}^1$  is nonempty if and only if the function  $Y : J \rightarrow \mathbb{R}$  defined by

$$Y(t) := \inf\{|v| : v \in F(t, y_t)\}$$

belongs to  $L^1(J, \mathbb{R})$  (see Hu and Papageorgiou [8]).

**Proof.** The proof is given in several steps.

**Step 1.** Consider the problem (1.1)-(1.3) on  $[-r, t_1]$

$$(3.1) \quad y' - Ay \in F(t, y_t), \quad \text{a.e. } t \in J_0,$$

$$(3.2) \quad y_0 = \phi.$$

We shall show that the possible mild solutions of (3.1)-(3.2) are *a priori* bounded, that is, there exists a constant  $b_0$  such that, if  $y \in \Omega$  is a mild solution of (3.1)-(3.2), then

$$\sup\{|y(t)| : t \in [-r, 0] \cup J_0\} \leq b_0.$$

Let  $y$  be a (possible) mild solution to (3.1)-(3.2). Then for each  $t \in [0, t_1]$

$$y(t) - T(t)\phi(0) \in \int_0^t T(t-s)F(s, y_s) ds.$$

From (H3) we get

$$|y(t)| \leq M\|\phi\| + M \int_0^t p(s)\psi(\|y_s\|) ds, \quad t \in [0, t_1].$$

We consider the function  $\mu_0$  defined by

$$\mu_0(t) = \sup\{|y(s)| : -r \leq s \leq t\}, \quad 0 \leq t \leq t_1.$$

Let  $t^* \in [-r, t]$  be such that  $\mu_0(t) = |y(t^*)|$ . If  $t^* \in [0, t_1]$ , by the previous inequality we have for  $t \in [0, t_1]$

$$\mu_0(t) \leq M\|\phi\| + M \int_0^t p(s)\psi(\mu_0(s)) ds.$$

If  $t^* \in [-r, 0]$  then  $\mu_0(t) = \|\phi\|$  and the previous inequality holds since  $M \geq 1$ .

Let us take the right-hand side of the above inequality as  $v_0(t)$ , then we have

$$v_0(0) = M\|\phi\| = N_0, \quad \mu_0(t) \leq v_0(t), \quad t \in [0, t_1]$$

and

$$v'_0(t) = Mp(t)\psi(\mu_0(t)), \quad t \in [0, t_1].$$

Using the nondecreasing character of  $\psi$  we get

$$v'_0(t) \leq Mp(t)\psi(v_0(t)), \quad t \in [0, t_1].$$

This implies for each  $t \in [0, t_1]$  that

$$\int_{N_0}^{v_0(t)} \frac{d\tau}{\psi(\tau)} \leq M \int_0^{t_1} p(s) ds.$$

In view of (H3), we obtain

$$|v_0(t^*)| \leq \Gamma_1^{-1} \left( M \int_0^{t_1} p(s) ds \right) := M_0.$$

Since for every  $t \in [0, t_1]$ ,  $\|y_t\| \leq \mu_0(t)$ , we have

$$\sup_{t \in [-r, t_1]} |y(t)| \leq \max(\|\phi\|, M_0) = b_0.$$

We transform this problem into a fixed point problem. A mild solution to (3.1)-(3.2) is a fixed point of the operator  $G : C([-r, t_1], E) \rightarrow 2^{C([-r, t_1], E)}$  defined by

$$G(y) := \left\{ h \in C([-r, t_1], E) : h(t) = \begin{cases} \phi(t), & t \in [-r, 0] \\ T(t)\phi(0) \\ + \int_0^t T(t-s)v(s) ds, & t \in J_0 \end{cases} \right\}$$

where  $v \in S_{F,y}^1$ . We shall show that  $G$  satisfies the assumptions of Lemma 2.2.

**Claim 1:**  $G(y)$  is convex for each  $y \in C(J_0, E)$ .

Indeed, if  $h, \bar{h}$  belong to  $G(y)$ , then there exist  $v \in S_{F,y}^1$  and  $\bar{v} \in S_{F,y}^1$  such that

$$h(t) = T(t)\phi(0) + \int_0^t T(t-s)v(s) ds, \quad t \in J_0$$

and

$$\bar{h}(t) = T(t)\phi(0) + \int_0^t T(t-s)\bar{v}(s) ds, \quad t \in J_0.$$

Let  $0 \leq l \leq 1$ . Then for each  $t \in J_0$  we have

$$[lh + (1-l)\bar{h}](t) = T(t)\phi(0) + \int_0^t T(t-s)[lv(s) + (1-l)\bar{v}(s)] ds.$$

Since  $S_{F,y}^1$  is convex (because  $F$  has convex values) then

$$lh + (1 - l)\bar{h} \in G(y).$$

**Claim 2:**  $G$  sends bounded sets into bounded sets in  $C(J_0, E)$ .

Let  $B_q := \{y \in C(J_0, E) : \|y\|_\infty = \sup_{t \in J_0} |y(t)| \leq q\}$  be a bounded set in  $C(J_0, E)$  and  $y \in B_q$ , then for each  $h \in G(y)$  there exists  $v \in S_{F,y}^1$  such that

$$h(t) = T(t)\phi(0) + \int_0^t T(t-s)v(s) ds, \quad t \in J_0.$$

Thus for each  $t \in J_0$  we get

$$\begin{aligned} |h(t)| &\leq M\|\phi\| + M \int_0^t |v(s)| ds \\ &\leq M\|\phi\| + M\|\phi_q\|_{L^1}. \end{aligned}$$

**Claim 3:**  $G$  sends bounded sets in  $C(J_0, E)$  into equicontinuous sets.

Let  $r_1, r_2 \in J_0$ ,  $r_1 < r_2$ ,  $B_q := \{y \in C(J_0, E) : \|y\|_\infty \leq q\}$  be a bounded set in  $C(J_0, E)$  as in Claim 2 and  $y \in B_q$ . For each  $h \in G(y)$  there exists  $v \in S_{F,y}^1$  such that

$$h(t) = T(t)\phi(0) + \int_0^t T(t-s)v(s) ds, \quad t \in J_0.$$

Hence,

$$\begin{aligned} |h(r_2) - h(r_1)| &\leq |(T(r_2)\phi(0) - T(r_1)\phi(0))| + \left| \int_0^{r_2} [T(r_2-s) - T(r_1-s)]v(s) ds \right| \\ &\quad + \left| \int_{r_1}^{r_2} T(r_1-s)v(s) ds \right| \\ &\leq |(T(r_2)\phi(0) - T(r_1)\phi(0))| + \left| \int_0^{r_2} [T(r_2-s) - T(r_1-s)]v(s) ds \right| \\ &\quad + M \int_{r_1}^{r_2} |v(s)| ds \\ &\leq |(T(r_2)\phi(0) - T(r_1)\phi(0))| + \left| \int_0^{r_2} [T(r_2-s) - T(r_1-s)]\phi_r(s) ds \right| \\ &\quad + M \int_{r_1}^{r_2} |\phi_r(s)| ds. \end{aligned}$$

The equicontinuity for the cases  $r_1 < r_2 \leq 0$  and  $r_1 \leq 0 \leq r_2$  are obvious.



Set

$$U = \{y \in C([-r, t_1], E) : \|y\|_\infty < b_0 + 1\}.$$

As a consequence of Claim 2 and Claim 3 together with the Arzela-Ascoli theorem we can conclude that  $G : \overline{U} \rightarrow 2^{C([-r, t_1], E)}$  is a compact multivalued map.

**Claim 4:**  $G$  has a closed graph.

Let  $y_n \rightarrow y_*$ ,  $h_n \in G(y_n)$  and  $h_n \rightarrow h_*$ . We shall prove that  $h_* \in G(y_*)$ .  $h_n \in G(y_n)$  means that there exists  $v_n \in S_{F, y_n}$  such that

$$h_n(t) = T(t)\phi(0) + \int_0^t T(t-s)v_n(s) ds, \quad t \in J_0.$$

We must prove that there exists  $v_* \in S_{F, y_*}^1$  such that

$$h_*(t) = T(t)\phi(0) + \int_0^t T(t-s)v_*(s) ds, \quad t \in J_0.$$

Consider the linear continuous operator  $\Gamma : L^1(J_0, E) \rightarrow C(J_0, E)$  defined by

$$(\Gamma v)(t) = \int_0^t T(t-s)v(s) ds.$$

We have

$$\|(h_n - T(t)\phi(0)) - (h_* - T(t)\phi(0))\|_\infty \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

From Lemma 2.1 it follows that  $\Gamma \circ S_F^1$  is a closed graph operator. Also from the definition of  $\Gamma$  we have that

$$h_n(t) - T(t)\phi(0) \in \Gamma(S_{F, y_n}^1).$$

Since  $y_n \rightarrow y_*$ , it follows from Lemma 2.1 that

$$h_*(t) = T(t)\phi(0) + \int_0^t T(t-s)v_*(s) ds, \quad t \in J_0$$

for some  $v_* \in S_{F, y_*}^1$ .

From the choice of  $U$  there is no  $y \in \partial U$  such that  $y \in \lambda G(y)$  for any  $\lambda \in (0, 1)$ .

As a consequence of Lemma 2.2 we deduce that  $G$  has a fixed point  $y_0 \in \overline{U}$  which is a mild solution of (3.1)–(3.2).

**Step 2.** Consider now the following problem on  $J_1 := [t_1, t_2]$

$$(3.3) \quad y' - Ay \in F(t, y_t), \quad \text{a.e. } t \in J_1,$$

$$(3.4) \quad y(t_1^+) = I_1(y(t_1^-)).$$

Let  $y$  be a (possible) mild solution to (3.3)–(3.4). Then for each  $t \in [t_1, t_2]$

$$y(t) - I_1(y(t_1^-)) \in \int_{t_1}^t T(t-s)F(s, y_s) ds.$$

From (H3) we get

$$|y(t)| \leq \sup_{t \in [-r, t_1]} |I_1(y_0(t^-))| + M \int_{t_1}^t p(s)\psi(\|y_s\|) ds, \quad t \in [t_1, t_2].$$

We consider the function  $\mu_1$  defined by

$$\mu_1(t) = \sup\{|y(s)| : t_1 \leq s \leq t\}, \quad t_1 \leq t \leq t_2.$$

Let  $t^* \in [t_1, t]$  be such that  $\mu_1(t) = |y(t^*)|$ . Then we have for  $t \in [t_1, t_2]$

$$\mu_1(t) \leq N_1 + M \int_{t_1}^t p(s)\psi(\mu_1(s)) ds.$$

Let us take the right-hand side of the above inequality as  $v_1(t)$ , then we have

$$v_1(t_1) = N_1, \quad \mu_1(t) \leq v_1(t), \quad t \in [t_1, t_2]$$

and

$$v_1'(t) = Mp(t)\psi(\mu_1(t)), \quad t \in [t_1, t_2].$$

Using the nondecreasing character of  $\psi$  we get

$$v_1'(t) \leq Mp(t)\psi(v_1(t)), \quad t \in [t_1, t_2].$$

This implies for each  $t \in [t_1, t_2]$  that

$$\int_{N_1}^{v_1(t)} \frac{d\tau}{\psi(\tau)} \leq M \int_{t_1}^{t_2} p(s) ds.$$

In view of (H3), we obtain

$$|v_1(t^*)| \leq \Gamma_2^{-1} \left( M \int_{t_1}^{t_2} p(s) ds \right) := M_1.$$

Since for every  $t \in [t_1, t_2]$ ,  $\|y_t\| \leq \mu_1(t)$ , we have

$$\sup_{t \in [t_1, t_2]} |y(t)| \leq M_1.$$

A mild solution to (3.3)-(3.4) is a fixed point of the operator  $G : C(J_1, E) \rightarrow 2^{C(J_1, E)}$  defined by

$$G(y) := \left\{ h \in C(J_1, E) : h(t) = I_1(y(t_1^-)) + \int_{t_1}^t T(t-s)v(s) ds : v \in S_{F,y}^1 \right\} .$$

Set

$$U = \{y \in C(J_1, E) : \|y\|_\infty < M_1 + 1\} .$$

As in Step 1 we can show that  $G : \bar{U} \rightarrow 2^{C(J_1, E)}$  is a compact multivalued map and u.s.c. From the choice of  $U$  there is no  $y \in \partial U$  such that  $y \in \lambda G(y)$  for any  $\lambda \in (0, 1)$ .

As a consequence of Lemma 2.2 we deduce that  $G$  has a fixed point  $y_1 \in \bar{U}$  which is a mild solution of (3.3)-(3.4).

**Step 3.** Continue this process and construct solutions  $y_k \in C(J_k, E), k = 2, \dots, m$  to

$$(3.5) \quad y' - Ay \in F(t, y_t), \quad \text{a.e. } t \in J_k,$$

$$(3.6) \quad y(t_k^+) = I_k(y(t_k^-)) .$$

Then

$$y(t) = \begin{cases} y_0(t), & t \in [-r, t_1] \\ y_1(t), & t \in (t_1, t_2] \\ \vdots \\ y_{m-1}(t), & t \in (t_{m-1}, t_m] \\ y_m(t), & t \in (t_m, b] \end{cases}$$

is a mild solution of (1.1)-(1.3). □

### REFERENCES

- [1] Agur, Z., Cojocaru, L., Mazur, G., Anderson, R.M. and Danon, Y.L., *Pulse mass measles vaccination across age cohorts*, Proc. Nat. Acad. Sci. USA **90** (1993), 11698–11702.
- [2] Banas, J. and Goebel, K., *Measures of Noncompactness in Banach Spaces*, Marcel Dekker, New York, 1980.
- [3] Corduneanu, C., *Integral Equations and Applications*, Cambridge Univ. Press, New York, 1990.
- [4] Deimling, K., *Multivalued Differential Equations*, De Gruyter, Berlin, 1992.
- [5] Dugundji, J. and Granas, A., *Fixed Point Theory*, Monografie Mat. PWN, Warsaw, 1982.
- [6] Erbe, L., Freedman, H.I., Liu, X.Z. and Wu, J.H., *Comparison principles for impulsive parabolic equations with applications to models of singles species growth*, J. Austral. Math. Soc. Ser. B **32** (1991), 382–400.
- [7] Goldbeter, A., Li, Y. X. and Dupont, G., *Pulsatile signalling in intercellular communication: Experimental and theoretical aspects*, Mathematics applied to Biology and Medicine, Werz. Pub. Winnipeg, Canada (1993), 429–439.

- [8] Hu, Sh. and Papageorgiou, N., *Handbook of Multivalued Analysis, Volume I: Theory*, Kluwer, Dordrecht, Boston, London, 1997.
- [9] Kirane, M. and Rogovchenko, Y. V., *Comparison results for systems of impulsive parabolic equations with applications to population dynamics*, *Nonlinear Anal.* **28** (1997), 263–276.
- [10] Lakshmikantham, V., Bainov, D. D. and Simeonov, P. S., *Theory of Impulsive Differential Equations*, World Scientific, Singapore, 1989.
- [11] Lasota, A. and Opial, Z., *An application of the Kakutani-Ky-Fan theorem in the theory of ordinary differential equations*, *Bull. Polish Acad. Sci., Ser. Sci. Math. Astronom. Phys.* **13** (1965), 781–786.
- [12] Liu, X., Sivaloganathan, S. and Zhang, S., *Monotone iterative techniques for time-dependent problems with applications*, *J. Math. Anal. Appl.* **237** (1999), 1–18.
- [13] Liu, X. and Zhang, S., *A cell population model described by impulsive PDEs-Existence and numerical approximation*, *Comput. Math. Appl.* **36** (8) (1998), 1–11.
- [14] Martelli, M., *A Rothe's type theorem for non compact acyclic-valued maps*, *Boll. Un. Mat. Ital.* **11** (Suppl. Fasc.) (1975), 70–76.
- [15] Pazy, A., *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, New York, 1983.
- [16] Samoilenko, A. M. and Perestyuk, N. A., *Impulsive Differential Equations*, World Scientific, Singapore, 1995.
- [17] Yosida, K., *Functional Analysis*, 6<sup>th</sup> edn. Springer-Verlag, Berlin, 1980.

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE SIDI BEL ABBÈS  
 BP 89, 22000 SIDI BEL ABBÈS, ALGÉRIE  
*E-mail:* [benchohra@univ.sba.dz](mailto:benchohra@univ.sba.dz)

DEPARTMENT OF MATHEMATICS, BAYLOR UNIVERSITY  
 WACO, TEXAS 76798-7328, USA  
*E-mail:* [Johny\\_Henderson@baylor.edu](mailto:Johny_Henderson@baylor.edu)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOANNINA  
 451 10 IOANNINA, GREECE  
*E-mail:* [sntouyas@cc.uoi.gr](mailto:sntouyas@cc.uoi.gr)