

Robert El Bashir; Tomáš Kepka  
Notes on slender prime rings

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 37 (1996), No. 2, 419--422

Persistent URL: <http://dml.cz/dmlcz/118847>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Notes on slender prime rings

ROBERT EL BASHIR, TOMÁŠ KEPKA

*Abstract.* If  $R$  is a prime ring such that  $R$  is not completely reducible and the additive group  $R(+)$  is not complete, then  $R$  is slender.

*Keywords:* ring, prime, slender

*Classification:* 16N60

The purpose of this short note is to discuss a few sufficient conditions for a prime ring to be slender. As concerns the concept of slenderness (various results, references, historical remarks, etc.), a reader is fully referred to [4, Chapter III].

### 1. Introduction

In the sequel,  $R$  is a non-zero associative ring with unit and modules are unitary left  $R$ -modules. The ring  $R$  is said to be prime (resp. a domain) if  $aRb \neq 0$  (resp.  $ab \neq 0$ ) for all  $a, b \in R$ ,  $a \neq 0 \neq b$ . Commutative domains are also called integral domains.

Let  $M$  be a module. By a filtration  $\mathcal{F}$  of  $M$  we mean any sequence  $M_i$ ,  $i < \omega$ , of submodules of  $M$  such that  $M_i \supseteq M_{i+1}$ . The filtration  $\mathcal{F}$  is said to be separating if  $\bigcap_{\mathcal{F}} M_i = 0$  and it is said to be discrete if  $0 \in \mathcal{F}$ . The filtration  $\mathcal{F}$  determines a linear closure operator on  $M$  and the module  $M$  is said to be  $\mathcal{F}$ -complete if every Cauchy  $\mathcal{F}$ -sequence of elements from  $M$  is convergent.

A module  $M$  is said to be complete if it is  $\mathcal{F}$ -complete for a non-discrete separating filtration  $\mathcal{F}$  of  $M$ .

A left (right) ideal  $I$  of  $R$  is said to be l. s.  $\cup$ -compact (r. s.  $\cup$ -compact) if every countable subset  $S$  of  $I$  is contained in a finitely generated left (right) ideal  $K \subseteq I$ .

The ring  $R$  is said to be left (right)  $\cap$ -compact if the left (right) module  ${}_R R$  ( $R_R$ ) possesses no non-discrete separating filtration.

We denote by  $\mathcal{T}_R$  the set of ideals  $I$  of  $R$  such that the factor  $R/I$  is completely reducible. For a module  $M$ , let  $Soct(M)$  be the set of all  $x \in M$  such that  $(0 : x)$  contains an ideal from  $\mathcal{T}_R$ . Finally, let  $V = R^\omega$ ,  $U = R^{(\omega)}$  and  $W = V/U$ . If  $i < \omega$ , then  $V[i] = \{a \in V; a(j) = 0 \text{ for every } j < i\}$ .

For further basic terminology concerning rings and modules, we refer to [1].

---

This research has been partially supported by the Grant Agency of the Czech Republic, grant #GAČR-2433

**2. Slender modules**

A module  $M$  is said to be slender if, for every homomorphism  $\varphi : V \rightarrow M$ ,  $\varphi(e_i) = 0$  for almost all  $i < \omega$ . The following result is implicitly contained in [5] and is proved in [3] for torsionfree modules over integral domains:

**2.1 Proposition.** *A module  $M$  is slender if and only if  $\text{Hom}_R(W, M) = 0$  and  $M$  is not complete.*

**2.2 Proposition.** *Let  $M$  be a module such that there exists a filtration  $I_i, i < \omega$ , of  $R$  satisfying the following conditions:*

- (1)  $I_i$  is a r. s.  $\cup$ -compact ideal for every  $i < \omega$ .
- (2) If  $i < \omega$  and  $0 \neq u \in M$ , then  $I_i u \neq 0$ .
- (3)  $\bigcap_{\omega} I_i M = 0$ .

*Then the module  $M$  is slender if and only if it is not complete.*

PROOF: The result is an immediate consequence of the following observation:

□

**2.3 Observation.** Let  $I_i, i < \omega$ , be a filtration of  $R_R$  such that all the right ideals  $I_i$  are r. s.  $\cup$ -compact. Put  $\mathcal{E} = \{I_i V[i]; i < \omega\}$ . Then  $\mathcal{E}$  is a separating filtration of  $V(+)$  and  $V(+)$  is  $\mathcal{E}$ -complete.

Now, let  $\varphi : V \rightarrow M$  be a (module) homomorphism. Put  $\mathcal{G} = \varphi(\mathcal{E}) = \{I_i \varphi(V[i])\}$ . Then  $\mathcal{G}$  is a filtration of  $M(+)$ ,  $\varphi$  is continuous and  $M(+)$  is  $\mathcal{G}$ -complete.

Assume  $\bigcap \mathcal{G} = 0$ . Then  $\text{Ker}(\varphi)$  is  $\mathcal{E}$ -closed in  $V$ . If  $\text{Ker}(\varphi)$  is  $\mathcal{E}$ -open, then  $I_m \varphi(V[m]) = 0$  for some  $m < \omega$ . If  $\text{Ker}(\varphi)$  is not  $\mathcal{E}$ -open, then  $\mathcal{G}$  is not discrete. Now,  $\mathcal{G}$  is a non-discrete separating filtration of  $M(+)$  and  $M(+)$  is  $\mathcal{G}$ -complete. In particular, if the right ideals  $I_i$  are two-sided, then  $M$  is a complete module.

**2.4 Corollary.** *Suppose that there exists a separating filtration  $I_i, i < \omega$ , of  $R$  such that  $I_i$  is an r. s.  $\cup$ -compact ideal and  $(0 : I_i)_r = 0$  for every  $i < \omega$ . Then the ring  $R$  is left slender if and only if it is not left complete.*

**2.5 Corollary.** *Let  $M$  be a module such that there exists a countable non-empty set  $\mathcal{M}$  of submodules of  $M$  satisfying the following properties:*

- (1)  $\bigcap \mathcal{M} = 0$ .
- (2)  $(0 : M/N)$  is r. s.  $\cup$ -compact for every  $N \in \mathcal{M}$ .
- (3)  $(0 : M/N) \in \mathcal{T}_R$  for every  $N \in \mathcal{M}$ .

*Then  $M$  is slender if and only if  $\text{Soct}(M) = 0$  and  $M$  is not complete.*

**2.6 Corollary.** *Suppose that  $R$  possesses a countable non-empty set  $\mathcal{M}$  of maximal left ideals such that  $\bigcap \mathcal{M} = 0$ ,  $(0 : R/I) \in \mathcal{T}_R$  and  $(0 : R/I)$  is r. s.  $\cup$ -compact for every  $I \in \mathcal{M}$ . Then  $R$  is left slender if and only if  $\text{Soct}_l(R) = 0$  and  $R$  is not left complete.*

### 3. Prime rings and slenderness

**3.1 Theorem.** *Let  $R$  be a prime ring.*

- (i) *If every right ideal is an ideal and  $R$  is not right  $\cap$ -compact, then  $R$  is a domain and  $R$  is left slender if and only if  $R$  is not left complete.*
- (ii) *If the additive group  $R(+)$  is not complete, then  $R$  is slender if and only if  $R$  is not isomorphic to a (full) matrix ring over a division ring.*
- (iii) *If  $\text{card}(R) \geq 2^\omega$  and the additive group  $R(+)$  is not complete, then  $R$  is slender.*

PROOF: (i) Clearly,  $R$  is a right uniform domain, and hence there is a separating filtration  $r_i R$ ,  $i < \omega$ , of non-zero principal right ideals and it remains to apply 2.4.

(ii) Let  $p$  denote the characteristic of  $R$ . If  $p > 0$ , then  $\text{card}(R) < 2^\omega$  (since  $R(+)$  is not complete) and we can use [2, Theorem 4.1]. If  $p = 0$  and  $R(+)$  is reduced, then  $R(+)$  is slender (see [6]) and consequently  $R$  is also slender. Assume finally that  $p = 0$  and the divisible part  $Q(+)$  of  $R(+)$  is non-zero.

Obviously,  $Q$  is an ideal of  $R$  and the factorgroup  $R(+)/Q(+)$  is slender ([6]), and hence the factormodule  ${}_R R/Q$  is slender, too. Now, it remains to show that the module  ${}_R Q$  is slender. However, since  $Q(+)$  is not complete, we have  $\text{card}(Q) < 2^\omega$  and then we can proceed similarly as in the proof of [2, Theorem 4.1].

- (iii) This assertion follows easily from (ii). □

**3.2 Proposition.** *Let  $R$  be a domain satisfying maximal condition on principal left ideals and such that  $R$  is not a division ring and that every right ideal of  $R$  is an ideal. Then  $R$  is left slender if and only if  $R$  is not left complete.*

PROOF: Clearly,  $R$  is not right  $\cap$ -compact and the result follows from 3.1 (i). □

**3.3 Proposition.** *Let  $R$  be an integral domain, not a field, satisfying at least one of the following conditions:*

- (1)  *$R$  is noetherian.*
- (2)  *$R$  is a unique factorization domain.*
- (3) *The quotient field of  $R$  is a countably generated  $R$ -module (see [3, Theorem 20]).*
- (4)  *$R$  is not  $\cap$ -compact.*

*Then  $R$  is slender if and only if it is not complete.*

PROOF: The first two cases follow from 3.2, the condition (3) implies (4) and, when (4) is true, the result follows from 3.1 (i). □

#### REFERENCES

- [1] Anderson F.W., Fuller K.R., *Rings and Categories of Modules*, 2<sup>nd</sup> edition, Springer, New York, 1992.
- [2] El Bashir R., Kepka T., *On when small semiprime rings are slender*, to appear.

- [3] Dimitrić R., *Slender modules over domains*, Commun. in Algebra **11** (1983), 1685–1700.
- [4] Eklof P., Mekler A., *Almost Free Modules*, North-Holland, New York, 1990.
- [5] Heinlein G., *Vollreflexive Ringe und schlanke Moduln*, Dissertation, Erlangen, 1971.
- [6] Nunke R., *Slender groups*, Acta Sci. Math. Szeged **23** (1962), 67–73.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY,  
SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECH REPUBLIC

*E-mail:* bashir@karlin.mff.cuni.cz      kepka@karlin.mff.cuni.cz

(Received May 4, 1995)