

Murray G. Bell

A compact ccc non-separable space from a Hausdorff gap and Martin's Axiom

Commentationes Mathematicae Universitatis Carolinae, Vol. 37 (1996), No. 3, 589--594

Persistent URL: <http://dml.cz/dmlcz/118865>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1996

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A compact ccc non-separable space from a Hausdorff gap and Martin’s Axiom

MURRAY BELL

Abstract. We answer a question of I. Juhász by showing that $MA + \neg CH$ does not imply that every compact ccc space of countable π -character is separable. The space constructed has the additional property that it does not map continuously onto I^{ω_1} .

Keywords: ccc, non-separable, Hausdorff gap, π -character

Classification: Primary 54D30, 54G20; Secondary 54A25, 54A35

1. Introduction

I. Juhász [Ju71] has proven that $MA(\omega_1)$ implies that every first countable compact ccc space is separable. This has been extended by Shapirovskii [Sh72] by replacing first countable with countable tightness. In Juhász [Ju77], the question is raised whether tightness can be replaced by π -character, i.e., whether $MA(\omega_1)$ implies that every compact ccc space of countable π -character is separable. We will show not. We present our space as a space whose points are certain ideals because this is the way that we found it; although the inclined reader should easily be able to identify the base set as a rather simple subset of $2^\omega \times \kappa$ (where κ is a certain regular cardinal $> \omega_1$) using a Hausdorff gap as a parameter.

2. General theory of total ideal spaces

Let $P = \bigcup_{A \subset \omega} 2^A$ and put $p \preceq q$ if q extends p . Then (P, \preceq) is a Dedekind complete partially ordered set. A subset F of P is **compatible** if $\bigcup F \in P$. We write $p \parallel q$ if $\{p, q\}$ is compatible and we write $p \perp q$ if $\{p, q\}$ is not compatible. A subset Q of P is **closed** in P if whenever F is a finite compatible subset of Q , then $\bigcup F \in Q$. For Q closed in P , a compatible and closed subset I of Q is called a **total ideal** of Q if

- (a) $\bigcup I$ has domain all of ω and
- (b) $p \in I$ and $q \in Q$ with $q \preceq p$ implies $q \in I$.

Let $\text{Fin} = \{p \in P : \text{dom}(p) \text{ is finite}\}$. The parameter for these ideal spaces will be a closed subset Q of P with $\text{Fin} \subset Q$. For such a Q , put $X(Q) = \{I \subset Q : I \text{ is a total ideal of } Q\}$. It is seen that $X(Q)$ is a closed subspace of 2^Q (where

The author gratefully acknowledges support from NSERC of Canada

$2 = \{0, 1\}$ has the discrete topology, 2^Q has the product topology, and points of $X(Q)$ are identified with their characteristic functions). For $B \subset Q$, put $B^+ = \{I \in X(Q) : B \subset I\}$ and $B^- = \{I \in X(Q) : B \cap I = \emptyset\}$. If $B = \{q\}$, then we simply write q^+ and q^- . Then, since Q is closed, a base for $X(Q)$ consists of the clopen sets $q^+ \cap B^-$ where $q \in Q$ and B is a finite subset of $\{r \in Q : q \prec r\}$. We note the helpful facts that

- (a) $q^+ \subset r^+$ iff $r \preceq q$,
- (b) $q^+ \cap r^+ \neq \emptyset$ iff $q \parallel r$ iff $[q^{-1}(1) \cup r^{-1}(1)] \cap [q^{-1}(0) \cup r^{-1}(0)] = \emptyset$,
- (c) $\lambda : X(Q) \rightarrow 2^\omega$ given by $\lambda(I) = \bigcup I$ is a continuous surjection.

Put $\mathcal{Q} = \{q^+ : q \in Q\}$ and for each $f \in 2^\omega$, let M_f be the maximal total ideal $\{q \in Q : q \preceq f\}$.

Fact 2.1. \mathcal{Q} is a T_0 -separating, binary π -base of $X(Q)$. Hence, $\pi w(X(Q)) = cf(Q, \preceq) = \min\{|D| : D \subset Q \text{ and } \forall q \in Q : \exists d \in D : (q \preceq d)\}$.

This was proved in [Be88] and [Be89]. T_0 -separating and binary are straightforward. The fact that \mathcal{Q} is a π -base is crucial. The reader will see a proof of this in Lemma 3.2 where we must prove a little bit more in order to achieve countable π -character.

Now we show two facts which delineate the kinds of Souslinean examples that we can get from these ideal spaces.

Fact 2.2. If $X(Q)$ is σ -linked, then $X(Q)$ is separable.

If $X(Q)$ is σ -linked, then $\mathcal{Q} = \bigcup_{n < \omega} \mathcal{Q}_n$ where for each $n < \omega$, \mathcal{Q}_n is linked. Since \mathcal{Q} is binary, by choosing $I_n \in \bigcap \mathcal{Q}_n$ for each $n < \omega$, we get that $\{I_n : n < \omega\}$ is dense in $X(Q)$.

We refer the reader to Todorcevic [To89] for the definition of the Open Colouring Axiom OCA.

Fact 2.3 (OCA). If $X(Q)$ is ccc, then $X(Q)$ is separable.

For each $q \in Q$ put $A_q = q^{-1}(1)$ and $B_q = q^{-1}(0)$. Then A_q and B_q are disjoint subsets of ω . For each $q \in Q$ let a_q, b_q be the characteristic functions of A_q, B_q respectively. Let $S = \{(a_q, b_q) : q \in Q\}$ have the subspace topology from $2^\omega \times 2^\omega$. Define a partition of $[S]^2$ by $\{(a_q, b_q), (a_r, b_r)\} \in K_0$ iff $q^+ \cap r^+ = \emptyset$ iff $(A_q \cup A_r) \cap (B_q \cup B_r) \neq \emptyset$. K_0 is open in $[S]^2$. Since $X(Q)$ is ccc, there does not exist a K_0 -homogeneous subset of S which has cardinality ω_1 . Hence, by OCA, $\mathcal{Q} = \bigcup_{n < \omega} \mathcal{Q}_n$ where for every n and for every q, r in \mathcal{Q}_n , $q^+ \cap r^+ \neq \emptyset$, i.e., $\{q^+ : q \in \mathcal{Q}_n\}$ is linked. We get that \mathcal{Q} is σ -linked, hence Fact 2.2 implies that $X(Q)$ is separable.

Remarks: We have learned that Fact 2.3 follows from a more general result Theorem 10.3* in Todorcevic and Farah [TF95]. We see from Fact 2.3, that if we want a ccc but not separable space $X(Q)$, then we must be in a model of set theory contradicting OCA. We did this in [Be89] under CH producing a first

countable Corson compact space which was ccc but did not have Property K . We also point out that interesting separable spaces $X(Q)$ of uncountable π -weight can be achieved in every model of set theory. In [Be88], for each regular cardinal κ for which there exists a κ -chain of clopen sets in $\beta\omega \setminus \omega$, we produced a separable space $X(Q)$ of π -weight κ that did not continuously map onto I^{ω_1} . So Problem 2 in Shapirovskii [Sh93] has a negative answer. Referring to the last comment in this paper, it seems that the “last word” in a large part of the theory of compact spaces has not yet been spoken.

3. The Hausdorff gap space

Our example will use a (κ, κ) Hausdorff gap where $\omega_1 < \kappa = cf(\kappa) \leq \mathfrak{c}$. Let $(A_\alpha, B_\alpha)_{\alpha < \kappa}$ be such that

Q1: $A_0 = \emptyset = B_0$ and $A_\alpha \cup B_\alpha \subset \omega$

Q2: $\alpha < \beta \Rightarrow (A_\alpha \subset^* A_\beta \text{ and } B_\alpha \subset^* B_\beta)$ (strict almost inclusion)

Q3: $A_\alpha \cap B_\alpha = \emptyset$

Q4: $\nexists A \subset \omega$ such that $\forall \alpha < \kappa (A_\alpha \subset^* A \text{ and } B_\alpha \subset^* \omega \setminus A)$.

Put $Q = \{p \in P : \exists \alpha < \kappa \text{ with } \text{dom}(p) =^* A_\alpha \cup B_\alpha \text{ and } p^{-1}(1) =^* A_\alpha\}$ and let $X = X(Q)$. For each $q \in Q$ define $\delta(q) =$ the unique $\alpha < \kappa$ with $\text{dom}(q) =^* A_\alpha \cup B_\alpha$ and extend δ so that $\delta : X \rightarrow \kappa$ by $\delta(I) = \sup\{\delta(q) : q \in I\}$. This definition of δ is well-defined because if $I \in X$, then by Q4, $\exists \alpha < \kappa$ such that either $A_\alpha \not\subset^* \lambda(I)^{-1}(1)$ or $B_\alpha \not\subset^* \lambda(I)^{-1}(0)$, hence if $\delta(q) > \alpha$, then $q \notin I$.

Lemma 3.1. *X can be partitioned into \mathfrak{c} many closed G_δ subspaces each of which is homeomorphic to an ordinal space $[0, \alpha]$ where $|\alpha| < \kappa$. Thus, X is G_δ -scattered (i.e., scattered in the G_δ topology) and so cannot map continuously onto I^{ω_1} .*

PROOF: Q1–Q4 allows us to easily identify, for each $f \in 2^\omega$, the closed G_δ subspace $\lambda^{-1}(f)$. If $\delta(M_f)$ is an isolated ordinal or if $\delta(M_f)$ is a limit ordinal which is not attained (i.e., $\delta(M_f) \notin M_f$), then $\lambda^{-1}(f) \approx$ the ordinal space $[0, \delta(M_f)]$. If $\delta(M_f)$ is a limit ordinal which is attained, then $\lambda^{-1}(f) \approx [0, \delta(M_f) + 1]$. Thus, X is partitioned into \mathfrak{c} many closed G_δ ordinal subspaces and so every non-empty subspace of X contains a relative G_δ -point. By a result of Shapirovskii [Sh80], X cannot map continuously onto I^{ω_1} . \square

We now partition Q into horizontal sections. For each $\alpha < \kappa$ put $Q^\alpha = \{q \in Q : \delta(q) = \alpha\}$ and put $\mathcal{Q}^\alpha = \{q^+ : q \in Q^\alpha\}$.

Lemma 3.2. *For each $\alpha < \kappa$, \mathcal{Q}^α is a π -base for $\{q^+ \cap B^- : \delta(q) \leq \alpha\}$, i.e., for every q and finite B with $q^+ \cap B^- \neq \emptyset$ and $\delta(q) \leq \alpha$ there exists $r \in \mathcal{Q}^\alpha$ with $r^+ \subset q^+ \cap B^-$.*

PROOF: Assume $q^+ \cap B^- \neq \emptyset$ and let $\delta(q) \leq \alpha$. Choose $I \in q^+ \cap B^-$ and put $f = \lambda(I)$. Put $A = \{p \in B : p \not\leq f\}$ and $C = \{p \in B : p \preceq f\}$. For each $p \in A$ choose $n_p \in \text{dom}(p)$ with $p(n_p) \neq f(n_p)$. Put $R = \text{dom}(q) \cup \{n_p : p \in A\}$

and put $r = f \upharpoonright R$. Since $B \cap I = \emptyset$, for each $p \in C$ and for each finite $H \subset \omega$ we have that $\text{dom}(p) \not\subset R \cup H$. This implies that we can choose, for each $p \in C$, an $m_p \in \text{dom}(p) \setminus R$ such that distinct p 's yield distinct m_p 's. Let s have domain $\{m_p : p \in C\}$ and satisfy that for each $p \in C$, $s(m_p) \neq p(m_p)$. Then, $\delta(r \cup s) = \delta(r) = \delta(q) \leq \alpha$ and $\emptyset \neq (r \cup s)^+ \subset q^+ \cap B^-$. □

Lemma 3.3. *X is ccc.*

PROOF: Assume not and choose an uncountable (meaning of cardinality ω_1 throughout this proof) $R \subset Q$ such that $r \neq s$ in $R \Rightarrow r \perp s$. Since $\delta : Q \rightarrow \kappa$ is $\leq \omega$ -to-1, choose an uncountable $R' \subset R$ such that $\delta \upharpoonright R'$ is 1-1. Since there exist only countably many finite collections of finite subsets of ω , choose an uncountable $R'' \subset R'$ and finite $F, A, G, B, H, C \subset \omega$ such that $p \in R''$ and $\delta(p) = \alpha \Rightarrow \text{dom}(p) = (A_\alpha \setminus F) \cup (B_\alpha \setminus G) \cup H$ and $p^{-1}(1) = (A_\alpha \setminus A) \cup B \cup C$ where F and A are disjoint subsets of A_α , G and B are disjoint subsets of B_α , and H and C are disjoint subsets of $\omega \setminus (A_\alpha \cup B_\alpha)$. Let $E = \{\delta(p) : p \in R''\}$. Since R'' consists of pairwise incompatible elements, we see that for $\alpha \neq \beta$ in E , $(A_\alpha \cup A_\beta) \cap (B_\alpha \cup B_\beta) \neq \emptyset$. Since $\text{cf}(\kappa) > \omega_1$, choose $\gamma < \kappa$ such that $\gamma > \sup(E)$. Choose $n < \omega$ and an uncountable $K \subset E$ such that for each $\alpha \in K$, $A_\alpha \setminus n \subset A_\gamma$ and $B_\alpha \setminus n \subset B_\gamma$. Since $A_\gamma \cap B_\gamma = \emptyset$, for every $\alpha \neq \beta$ in K , $(A_\alpha \cup A_\beta) \cap (B_\alpha \cup B_\beta) \cap n \neq \emptyset$. So we have a finite partition $[K]^2 = \bigcup_{i \in n} \{\{\alpha, \beta\} : i \in (A_\alpha \cup A_\beta) \cap (B_\alpha \cup B_\beta)\}$. By Ramsey's Theorem, get $j < n$ and $\alpha < \beta < \eta$ in K such that $\{\alpha, \beta, \eta\}$ is homogeneous for j . This contradicts that $A_i \cap B_i = \emptyset$ for $i = \alpha, \beta, \eta$. Lemma 3.3 is proved. □

Lemma 3.4. *Q is a point- $< \kappa$ collection, i.e., if $I \in X$, then $\{q^+ : I \in q^+\}$ has cardinality $< \kappa$. Consequently, X does not have Property K_κ (Q is a collection of κ many clopen sets which does not have a linked subcollection of cardinality κ).*

PROOF: If $I \in q^+$ for κ many q 's, then $\lambda(I)^{-1}(1)$ would fill our Hausdorff gap $(A_\alpha, B_\alpha)_{\alpha < \kappa}$ contradicting Q4. □

The above lemma tells us that X is not separable and also that X is not the support of a measure algebra (as these all have Property K_κ for every regular κ).

Lemma 3.5. *X has countable π -character.*

PROOF: Let $I \in X$ and put $\alpha = \delta(I)$. Lemma 3.2 implies that for every neighbourhood $q^+ \cap B^-$ of I , there exists $r^+ \in Q^\alpha$ such that $r^+ \subset q^+ \cap B^-$. Since $|Q^\alpha| = \omega$, we are done. □

So, we have shown

Theorem 3.6. *If there exists a (κ, κ) Hausdorff gap where $\kappa = \text{cf}(\kappa) > \omega_1$, then there exists a compact, ccc, non-separable space X which has countable π -character, character = $\sup\{\lambda : \lambda < \kappa\}$, and which does not continuously map onto I^{ω_1} .*

Corollary. $MA(\omega_1)$ does not imply any of the following:

- (a) Every compact ccc space of countable π -character is separable.
- (b) Every compact ccc space of tightness (or even character) $\leq \omega_1$ is separable.
- (c) Every compact ccc non-separable space continuously maps onto I^{ω_1} .

PROOF: We can apply the theorem because Kunen (cf. Baumgartner [Ba84]) has proved that Martin's Axiom is consistent with $\mathfrak{c} = \omega_2$ + there exists a (ω_2, ω_2) Hausdorff gap. \square

The above (a) answers the question of Juhasz [Ju77] (this question was also repeated on page 209 in Fremlin [Fr84]). The above (b) is a different kind of example showing that the theorem of Shapirovskii [Sh72]:

$MA(\omega_1) \Rightarrow$ Every compact ccc space of tightness $< \omega_1$ is separable cannot be improved in the tightness direction. It is quite different from the first published example (Bell [Be80]); that one was covered by Cantor cubes of uncountable weight. The above (c) is of interest because of the following: Let \mathbf{A} represent the axiom of Todorcevic "Every compact ccc non-separable space maps onto I^{ω_1} ". One use of axiom \mathbf{A} is that it resolves several problems in the literature. S. Todorcevic has shown that $\mathbf{A} \Rightarrow MA(\omega_1)$. What we have shown is that $MA(\omega_1) \not\Rightarrow \mathbf{A}$.

In conclusion, we mention that the question of whether every model of set theory contains an example of a compact ccc non-separable space with countable π -character remains open.

REFERENCES

- [Ba84] Baumgartner J.E., *Applications of the Proper Forcing Axiom*, Handbook of Set-Theoretic Topology, editors K. Kunen and J. Vaughan, North-Holland, 1984, pp. 913–959.
- [Be80] Bell M., *Compact ccc non-separable spaces of small weight*, Topology Proceedings **5** (1980), 11–25.
- [Be88] Bell M., *G_κ subspaces of hyadic spaces*, Proc. Amer. Math. Soc. **104**, No. 2 (1988), 635–640.
- [Be89] Bell M., *Spaces of Ideals of Partial Functions*, Set Theory and its Applications, Lecture Notes in Mathematics **1401**, Springer-Verlag, 1989, pp. 1–4.
- [Fr84] Fremlin D.H., *Consequences of Martin's Axiom*, Cambridge Tracts in Mathematics **84**, Cambridge University Press, 1984.
- [Ju71] Juhasz I., *Cardinal Functions in Topology*, Mathematical Centre Tracts **34**, Mathematisch Centrum, Amsterdam, 1971.
- [Ju77] Juhasz I., *Consistency Results in Topology*, Handbook of Mathematical Logic, editor J. Barwise, North-Holland, 1977, pp. 503–522.
- [Sh72] Shapirovskii B., *On separability and metrizability of spaces with Souslin condition*, Soviet Math. Dokl. **13** (1972), 1633–1637.
- [Sh80] Shapirovskii B., *Maps onto Tikhonov cubes*, Russ. Math. Surv. **35.3** (1980), 145–156.
- [Sh93] Shapirovskii B. (presented by P.Nyikos and J.Vaughan), *The Equivalence of Sequential Compactness and Pseudoradialness in the Class of Compact T_2 Spaces, Assuming CH*,

Papers on General Topology and Applications, *Annals of the New York Academy of Sciences* **704**, 1993, pp. 322–327.

[To89] Todorčević S., *Partition Problems in Topology*, Contemporary Mathematics **84**, American Mathematical Society, Providence, Rhode Island, 1989.

[TF95] Todorčević S., Farah I., *Some Applications of the Method of Forcing*, Yenisey Publ. Co., Moscow, 1995.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MANITOBA, FORT GARRY CAMPUS,
WINNIPEG R3T 2N2, CANADA

E-mail: mbell@cc.umanitoba.ca

(Received January 19, 1996)