

Tomáš Kepka; Petr Němec
Connected transversals -- the Zassenhaus case

Commentationes Mathematicae Universitatis Carolinae, Vol. 41 (2000), No. 2, 299--300

Persistent URL: <http://dml.cz/dmlcz/119165>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2000

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Connected transversals — the Zassenhaus case

TOMÁŠ KEPKA, PETR NĚMEC

Abstract. In this short note, it is shown that if A, B are H -connected transversals for a finite subgroup H of an infinite group G such that the index of H in G is at least 3 and $H \cap H^u \cap H^v = 1$ whenever $u, v \in G \setminus H$ and $uv^{-1} \in G \setminus H$ then $A = B$ is a normal abelian subgroup of G .

Keywords: group, subgroup, connected transversals, core

Classification: 20F12, 20D60

Throughout this extremely short note, let H be a subgroup of a group G such that the index of H in G is at least 3 and $H \cap H^u \cap H^v = 1$ whenever $u, v \in G \setminus H$ and $uv^{-1} \in G \setminus H$. Furthermore, let A, B be subsets of G such that $AH = G = BH$ and $a^{-1}b^{-1}ab \in H$ for all $a \in A$ and $b \in B$. In his remarkable paper [3], A. Drápal showed that then $A = B$ is an abelian subgroup of G , provided that G is finite. The purpose of the present modest note is to check that $A = B$ is a normal abelian subgroup of G , provided that H is finite and G is infinite (notice that if H is infinite then neither A nor B need to be a subgroup of G — see [1] and [2]). The kind reader is fully referred to [1], [2] and [3] as concerns all the necessary background and many further useful details and connections.

In the rest of this note, assume that H is finite and G is infinite. If $G_1 = \langle A, B \rangle$ and $H_1 = H \cap G_1$ then $AH_1 = G_1 = BH_1$ and we show first that the centralizer K of H_1 in G_1 is of finite index in G_1 .

Since $n = |H_1|$ is finite, we have $H_1 \subseteq \langle C \rangle$ for a finite subset $C \subseteq A \cup B$. Now, take $c \in C$ and put $K_c = \{x \in G_1; xc = cx\}$ and $B_u = \{b \in B; c^{-1}b^{-1}cb = u\}$ for every $u \in H_1$ (here we assume $c \in A$, the other case being similar). Then $B = \bigcup B_u$, this union is disjoint and $b_2b_1^{-1} \in K_c$ for all $b_1, b_2 \in B_u$. Further, for every $u \in H_1$ such that $B_u \neq \emptyset$, choose $b_u \in B_u$ and put $D_c = \{b_u; u \in H_1\}$. Then $G_1 = K_c D_c H_1$, $|D_c H_1| \leq n^2$ and it follows easily that the index of K_c in G_1 is at most n^2 . On the other hand, $K = \bigcap K_c$ and consequently the index of K in G_1 is finite, too.

This research has been supported by the Grant Agency of the Czech Republic, grant # GAČR-201/99/0263.

Since H is finite, G_1 is of finite index in G and we conclude that also the index of K in G is finite. Finally, if $L = K_G$ denotes the core of K in G then G/L is a finite group. Since G is infinite, L must be so and the finiteness of H implies that we can find elements $u_1, v_1 \in L \setminus H$ such that $u_1 v_1^{-1} \notin H$. Now, $H_1 = H_1 \cap H_1^{u_1} \cap H_1^{v_1} \subseteq H \cap H^{u_1} \cap H^{v_1} = 1$ and we have proven that $H_1 = 1$. Consequently, $ab = ba$ for all $a \in A$ and $b \in B$.

Next, we show that both A and B are subgroups of G . Indeed, if $H_2 = \langle A \rangle \cap H$ then $H_2^b = H_2 \subseteq H$ for every $b \in B$ and we see that $H_2 \subseteq H_G = 1$, where H_G is the core of H in G . Thus $H_2 = 1$ and it follows easily that $\langle A \rangle = A$. Quite similarly, B is a subgroup of G .

Since the index of $A \cap B$ in G is finite, we can find $e \in A \cap B$, $e \neq 1$. If $a \in A$ and $b \in B$ are such that $a^{-1}b \in H$ then $a^{-1}b \in H \cap H^u \cap H^v$, where $u = a^{-1}$ and $v = (a^{-1}ea)^{-1}$. Assume, for a moment, that $a \neq b$. Then $u, v \in G \setminus H$, $uv^{-1} \in G \setminus H$, and therefore $H \cap H^u \cap H^v = 1$ and $a = b$, a contradiction. Thus $a = b$ and we have shown that $A = B$. Clearly, this subgroup is abelian.

It remains to show that A is normal in G . Since A is of finite index in G , the core A_G is infinite. If E denotes the centralizer of A_G , then $A \subseteq E$, E is normal in G and we have $E = AH_3$ for some $H_3 \subseteq H$. Finally, $H_3 \subseteq H^w$ for every $w \in A_G$, and therefore $H_3 = 1$ and $A = E$.

REFERENCES

- [1] Drápal A., *Multiplication groups of free loops I*, Czech. Math. J. **46** (121) (1996), 121–131.
- [2] Drápal A., *Multiplication groups of free loops II*, Czech. Math. J. **46** (121) (1996), 201–220.
- [3] Drápal A., *Multiplication groups of finite loops that fix at most two points*, submitted.

DEPARTMENT OF ALGEBRA, FACULTY OF MATHEMATICS AND PHYSICS, CHARLES UNIVERSITY,
SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail: keпка@karlin.mff.cuni.cz

DEPARTMENT OF MATHEMATICS, ČZU, KAMÝČKÁ 129, 165 21 PRAHA 6 – SUCHDOL, CZECH
REPUBLIC

E-mail: nemeц@tf.czu.cz

(Received October 4, 1999)