

Jakub Černý; Jan Kára; Daniel Král; Pavel Podbrdský; Miroslava Sotáková;  
Robert Šámal

On the number of intersections of two polygons

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 44 (2003), No. 2, 217–228

Persistent URL: <http://dml.cz/dmlcz/119381>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## On the number of intersections of two polygons

JAKUB ČERNÝ, JAN KÁRA, DANIEL KRÁL',  
PAVEL PODBRDSKÝ, MIROSLAVA SOTÁKOVÁ, ROBERT ŠÁMAL

*Abstract.* We study the maximum possible number  $f(k, l)$  of intersections of the boundaries of a simple  $k$ -gon with a simple  $l$ -gon in the plane for  $k, l \geq 3$ . To determine the number  $f(k, l)$  is quite easy and known when  $k$  or  $l$  is even but still remains open for  $k$  and  $l$  both odd. We improve (for  $k \leq l$ ) the easy upper bound  $kl - l$  to  $kl - \lceil k/6 \rceil - l$  and obtain exact bounds for  $k = 5$  ( $f(5, l) = 4l - 2$ ) in this case.

*Keywords:* geometry, polygon, intersection, combinatorial complexity

*Classification:* 52C45, 52C10

### 1. Introduction

To determine the maximum complexity of union of two or more geometric objects in the plane is among basic extremal geometric problems, see e.g. [1], [2], [3], [5], [6] and [7]. Let  $k, l \geq 3$  be given integer numbers. We are interested in the problem of determining the maximum possible number  $f(k, l)$  of intersections of a simple  $k$ -gon and a simple  $l$ -gon. This problem was studied in [2] — the cases when  $k$  or  $l$  is even are solved there, but an unrecoverable error appears in the case of  $k$  and  $l$  being both odd. Similar problem was also studied in [4].

We introduce basic definitions and notation in Section 2. The difficulty of determining the number  $f(k, l)$  depends on the parity of  $k$  and  $l$ . If one of these numbers is even, the problem is quite easy and solved; we survey the previous results in Section 3.

We deal with the case of  $k$  and  $l$  being both odd in Section 4. The bounds  $kl - k - l + 3 \leq f(k, l) \leq kl - l$  (for  $k \leq l$ ) are proved in [2]. If  $k, l \geq 7$  we improve the upper bound and prove  $f(k, l) \leq kl - \lceil \frac{k}{6} \rceil - l$  for  $k \leq l$  (Theorem 4). The conjecture is that the lower bound  $kl - k - l + 3$  is tight.

We focus on the number of intersections of an  $l$ -gon and a pentagon for  $l$  odd in Section 5. We prove that  $f(5, l) = 4l - 2$  (Theorem 5).

The general problem of determining  $f(k, l)$  for  $k$  and  $l$  both odd,  $k, l \geq 7$  remains open.

---

This research was supported by GAUK 158/99 and GAČR 201/99/0242. Institute for Theoretical Computer Science (ITI) is supported as project LN00A056 by the Ministry of Education of Czech Republic.

## 2. Definitions and notation

We mean by a *segment* a closed line segment. If  $A$  and  $B$  are points of the plane, then we write  $AB$  for the segment connecting them. We assume throughout the whole paper that all the end-points of all the segments are in general position. Two segments *intersect* if they share an interior point and we call that point their *intersection*. The set of segments is *intersection-free* if no two of segments of the set intersect. Two segments *miss* each other if they do not intersect.

The sequence of non-intersecting segments  $A_1A_2, A_2A_3, \dots, A_kA_{k+1}$  where  $A_i \neq A_j$  for  $i \neq j$  is called a *path*; we write shortly  $A_1A_2 \dots A_kA_{k+1}$  instead of  $A_1A_2, A_2A_3, \dots, A_kA_{k+1}$  in the paper. The *length* of the path is the number of segments which it contains. If  $A_{k+1} = A_1$ , then the sequence forms a *k-gon*; if we do not want to emphasize the length of the sequence, we say a *polygon* instead of a *k-gon*. *Intersections of two polygons* are the intersections of segments which form their boundaries. For given  $k, l \geq 3$  we denote  $f(k, l)$  the maximum number of intersections of some *k-gon* and *l-gon*.

Let  $\mathcal{A}$  and  $\mathcal{B}$  be two intersection-free sets of segments. Then their *non-intersection graph* is a bipartite graph with the vertex set  $\mathcal{A} \cup \mathcal{B}$  such that a segment of  $\mathcal{A}$  is joined by an edge to a segment of  $\mathcal{B}$  iff these two segments do not intersect; the segments of  $\mathcal{A}$  ( $\mathcal{B}$ ) are not mutually joined by edges. We state two easy observations to get familiar with these definitions:

**Observation 1.** *Let  $\mathcal{A}$  and  $\mathcal{B}$  be two intersection-free sets of segments; let  $\mathcal{A}' \subseteq \mathcal{A}$  and  $\mathcal{B}' \subseteq \mathcal{B}$ . Then the non-intersection graph of  $\mathcal{A}'$  and  $\mathcal{B}'$  is the subgraph of the non-intersection graph of  $\mathcal{A}$  and  $\mathcal{B}$  induced by the vertex set  $\mathcal{A}' \cup \mathcal{B}'$ .*

**Observation 2.** *Let  $ABCD$  be a path which forms together with the segment  $DA$  a convex quadrilateral and let  $p$  be any segment in the plane. Then the vertex  $p$  has degree at least 1 in the non-intersection graph of  $\mathcal{A} = \{AB, BC, CD\}$  and  $\mathcal{B} = \{p\}$ .*

PROOF: If the degree of  $p$  is zero, then  $p$  intersects all the three segments  $AB$ ,  $BC$  and  $CD$  which is impossible since  $A$ ,  $B$ ,  $C$  and  $D$  form a convex quadrilateral. □

## 3. Intersections with $N$ -gons for even $N$ 's

The following theorems and lemmas are proved (using a different notation) in [2]. Note that the upper bounds in Theorem 1 and Theorem 2 easily follow from Lemma 1.

**Lemma 1.** *Let  $K$  be a  $k$ -gon and let  $p$  be a line. The number of intersections of  $p$  with the segments of  $K$  is even and at most  $k$ . Thus no line intersects all the segments of a polygon with odd number of vertices.*

**Theorem 1.** *Let  $k, l \geq 3$  be two even integers. Then  $f(k, l) = kl$  (the lower bound is shown by Figure 1).*




Figure 1: A  $k$ -gon  $K$  and an  $l$ -gon  $L$  with  $kl$  intersections for  $k$  and  $l$  both even.

**Theorem 2.** *Let  $k \geq 3$  be even and let  $l \geq 3$  be odd. Then  $f(k, l) = k(l - 1)$  (the lower bound is shown by Figure 2).*




Figure 2: A  $k$ -gon  $K$  and an  $l$ -gon  $L$  with  $k(l - 1)$  intersections for even  $k$  and odd  $l$ .

*Note:* Figure 1 and Figure 2 can be generalized for arbitrary  $k, l$  by an easy trick. We can substitute one segment of the polygon  $K$  (or  $L$ ) which looks like I by three segments which look like a narrow N. This can obviously be done in such a way that we obtain sufficiently many new intersections.

#### 4. Intersections with $N$ -gons for odd $N$ 's

We assume  $k \leq l$  in the whole section.

The following lower bound on  $f(k, l)$  is also shown in [2].

**Theorem 3.** *Let  $k, l \geq 3$  be two odd integers. Then  $f(k, l) \geq (k-1)(l-1) + 2 = kl - k - l + 3$  (see Figure 3).*




Figure 3: A  $k$ -gon  $K$  and an  $l$ -gon  $L$  with  $kl - k - l + 3$  intersections for both  $k$  and  $l$  odd.

We focus our attention on proving an upper bound on the number of intersections of two polygons with odd number of vertices in the rest of this section. There is an easy upper bound  $kl - l$  (Theorem 2) which gives the exact value for  $k = 3$  (in this case, it is equal to the lower bound). For  $k \geq 7$  we improve this bound to  $kl - \lceil k/6 \rceil - l$  in Theorem 4. We first prove several lemmas on the number of intersections of a path and a non-intersecting set of segments:

**Lemma 2.** *Let  $ABCDE$  be a path such that the point  $D$  is inside the triangle  $ABC$ . Then there is no intersection-free set  $\mathcal{B}$  of four segments such that the non-crossing graph of  $\mathcal{A} = \{AB, BC, CD, DE\}$  and  $\mathcal{B}$  forms a perfect matching.*

PROOF: Let us suppose there exists such a  $\mathcal{B}$ . Let  $p$  be the segment which misses only  $CD$  and let  $q$  be the segment which misses only  $DE$ . We can suppose that  $p$  and  $q$  do not share an end-point, since we could shorten them otherwise. Note that  $p$  and  $q$  do not intersect each other ( $\mathcal{B}$  is intersection-free) and they both intersect  $AB$  and  $BC$ . Hence the segments  $p, q, AB$  and  $BC$  split the plane into three regions (see Figure 4); let  $T$  be the triangle and  $Q$  be the quadrilateral. The point  $E$  has to lie inside the triangle  $T$ , since  $DE$  intersects  $p$  but not  $q$ . Thus the segments  $CD$  and  $DE$  split the quadrilateral  $Q$  into two pentagons; let  $P$  be the convex one of them. It is either  $P \cap AB = \emptyset$  or  $P \cap BC = \emptyset$ .

Let us suppose first that  $P \cap AB = \emptyset$ . Let the vertices of  $P$  be  $B', C', D', D$  and  $E'$  where  $B', C' \in BC, D' \in CD$  and  $E' \in DE$ . Let  $r$  be the segment of  $\mathcal{B}$  which misses only  $AB$ ;  $r$  does not intersect  $p$  or  $q$ , but it intersects  $BC, CD$  and  $DE$ ; thus it has to intersect  $CD$  between  $D'$  and  $D$  and  $DE$  between  $D$  and  $E'$ . Then  $r$  intersects the segments of the convex pentagon  $P$  more than twice, but this is impossible. The case that  $P \cap BC = \emptyset$  is symmetric. We have just proved that no such set  $\mathcal{B}$  can exist.  $\square$




Figure 4: The triangle  $ABC$ , the point  $D$  inside it, the triangle  $T$  and quadrilateral  $Q$ , and the forced position of the point  $E$ .

**Lemma 3.** *Let  $ABCDEFG$  be a path. Then there is no intersection-free set  $\mathcal{B}$  of six segments such that the non-crossing graph of  $\mathcal{A} = \{AB, BC, CD, DE, EF, FG\}$  and  $\mathcal{B}$  forms a perfect matching.*

PROOF: Let us suppose the existence of such a  $\mathcal{B}$ . The lines  $BC, CD$  and  $BD$  split the plane into seven regions (see Figure 5):  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  and  $\eta$ .




Figure 5: The seven regions  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  and  $\eta$  in which the lines  $BC, CD$  and  $BD$  split the plane.

The point  $E$  cannot lie in  $\alpha$ , since  $BC$  and  $DE$  do not intersect, it cannot lie in  $\beta$  due to Observation 1 and Lemma 2 used for the path  $EDCBA$ , it cannot lie in  $\gamma$  due to Observation 2 used for the path  $BCDE$ , it cannot lie in  $\epsilon$ , since any

segment intersecting both  $BC$  and  $DE$  would not miss  $CD$ , and it cannot lie in  $\eta$  due to Observation 1 and Lemma 2 used for the path  $BCDEF$ . Thus the point  $E$  is either in  $\delta$  or in  $\zeta$ .




Figure 6: The regions  $\xi$ ,  $\psi$  and  $\omega$ ;  $\xi$  and  $\psi$  are marked by gray.

We first deal with the case that the point  $E$  lies in  $\zeta$ . Consider the regions  $\xi$ ,  $\psi$  and  $\omega$  as shown in Figure 6. The point  $F$  cannot lie in  $\xi$  due to Observation 1 and Lemma 2 used for the path  $CDEFG$ , it cannot lie in  $\psi$  due to Observation 1 and Lemma 2 used for the path  $FEDCB$  and it cannot lie in  $\omega$ , since  $EF$  crosses neither  $BC$  nor  $CD$ . If  $F$  were outside  $\xi$ ,  $\psi$  and  $\omega$ , it would be impossible to intersect all the segment except for  $DE$  by a segment. We conclude that  $E$  cannot lie in  $\zeta$ .




Figure 7: The regions  $\sigma$ ,  $\tau$ ,  $v$ ,  $\phi$ ,  $\chi$ ,  $\psi$ ,  $\rho$  and  $\omega$ ; the regions  $\sigma$ ,  $v$ ,  $\phi$  and  $\chi$  are marked by gray.

The point  $E$  has to lie in  $\delta$  according to the discussion above. Consider the regions  $\sigma$ ,  $\tau$ ,  $v$ ,  $\phi$ ,  $\chi$ ,  $\psi$ ,  $\rho$  and  $\omega$  as shown in Figure 7. We discuss the position of  $F$ .

The point  $F$  cannot lie in  $\sigma$  due to Observation 1 and Lemma 2 used for the path  $CDEFG$ , it cannot lie in  $\tau$  due to Observation 2 used for the path  $CDEF$  and it cannot lie in  $\nu$  due to Observation 1 and Lemma 2 used for the path  $FEDCB$ . The point  $F$  cannot lie in  $\phi$  or  $\omega$ , since there would be no segment intersecting  $BC$ ,  $DE$  and  $EF$  which misses  $CD$  and the point  $F$  cannot lie in  $\rho$ , since there would be no segment intersecting  $BC$ ,  $CD$  and  $EF$  which misses  $DE$ . Thus the point  $F$  has to lie either in  $\psi$  or  $\chi$ ; these two cases are “symmetric” with respect to changing the orientation of the given path (we take the path  $GFEDCBA$  instead of  $ABCDEFG$ ) — see Figure 8. We may thus assume w.l.o.g. that  $F$  is in  $\psi$ .




Figure 8: The cases that the point  $F$  is in  $\psi$  or  $\chi$  are symmetric.




Figure 9: The segments  $p$  and  $q$  and the triangle  $T'$ .

The situation is shown in Figure 9. Let  $B'$  be the intersection of the lines  $BC$  and  $EF$  and let  $T$  be the triangle  $B'CE$ . Every line intersecting both  $CD$  and  $DE$  has to intersect also  $B'C$  and  $EF$ . If  $A$  were outside  $T$ , then there could not be a segment which misses only  $BC$  and a segment which misses only  $EF$



at once. A similar argument holds also for the point  $G$ . Thus both the points  $A$  and  $G$  are inside  $T$ . Let  $p$  be the segment of  $\mathcal{B}$  which misses only  $AB$  and let  $q$  be the segment which misses only  $FG$ . Since  $p$  ( $q$ ) intersects  $CD$  and  $DE$ , the intersection with  $BC$  is between  $B'$  and  $C$ . The triangle  $T$  is splitted into three regions by  $p$  and  $q$ , see Figure 9; let  $T'$  be the triangular one. We first deal with the case that one side of  $T'$  is formed by  $q$ . Since  $FG$  misses  $q$ , the point  $G$  has to lie inside  $T'$ . But then, the segments  $FG$  and  $p$  do not intersect. If one side of  $T'$  is formed by  $p$ , then  $A$  has to lie inside  $T'$  and  $AB$  and  $q$  do not intersect. Both cases are impossible. But this was the very last possible configuration and thus the set  $\mathcal{B}$  does not exist.  $\square$

Lemma 3 is the best possible in the sense that there is a path of length five and a set of five non-intersecting segments such that their non-intersection graph is a perfect matching (see Figure 10).




Figure 10: A path of length five and a set of five non-intersecting segments such that their non-intersection graph is a perfect matching.

**Lemma 4.** *Let  $ABCDEFGF$  be a path and let  $\mathcal{B}$  be a set of non-intersecting segments. The following holds for at least one segment  $r$  among  $AB$ ,  $BC$ ,  $CD$ ,  $DE$ ,  $EF$  and  $FG$ : If a segment  $q$  of  $\mathcal{B}$  intersects at least five segments of the path, then it intersects  $r$ .*

PROOF: If the lemma does not hold, then for each segment  $p$  of  $AB$ ,  $BC$ ,  $CD$ ,  $DE$ ,  $EF$  and  $FG$ , there exists a segment of  $\mathcal{B}$  which misses only  $p$ . But then these six segments of  $\mathcal{B}$  and the path  $ABCDEFGF$  contradict Lemma 3.  $\square$

**Theorem 4.** *Let  $k, l \geq 7$  be odd. Then  $f(k, l) \leq kl - \lceil k/6 \rceil - l$ .*

PROOF: Let  $K$  be any  $k$ -gon and let  $L$  be any  $l$ -gon. Let  $\mathcal{A}$  be the set of segments forming the polygon  $K$  and let  $\mathcal{B}$  be the set of segments forming the polygon  $L$ ; let  $G$  be a non-intersection graph of  $\mathcal{A}$  and  $\mathcal{B}$ . Each vertex of  $G$  has degree at least one, since no segment can intersect all the segments of a polygon with odd number of vertices due to Lemma 1.

Let  $p_1, \dots, p_k$  be the segments of  $K$ . Let  $r_1$  be the segment for the path  $p_1p_2p_3p_4p_5p_6$  with the properties described in Lemma 4,  $r_1 = p_{i_1}$  for some  $1 \leq i_1 \leq 6$ . Let  $r_2$  be the segment for the path  $p_{i_1+1}p_{i_1+2}p_{i_1+3}p_{i_1+4}p_{i_1+5}p_{i_1+6}$  with the properties described in Lemma 4,  $r_2 = p_{i_2}$  for some  $i_1 + 1 \leq i_2 \leq i_1 + 6$ . By repeating this procedure we can find segments  $r_j, j = 1, 2, \dots, \lceil k/6 \rceil$ , it is easy to see that the path for  $r_{\lceil k/6 \rceil}$  contains none of the segments  $r_1, r_2, \dots, r_{\lceil k/6 \rceil - 1}$ .

Each segment of  $\mathcal{B}$  misses at least one segment of  $\mathcal{A}$  different from the segments  $r_1, r_2, \dots, r_{\lceil k/6 \rceil}$ ; otherwise due to Lemma 4 it would have to intersect also  $r_{\lceil k/6 \rceil}, r_{\lceil k/6 \rceil - 1}, \dots, r_2$  and  $r_1$  hence it would have to intersect all the segments of  $\mathcal{A}$  which is impossible due to Lemma 1 since the segments of  $\mathcal{A}$  form a polygon with odd number of vertices.

Let  $\mathcal{A}' \subseteq \mathcal{A}$  be the set  $\{r_1, \dots, r_{\lceil k/6 \rceil}\}$ . The degree of each vertex of  $\mathcal{B}$  in the subgraph of  $G$  induced by  $(\mathcal{A} \setminus \mathcal{A}') \cup \mathcal{B}$  is at least one, since each segment of  $\mathcal{B}$  misses at least one segment of  $\mathcal{A} \setminus \mathcal{A}'$ ; thus this subgraph contains at least  $l$  edges. The degree of each vertex of  $\mathcal{A}'$  in the subgraph of  $G$  induced by  $\mathcal{A}' \cup \mathcal{B}$  is at least one, since no segment can intersect all the segments of  $\mathcal{B}$ ; this subgraph contains at least  $\lceil k/6 \rceil$  edges. The whole graph  $G$  thus contains at least  $\lceil k/6 \rceil + l$  edges and thus the number of intersections of  $K$  and  $L$  is at most  $kl - \lceil k/6 \rceil - l$ .  $\square$

### 5. Intersections with pentagons

**Observation 3.** *Let  $K$  be a polygon. Then there exist three consecutive vertices  $A, B, C$  of  $K$  such that the interior angle  $\angle ABC$  is convex and no vertex of  $K$  lies in the interior of the triangle  $\triangle ABC$ .*

PROOF: As a well known fact every (not necessarily convex) polygon has a triangulation. Let us consider any triangulation  $T$  of  $K$ . Let  $G$  be a graph with the vertex set consisting of all triangles of  $T$ ; two distinct triangles are connected by an edge iff they are sharing a common segment. Then  $G$  is obviously a tree and we can take  $A, B, C$  to be vertices of any leaf of  $G$ .  $\square$

**Lemma 5.** *Let  $K$  be a  $k$ -gon for  $k \geq 5$  odd. Then there are four consecutive vertices  $A, B, C, D$  of  $K$  such that one of the four following conditions holds:*

- (1) *No line  $p$  intersects all the segments  $AB, BC$  and  $CD$ .*
- (2) *Every line  $p$  which intersects the segments  $AB$  and  $BC$  intersects also the segment  $CD$ .*
- (3) *Every line  $p$  which intersects the segments  $AB$  and  $CD$  intersects also the segment  $BC$ .*
- (4) *Every line  $p$  which intersects the segments  $BC$  and  $CD$  intersects also the segment  $AB$ .*

PROOF: Take the points  $A, B, C$  constructed in the Observation 3 and let  $D$  be the vertex neighboring with  $C$  (different from  $B$ ). The lines  $AB, BC$  and  $CA$  split the plane into the six regions  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  where the vertex  $D$  can lie (due to the choice of the points  $A, B, C$  no vertex of  $K$  lies in the interior of the triangle  $\triangle ABC$ ) (see Figure 11). We can w.l.o.g. suppose that  $D$  does not lie in the region  $\zeta$  (if it does we can take  $A' = C, B' = B, C' = A$  and  $D'$  the vertex neighboring with  $C'$  different from  $B'$ , then  $D'$  cannot lie in  $\zeta' = \delta$  unless the segments  $CD$  and  $C'D'$  intersect which is impossible). The vertex  $D$  cannot lie in the region  $\alpha$  because  $AB$  and  $CD$  cannot intersect. If  $D$  lies in  $\beta$  then condition (2) holds. If  $D$  lies in  $\gamma$  then condition (3) holds. If  $D$  lies in  $\delta$  then condition (4) holds. Finally if  $D$  lies in  $\epsilon$  then condition (1) holds.  $\square$




Figure 11: The six regions  $\alpha, \beta, \gamma, \delta, \epsilon, \zeta$  where the vertex  $D$  can lie.

**Corollary 1.** *Let  $K = ABCDE$  be a pentagon. There are four segments among  $AB, BC, CD, DE$  and  $EA$  such that no line intersects all of these four segments.*

PROOF: Suppose that  $A, B, C, D$  are the vertices constructed in Lemma 5. If condition (1) of Lemma 5 holds then no line intersects all the segments  $AB, BC, CD$  hence we can take e.g. the segments  $AB, BC, CD$  and  $DE$ . If condition (2) holds then no line intersects all the segments  $AB, BC, DE, EA$  (if such line exists then it also intersects the segment  $CD$  by condition (2) which is impossible due to Lemma 1). The cases when conditions (3) or (4) hold are similar.  $\square$

**Theorem 5.** *Let  $l \geq 5$  be odd. Then  $f(5, l) = 4l - 2$ .*

PROOF: The inequality  $f(5, l) \geq 4l - 2$  is assured by Theorem 3. It remains to prove the upper bound for the number of intersections. Let  $L$  be any  $l$ -gon and  $K$

be any pentagon. Each segment of  $L$  has at most 4 intersections with segments of  $K$  due to Lemma 1. Thus  $K$  and  $L$  cannot have more than  $4l$  intersections. The number of intersections of  $K$  and  $L$  is even: Imagine that we pass along the boundary of  $K$ ; each time we intersect a segment of  $L$ , we either enter or leave the interior region of  $L$  and we cannot enter or leave it in any other way. Thus if the number of intersections is more than  $4l - 2$ , it has to be  $4l$  and each segment of  $L$  intersects exactly four segments of  $K$ . But there are four segments of  $K$  which cannot be intersected by a line (more likely by a segment) due to Corollary 1; let  $p$  be the remaining segment of  $K$ . Each of the segments of  $L$  has to intersect  $p$  (otherwise it would intersect the remaining four segments of  $K$  which is impossible). But  $p$  cannot intersect all the segments of  $L$  due to Lemma 1 — thus  $K$  and  $L$  have at most  $4l - 2$  intersections.  $\square$

*Note:* It is quite obvious that Corollary 1 and thus the upper bound in Theorem 5 can be easily generalized to all  $k$ -gons for  $k \geq 5$  odd. We obtain  $f(k, l) \leq kl - l - 2$  for  $5 \leq k \leq l$  and  $k, l$  odd. However for  $k > 5$  Theorem 4 gives us the same or even better bound.

## 6. Conclusion

We have focused on the case of  $k$  and  $l$  being both odd. We proved an exact bounds on  $f(k, l)$  when  $k$  is 3 or 5 and  $l$  is arbitrary odd. The case that  $k$  and  $l$  are both odd and at least seven remains open also for  $k = l = 7$ . We improved the simple upper bound of  $kl - l$  to  $kl - \lceil k/6 \rceil - l$ . The original conjecture is that there are no  $k$ -gon and  $l$ -gon with more than  $kl - k - l + 3$  intersections; the construction of a  $k$ -gon and an  $l$ -gon with  $kl - k - l + 3$  intersections is described in Theorem 3. The gap between the lower and the upper bound is still linear in the number of vertices thus the natural task might be: Improve the upper (lower) bound on the maximum possible number of intersections of two polygons.

**Acknowledgment.** We would like to thank Peter Brass, Jan Kratochvíl and Pavel Valtr for bringing this problem to our attention and for many helpful discussions.

## REFERENCES

- [1] Aronov B., Sharir M., *The common exterior of convex polygons in the plane*, Comput. Geom. **8** (3) (1997), 139–149.
- [2] Dillencourt M.B., Mount D.M., Saalfeld A.J., *On The Maximum Number of Intersections of Two Polyhedra in 2 and 3 Dimensions*, Proceedings of the Fifth Canadian Conference on Computational Geometry, Waterloo, Ontario, August, 1993, pp. 49–54.
- [3] Efrat A., Sharir M., *On the complexity of the union of fat convex objects in the plane*, Discrete Comput. Geom. **23** (2000), 171–189.
- [4] Grünbaum B., *Selfintersections of polygons*, Geombinatorics **8** (1998), no. 2, 37–45.
- [5] van Kreveld M., *On fat partitioning, fat covering and the union size of polygons*, Comput. Geom. **9** (1994), no. 4, 197–210.

- [6] Matoušek J., Pach J., Sharir M., Sifrony S., Welzl E., *Fat triangles determine linearly many holes*, SIAM J. Comput. **23** (1994), no. 1, 154–169.
- [7] Pach J., Sharir M., *On the boundary of the union of planar convex sets*, Discrete Comput. Geom. **3** (1999), 321–328.

DEPARTMENT OF APPLIED MATHEMATICS AND INSTITUTE FOR THEORETICAL COMPUTER SCIENCE (ITI), CHARLES UNIVERSITY, MALOSTRANSKÉ NÁM. 25, 118 00 PRAGUE 1, CZECH REPUBLIC

*E-mail:* kuba@atrey.karlin.mff.cuni.cz  
jack@atrey.karlin.mff.cuni.cz  
kral@atrey.karlin.mff.cuni.cz  
podbrdsk@karlin.mff.cuni.cz  
sotam0am@artax.karlin.mff.cuni.cz  
samal@atrey.karlin.mff.cuni.cz

(Received May 24, 2002, revised February 14, 2003)