

Josef Král; Beloslav Riečan

Note on the Stokes Formula for 2-Dimensional Integrals in n -Space

Matematicko-fyzikálny časopis, Vol. 12 (1962), No. 4, 280--292

Persistent URL: <http://dml.cz/dmlcz/126330>

Terms of use:

© Mathematical Institute of the Slovak Academy of Sciences, 1962

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

NOTE ON THE STOKES FORMULA FOR 2-DIMENSIONAL INTEGRALS IN n -SPACE

JOSEF KRÁL, Praha, BELOSLAV RIEČAN, Bratislava

In present note some theorems of the Stokes type concerning curvilinear and 2-dimensional integrals in n -space are established.

1. Introduction. The term path (on $\langle a, b \rangle$) is taken to mean a continuous mapping f of $\langle a, b \rangle = \{t; t \in E_1, a \leq t \leq b\}$ into E_n , the Euclidean n -space; f will be termed closed provided $f(a) = f(b)$. (For $n = 2$ we shall speak of a plane path.) The length of f on $\langle a, b \rangle$ is defined as usual; we say that f is rectifiable if its length is finite. Let f be a plane path on $\langle a, b \rangle$ and let $\Phi = [\Phi_1, \dots, \Phi_n]$, $\Psi = [\Psi_1, \dots, \Psi_n]$ be continuous mappings of $[f] = f(\langle a, b \rangle)$ into E_n . We put

$$\int_f \Phi d\Psi = \sum_{i=1}^n \int_a^b \Phi_i(f(t)) d\Psi_i(f(t))$$

provided the Stieltjes integrals on the right-hand side exist. If $K = \langle \alpha, \beta \rangle \times \langle \gamma, \delta \rangle$ is a rectangle then f_K will stand for the closed plane path describing simply the boundary of K in positive sense.

Let now Φ, Ψ be continuous mappings of K into E_n . General conditions are known which, imposed on Φ and Ψ , secure the existence of an integrable function γ on K with

$$\int_{f_K} \Phi d\Psi = \iint_K \gamma \tag{1}$$

(the integral on the right-hand side is taken in the sense of Lebesgue). The aim of the present paper is, roughly speaking, to extend the validity of (1) to the case where f_K is replaced by a finite number of rectifiable closed plane paths with any number of self-intersections.

If $G \subset E_2$ is an open set and Φ, Ψ are mappings of G into E_n , then

$$\gamma = \text{rot}(\Phi, \Psi) \text{ in } G$$

means that (1) holds for every rectangle $K \subset G$.

Given a closed plane path f and a point $z \in E_2 - [f]$ we shall denote by $\text{ind}(z, f)$ the index of z with respect to f . (The reader may consult T. Radó's monograph [1], II. 4. 34 and IV. 1. 24 for a precise definition.) Our main objective is to prove the following theorem.

1,1. Theorem. Let f^1, \dots, f^m be rectifiable closed plane paths and put $C = \bigcup_{k=1}^m [f^k]$,

$\omega(z) = \sum_{k=1}^m \text{ind}(z; f^k) \quad (z \in E_2 - C), \quad G = \{z; z \in E_2 - C, \omega(z) \neq 0\}, \quad G_p =$
 $= \{z; z \in E_2 - C, \omega(z) = p\}$. Let Φ, Ψ be continuous mappings of $C \cup G$ into E_n
and suppose that Ψ is Lipschitzian on $C \cup G$ and $\gamma = \text{rot}(\Phi, \Psi)$ in G .

Then

$$\sum_{k=1}^m \int_{f^k} \Phi \, d\Psi = \sum_{l=1}^{\infty} \left[\sum_{p \geq l} \left(\iint_{G_p} \gamma - \iint_{G_{-p}} \gamma \right) \right] \quad (2)$$

provided the Lebesgue integrals $\iint_{G_p} \gamma$ ($p \neq 0$) exist.

1,2. Remark. A sort of formula (2) still holds even if the Lebesgue integrals $\iint_{G_p} \gamma$ ($p \neq 0$) fail to exist (cf. theorem 11,1 below).

The right-hand side in (2) may be replaced by the series

$$\sum_{p=1}^{\infty} p \left(\iint_{G_p} \gamma - \iint_{G_{-p}} \gamma \right) \quad (3)$$

provided (3) is convergent (possibly, non-absolutely). In [2], p. 595, an example is given showing that (3) may actually diverge even in the relatively simple case where $n = 2$ and Ψ is the identity map. If the integral

$$\iint_G \omega \gamma \quad (4)$$

happens to exist, then, in (2), we may write simply (4) instead of $\sum_{l=1}^{\infty} \left[\sum_{p \geq l} (\dots) \right]$.

1,3. Remark. From 1,1 we obtain as a corollary a theorem of the Stokes type for 2-dimensional Lipschitzian surfaces in E_n bounded by a finite number of rectifiable curves (cf. remark 11,4 below). The reader may consult H. Whitney's monograph [3] for the rôle of analogous theorems dealing with k -dimensional integrals. An extensive bibliography concerning the Stokes formula together with corresponding comments on the subject is given in K. Krickeberg's article [4].

2. Before going into the proof of our main theorem we shall establish several auxiliary results. Let us agree to accept the following notation. H_r will stand for the r -dimensional Hausdorff measure. Given $z = [z_1, \dots, z_{r+1}] \in E_{r+1}$ and a positive integer $i \in \langle 1, r+1 \rangle$ we put $\hat{z}^i = [z_1, \dots, z_{i-1}, z_{i+1}, \dots, z_{r+1}]$. For $F \subset E_{r+1}$ put $\hat{F}^i = \{\hat{z}^i; z \in F\}$. For every $x \in E_r$ denote by $N_i(F, x)$ the number (possibly zero or infinite) of points in $\{z; z \in F, \hat{z}^i = x\}$.

2,1. Lemma. Let F be an analytic set in E_{r+1} . Then $N_i(F, x)$ is Lebesgue measurable with respect to x on E_r and

$$H_r(F) \geq \int_{E_r} N_i(F, x) \, dx.$$

Proof. Write F_{nk} for the set of all $z = [z_1, \dots, z_{r+1}] \in \hat{F}$ with $k \cdot 2^{-n} \leq z_i < (k+1) \cdot 2^{-n}$. Every F_{nk} is analytic and, consequently, \hat{F}_{nk}^i is Lebesgue measurable. Clearly, $H_r(\hat{F}_{nk}^i) \leq H_r(F_{nk})$. Denoting by χ_{nk} the characteristic function of \hat{F}_{nk}^i on E_r we obtain $\sum_{k=-\infty}^{\infty} \chi_{nk}(x) \nearrow N_i(F, x)$ ($n \rightarrow \infty$) and

$$\begin{aligned} \int_{E_r} N_i(F, x) dx &= \lim_{n \rightarrow \infty} \int_{E_r} \left(\sum_{k=-\infty}^{\infty} \chi_{nk}(x) \right) dx = \lim_{n \rightarrow \infty} \sum_{k=-\infty}^{\infty} H_r(\hat{F}_{nk}^i) \leq \\ &\leq \lim_{n \rightarrow \infty} \sum_{k=-\infty}^{\infty} H_r(F_{nk}) = H_r(F). \end{aligned}$$

3. Some background material. \mathbf{D} will be used to denote the set of all infinitely differentiable functions with compact support on E_{r+1} . Let \mathbf{A}_i be the system of all Lebesgue measurable sets $A \subset E_{r+1}$ with

$$+\infty > \|A\|_i = \sup_{\varphi \in \mathbf{D}} \int_A \frac{\partial \varphi(z)}{\partial z_i} dz, \quad \varphi \in \mathbf{D}, \quad \max_z |\varphi(z)| \leq 1.$$

A measurable set A belongs to \mathbf{A}_i if and only if such a finite signed Borel measure \mathbf{P}_i^A exists over the boundary \hat{A} of A that

$$\varphi \in \mathbf{D} \Rightarrow \int_{\hat{A}} \varphi d\mathbf{P}_i^A = \int_A \frac{\partial \varphi(z)}{\partial z_i} dz.$$

$\|A\|_i$ is equal to the variation of \mathbf{P}_i^A on A whenever $\hat{A} \in \mathbf{A}_i$. Further put $\mathbf{A} = \bigcap_{i=1}^{r+1} \mathbf{A}_i$.

\mathbf{A} is the system of all measurable $A \subset E_{r+1}$ for which the following is true: Such a vector-valued measure $\mathbf{P}^A = [\mathbf{P}_1^A, \dots, \mathbf{P}_{r+1}^A]$ exists over \hat{A} that

$$\int_{\hat{A}} v d\mathbf{P}^A \left(= \sum_{i=1}^{r+1} \int_{\hat{A}} v_i d\mathbf{P}_i^A \right) = \int_A \operatorname{div} v(z) dz$$

for every vector-valued function $v = [v_1, \dots, v_{r+1}]$ with $v_i \in \mathbf{D}$, $1 \leq i \leq r+1$. Writing \mathbf{V}^1 for the set of all $v = [v_1, \dots, v_{r+1}]$ with $v_i \in \mathbf{D}$ ($1 \leq i \leq r+1$), $|v(z)| = \left(\sum_{i=1}^{r+1} v_i^2(z) \right)^{\frac{1}{2}} \leq 1$ on E_{r+1} , we have for a measurable set $A \subset E_{r+1}$

$$+\infty > \|A\| = \sup_{v \in \mathbf{V}^1} \int_{\hat{A}} \operatorname{div} v(z) dz, \quad v \in \mathbf{V}^1,$$

if and only if $A \in \mathbf{A}$. $\|A\|$ coincides with the total variation of the vector-valued measure \mathbf{P}^A on \hat{A} whenever $A \in \mathbf{A}$. \mathbf{A}_i and \mathbf{A} are Boolean algebras.

\mathbf{A}_i includes all measurable sets A with $\int_{E_r} N_i(\hat{A}, x) dx < +\infty$. In particular, every $A \subset E_{r+1}$ with $H_r(\hat{A}) < +\infty$ belongs to \mathbf{A} and $\|A\| \leq H_r(\hat{A})$.

3,1. Remark. The systems \mathbf{A} and \mathbf{A}_i were, from different points of view, introduced by E. De Giorgi and J. Mařík. Their properties were studied by several authors. Interested reader is referred to [5] for a bibliography on the subject.

4. A set in E_{r+1} which can be represented as a union of a finite number of compact $(r+1)$ -dimensional intervals which are allowed to have a void interior will be called a figure. \bar{A} , A° , \dot{A} and $\text{diam } A$ will stand for the closure, the interior, the boundary and the diameter of A ($\subset E_{r+1}$) respectively. \mathbf{L} will denote the Lebesgue measure in E_{r+1} .

4.1. Lemma.* Let $A \subset E_{r+1}$ be a bounded set, $A^\circ \neq \emptyset$, $H_r(\dot{A}) < +\infty$. Then there exists a sequence of figures A_k ($k = 1, 2, \dots$) such that $\sup_k H_r(\dot{A}_k) < +\infty$ and

$$A_k \subset A_{k+1} \quad (k = 1, 2, \dots), \quad \bigcup_k A_k = A^\circ. **$$

Proof. For every positive integer p there exists a sequence $\{K_{jp}\}_{j=1}^\infty$ of open $(r+1)$ -dimensional cubes such that $\dot{A} \subset \bigcup_{j=1}^\infty K_{jp}$, $\text{diam } K_{jp} < \frac{1}{p}$ ($j = 1, 2, \dots$) and

$$\sum_j \text{diam}^r K_{jp} < 1 + cH_r(\dot{A}), \quad (5)$$

where $c > 0$ is a constant independent of p . Rearranging the sequence $\{K_{jp}\}_{j=1}^\infty$, if necessary, we can fix a $j(p)$ such that

$$\dot{A} \subset \bigcup_{j=1}^{j(p)} K_{jp}, \quad A \cap K_{jp} \neq \emptyset \quad \text{whenever} \quad j \in \langle 1, j(p) \rangle.$$

Denote by p_1 the least p with $A - \bigcup_{j=1}^{j(p)} K_{jp} \neq \emptyset$ and put $A_1 = A - \bigcup_{j=1}^{j(p_1)} K_{jp_1}$. Clearly, $A_1 \subset A^\circ$ and A_1 is a figure. Suppose now that figures $A_1 \subset \dots \subset A_k$ have already been constructed. Denote by p_{k+1} the least p for which $\bigcup_{j=1}^{j(p)} K_{jp}$ has a positive distance from A_k and put $A_{k+1} = A - \bigcup_{j=1}^{j(p_{k+1})} K_{jp_{k+1}}$. Repeating this procedure infinitely many times we arrive at a sequence of figures $A_k \nearrow A^\circ$ ($k \rightarrow \infty$). Taking (5) into account we see that $H_r(\dot{A}_k) \leq \sum_k H_r(\dot{K}_{jp_k}) \leq 2(r+1) \sum_j \text{diam}^r K_{jp_k} < 2(r+1) \times [1 + cH_r(\dot{A})]$ for every k . Thus the proof is complete.

4.2. Lemma. Let $A \subset E_{r+1}$ be a bounded set and suppose that there exist $A_k \in \mathbf{A}$ ($k = 1, 2, \dots$) such that $A_k \subset A$, $\lim_{k \rightarrow \infty} \mathbf{L}(A - A_k) = 0$, $\limsup_{k \rightarrow \infty} \|A_k\| = c < +\infty$.

* Cf. also [13], lemma 19, 26, p. 154.

** This will be expressed symbolically in the form $A_k \nearrow A^\circ$ ($k \rightarrow \infty$).

Then $\|A\| \leq c$ and

$$\int_{\bar{A}} v \, d\mathbf{P}^A = \lim_{k \rightarrow \infty} \int_{\bar{A}_k} v \, d\mathbf{P}^{A_k} \quad (6)$$

for every continuous $(r+1)$ -dimensional vector-valued function v on \bar{A} .*

Proof. We have for $v \in \mathbf{V}^1$

$$\begin{aligned} \int_{\bar{A}} \operatorname{div} v(z) \, dz &= \lim_{k \rightarrow \infty} \int_{\bar{A}_k} \operatorname{div} v(z) \, dz = \lim_{k \rightarrow \infty} \int_{\bar{A}_k} v \, d\mathbf{P}^{A_k} \leq \\ &\leq \limsup_{k \rightarrow \infty} \|A_k\| = c. \end{aligned}$$

Consequently, $\|A\| \leq c$.

Noticing that

$$\left| \int_{\bar{A}} v \, d\mathbf{P}^A - \int_{\bar{A}} \tilde{v} \, d\mathbf{P}^A \right| \leq \varepsilon c, \quad \left| \int_{\bar{A}_k} v \, d\mathbf{P}^{A_k} - \int_{\bar{A}_k} \tilde{v} \, d\mathbf{P}^{A_k} \right| \leq \varepsilon c$$

whenever v, \tilde{v} are continuous vector-valued functions on \bar{A} with $\max_{z \in \bar{A}} |v(z) - \tilde{v}(z)| \leq \varepsilon$,

we see at once that it is sufficient to prove (6) for $v = [v_1, \dots, v_{r+1}]$ with $v_i \in \mathbf{D}$ ($1 \leq i \leq r+1$) only. For such a v

$$\int_{\bar{A}} v \, d\mathbf{P}^A = \int_{\bar{A}} \operatorname{div} v(z) \, dz = \lim_{k \rightarrow \infty} \int_{\bar{A}_k} \operatorname{div} v(z) \, dz = \lim_{k \rightarrow \infty} \int_{\bar{A}_k} v \, d\mathbf{P}^{A_k}.$$

5. The scalar product of vectors $u, v \in E_n$ will be denoted by $u \cdot v$. Given $M \subset E_2$ we shall denote by $\mathbf{C}_n^{(0)}(M)$ the system of all continuous mappings of M into E_n . If M happens to be open, then $\mathbf{C}_n^{(1)}(M)$ will stand for the system of all $\Phi = [\Phi_1, \dots, \Phi_n] \in \mathbf{C}_n^{(0)}(M)$ whose components Φ_i ($1 \leq i \leq n$) have continuous first order partial derivatives in M . We shall write simply $\mathbf{C}_n^{(1)}$ instead of $\mathbf{C}_n^{(1)}(E_2)$ and ι will be used to denote the identity map of E_2 onto itself. $V(a, b)$ is the system of all rectifiable plane paths on $\langle a, b \rangle$, $V_0(a, b)$ is the subsystem of all $f \in V(a, b)$ with $(a) = f(b)$ (i. e. of all closed paths in $V(a, b)$).

5.1. Lemma. Let $f \in V(a, b)$, $\Psi \in \mathbf{C}_n^{(0)}([f])$, $\Phi \in \mathbf{C}_n^{(1)}(O)$, where O is some neighbourhood of $[f]$ in E_2 . Define the mapping $\chi = [\chi_1, \chi_2]$ of $[f]$ into E_2 by

$$\chi_1 = \Psi \circ \frac{\partial \Phi}{\partial x}, \quad \chi_2 = \Psi \circ \frac{\partial \Phi}{\partial y}. \quad ** \quad (7)$$

Then $\chi \in \mathbf{C}_2^{(0)}([f])$ and

$$\int_f \Psi \, d\Phi = \int_f \chi \, d\iota.$$

* This assertion was communicated to us by prof. J. Mařík, compare also [13], lemma 19, 21, pp. 150—151.

** We write $\frac{\partial \Phi}{\partial x} = \left[\frac{\partial \Phi_1}{\partial x}, \dots, \frac{\partial \Phi_n}{\partial x} \right]$ for $\Phi(x, y) = \Phi = [\Phi_1, \dots, \Phi_n]$; $\frac{\partial \Phi}{\partial y}$ has a similar meaning.

Proof. Let $f = [f_1, f_2]$ and put $t_k^m = a + k(b - a)m^{-1}$, $z_k^m = f(t_k^m)$ ($k = 0, \dots, m$; $m = 1, 2, \dots$). It is easily seen that $\Phi(f)$ is rectifiable on $\langle a, b \rangle$ and

$$\begin{aligned} \Phi(z_k^m) - \Phi(z_{k-1}^m) &= \frac{\partial(\Phi z_{k-1}^m)}{\partial x} [f_1(t_k^m) - f_1(t_{k-1}^m)] + \\ &+ \frac{\partial\Phi(z_{k-1}^m)}{\partial y} [f_2(t_k^m) - f_2(t_{k-1}^m)] + [f(t_k^m) - f(t_{k-1}^m)] \cdot \vec{o}_{mk}, \end{aligned}$$

where $\max_k |\vec{o}_{mk}| \rightarrow 0$ as $m \rightarrow \infty$. Hence

$$\begin{aligned} \int_f \Psi \, d\Phi &= \lim_{m \rightarrow \infty} \sum_{k=1}^m \Psi(z_{k-1}^m) \circ [\Phi(z_k^m) - \Phi(z_{k-1}^m)] = \\ &= \lim_{m \rightarrow \infty} \sum_{k=1}^m \chi(z_{k-1}^m) \circ [t(z_k^m) - t(z_{k-1}^m)] = \int_f \chi \, dt. \end{aligned}$$

6. In section 3 we have recalled some basic properties of the systems \mathbf{A}_i and \mathbf{A} of subsets in E_{r+1} . Since no simplification could have been acquired by specialization to $r = 1$, we described the general situation for any $r \geq 1$. However, the special case $r = 1$ is the only one we shall deal with in the sequel. Let us agree that, from now on, the systems \mathbf{A}_i and \mathbf{A} will be considered with respect to E_2 only. (Thus every set of \mathbf{A}_i , \mathbf{A} to be met below is a subset in E_2 .) Further denote by $\tilde{\mathbf{A}}$ the subsystem of all $A \in \mathbf{A}$ whose boundary \dot{A} is compact.

6.1. Definition. Let $A \in \tilde{\mathbf{A}}$, $\Phi \in \mathbf{C}_n^{(1)}$, $\Psi \in \mathbf{C}_n^{(0)}(\dot{A})$.

We put

$$P(A, \Phi, \Psi) = \int_{\dot{A}} \tilde{\chi} \, d\mathbf{P}^A,$$

where $\tilde{\chi} = [-\chi_2, \chi_1]$ and χ_1, χ_2 are defined by (7).

6.2. Lemma. Let $f \in V_0(a, b)$, $A \subset E_2$ and suppose that

$$\{z; \text{ind}(z; f) = 1\} = A, \{z; \text{ind}(z; f) = 0\} = E_2 - \bar{A}.$$

Then $A \in \tilde{\mathbf{A}}$ and

$$P(A, \Phi, \Psi) = \int_f \Phi \, d\Psi$$

whenever $\Phi, \Psi, \frac{\partial\Phi}{\partial x}, \frac{\partial\Phi}{\partial y} \in \mathbf{C}_n^{(1)}$.

Proof. Since $\dot{A} \subset [f]$ and f is rectifiable, we have $H_1(\dot{A}) < +\infty$. Consequently, $A \in \tilde{\mathbf{A}}$. Using Green's formula (cf. [6]) and lemma 5.1 we obtain

$$P(A, \Phi, \Psi) = \int_{\dot{A}} \tilde{\chi} \, d\mathbf{P}^A = \iint_{\dot{A}} \text{div} \tilde{\chi} = - \iiint_{\dot{A}} \left(\frac{\partial\chi_2}{\partial x} - \frac{\partial\chi_1}{\partial y} \right) = - \int_f \chi \, dt = - \int_f \Psi \, d\Phi.$$

Finally, integration by parts for Stieltjes integrals yields $-\int_f \Psi d\Phi = \int_f \Phi d\Psi$.

6,3. Remark. In 6,2, the assumption $\frac{\partial\Phi}{\partial x}, \frac{\partial\Phi}{\partial y}, \Psi \in \mathbf{C}_n^{(1)}$ could be generalized to $\Psi \in \mathbf{C}_n^{(0)}(A), \Phi \in \mathbf{C}_n^{(1)}$. As lemma 6,2 shows, $P(A, \Phi, \Psi)$ can be considered as an analogue of $\int_f \Phi d\Psi$. Indeed, if f is a positively oriented rectifiable simple closed curve bounding A , then these two quantities coincide with each other.

6,4. Lemma. Let $A \in \tilde{\mathbf{A}}$ and let $\Phi, \Psi \in \mathbf{C}_n^{(1)}$. Then

$$P(A, \Phi, \Psi) = -P(A, \Psi, \Phi).$$

Proof. Since $P(A, \dots) = -P(E_2 - A, \dots)$, we may assume that A is bounded. Let us recall that for $h \in \mathbf{C}_1^{(1)}$ and a solenoidal vector-valued function $v \in \mathbf{C}_2^{(1)}$ the formula

$$\int_A hv d\mathbf{P}^A = \iint_A \text{grad } h \circ v \quad (8)$$

is true (cf. [7], theorem 48, p. 554). Applying (8) to $h = \Psi_i, v = \left[-\frac{\partial\Phi_i}{\partial y}, \frac{\partial\Phi_i}{\partial x} \right]$ ($i = 1, \dots, n$), we obtain

$$P(A, \Phi, \Psi) = \sum_{i=1}^n \iint_A \left(-\frac{\partial\Psi_i}{\partial x} \cdot \frac{\partial\Phi_i}{\partial y} + \frac{\partial\Psi_i}{\partial y} \cdot \frac{\partial\Phi_i}{\partial x} \right).$$

In a similar way

$$P(A, \Psi, \Phi) = \sum_{i=1}^n \iint_A \left(-\frac{\partial\Phi_i}{\partial x} \cdot \frac{\partial\Psi_i}{\partial y} + \frac{\partial\Phi_i}{\partial y} \cdot \frac{\partial\Psi_i}{\partial x} \right).$$

whence our lemma follows at once.

7. Given a $M \subset E_2$ and a mapping Φ of M into E_n we put for any $N \subset M$

$$\|\Phi\|_N = \sup_{z \in N} |\Phi(z)|.$$

We say that Φ is Lipschitzian on N with constant λ provided $|\Phi(u) - \Phi(v)| \leq \lambda |u - v|$ whenever $u, v \in N$.

7,1. Definition. Let $A \in \tilde{\mathbf{A}}, \Psi \in \mathbf{C}_n^{(0)}(A)$. We define

$$\alpha(A, \Psi) = \sup_{\Phi} P(A, \Phi, \Psi),$$

Φ ranging over the class of all $\Phi \in \mathbf{C}_n^{(1)}$ with $\|\Phi\|_{E_2} \leq 1$.

7,2. Lemma. Let $A \in \tilde{\mathbf{A}}, \Psi = [\Psi_1, \dots, \Psi_n] \in \mathbf{C}_n^{(1)}$ and suppose that

$$\left| \frac{\partial\Psi_i(z)}{\partial x} \right| \leq \lambda, \quad \left| \frac{\partial\Psi_i(z)}{\partial y} \right| \leq \lambda \quad (i = 1, \dots, n)$$

whenever $z \in \dot{A}$. Then, for every $\Phi \in \mathbf{C}_n^{(1)}$,

$$|P(A, \Phi, \Psi)| \leq \lambda \sqrt{2} \|\Phi\|_{\dot{A}} \cdot \|A\|. \quad (9)$$

In particular,

$$\alpha(A, \Psi) \leq \lambda \sqrt{2} \|A\|. \quad (10)$$

Proof. Writing $v = \left[-\Phi, \frac{\partial \Psi}{\partial y}, \Phi \circ \frac{\partial \Psi}{\partial x} \right]$ we obtain by 6,4 and 6,1

$$|P(A, \Phi, \Psi)| = |P(A, \Psi, \Phi)| = \left| \int_{\dot{A}} v \, d\mathbf{P}^A \right| \leq \|v\|_{\dot{A}} \cdot \|A\|.$$

Clearly, $\|v\|_{\dot{A}} \leq \lambda \sqrt{2} \|\Phi\|_{\dot{A}}$.

7,3. Lemma. Let h be a function which is Lipschitzian on E_2 with constant λ . Then there exists a sequence of functions $h_k \in \mathbf{C}_1^{(1)}$ ($k = 1, 2, \dots$) such that $h_k \rightarrow h$ uniformly on E_2 as $k \rightarrow \infty$ and

$$\left| \frac{\partial h_k}{\partial x} \right| \leq \lambda, \quad \left| \frac{\partial h_k}{\partial y} \right| \leq \lambda \quad (k = 1, 2, \dots).$$

Proof. This lemma is well known.

7,4. Proposition. Let $A \in \tilde{\mathbf{A}}$ and let Ψ be a mapping of \dot{A} into E_n , which is Lipschitzian on \dot{A} with constant λ . Then (10) is valid.

Proof. We may assume that $\Psi = [\Psi_1, \dots, \Psi_n]$, where Ψ_i ($1 \leq i \leq n$) are Lipschitzian on E_2 with constant λ (cf. [8], lemma 1, p. 341). According to 7,2 we have a sequence $\Psi_i^k \in \mathbf{C}_1^{(1)}$ ($k = 1, 2, \dots$) such that $\Psi_i^k \rightarrow \Psi_i$ ($k \rightarrow \infty$) uniformly on E_2 and $\left| \frac{\partial \Psi_i^k}{\partial x} \right| \leq \lambda$, $\left| \frac{\partial \Psi_i^k}{\partial y} \right| \leq \lambda$. Put $\Psi^k = [\Psi_1^k, \dots, \Psi_n^k]$. Clearly, $\Psi^k \in \mathbf{C}_n^{(1)}$ and, in view of 7,2, $|P(A, \Phi, \Psi^k)| \leq \lambda \sqrt{2} \|\Phi\|_{\dot{A}} \cdot \|A\|$ for an arbitrary $\Phi \in \mathbf{C}_n^{(1)}$. Making $k \rightarrow \infty$ we obtain (9) (cf. the definition 6,1). Hence (10) easily follows.

8,1. Lemma. Let $A \in \tilde{\mathbf{A}}$ and suppose that $\Psi \in \mathbf{C}_n^{(0)}(\dot{A})$, $\alpha(A, \Psi) < \infty$. Then, for every $\Phi \in \mathbf{C}_n^{(1)}$,

$$|P(A, \Phi, \Psi)| \leq \|\Phi\|_{\dot{A}} \cdot \alpha(A, \Psi).$$

Proof. Given $\varepsilon > 0$ and $\Phi \in \mathbf{C}_n^{(1)}$ we can fix a $\tilde{\Phi} \in \mathbf{C}_n^{(1)}$ such that $\|\tilde{\Phi}\|_{E_2} \leq \varepsilon + \|\Phi\|_{\dot{A}}$ and $\tilde{\Phi} = \Phi$ in some neighbourhood of \dot{A} (cf. lemma 5 in [7]). According to the definition 7,1 we have $|P(A, \Phi, \Psi)| = |P(A, \tilde{\Phi}, \Psi)| \leq \|\tilde{\Phi}\|_{E_2} \cdot \alpha(A, \Psi) \leq (\varepsilon + \|\Phi\|_{\dot{A}}) \cdot \alpha(A, \Psi)$. Since ε was an arbitrary positive number, the proof is complete.

8,2. Remark. Let $A \in \tilde{\mathbf{A}}$, $\Psi \in \mathbf{C}_n^{(0)}(\dot{A})$, $\alpha(A, \Psi) < +\infty$. Fix $\Phi \in \mathbf{C}_n^{(0)}(A)$ and suppose that $\Phi^k \in \mathbf{C}_n^{(1)}$ ($k = 1, 2, \dots$),

$$\lim_{k \rightarrow \infty} \|\Phi - \Phi^k\|_{\dot{A}} = 0. \quad (11)$$

It follows easily from 8.1 that the limit $\lim_{k \rightarrow \infty} P(A, \Phi^k, \Psi)$ exists and is independent of the choice of the sequence $\{\Phi^k\}_{k=1}^\infty$ fulfilling (11). We are thus justified to introduce the following definition:

8.3. Definition. Let $A \in \tilde{\mathbf{A}}$, $\Psi \in \mathbf{C}_n^{(0)}(\dot{A})$, $\alpha(A, \Psi) < +\infty$. For any $\Phi \in \mathbf{C}_n^{(0)}(\dot{A})$ put

$$P(A, \Phi, \Psi) = \lim_{k \rightarrow \infty} P(A, \Phi^k, \Psi),$$

where $\{\Phi^k\}_{k=1}^\infty$ is a sequence of mappings in $\mathbf{C}_n^{(1)}$ fulfilling (11).

9. The symbols $f^1, \dots, f^m, C, \omega, G_p, G$ will have the same meaning as in the theorem 1.1. Further put $U_l = \{z; z \in E_2 - C, \omega(z) \geq l\}$.

9.1. Lemma.

$$\sum_{l=-\infty}^{\infty} \|U_l\| < +\infty, \quad \sum_{p=-\infty}^{\infty} \|G_p\| < +\infty. \quad (12)$$

In particular, $U_p, G_p \in A$ for every integer p and, by proposition 7.4,

$$\sum_{l=-\infty}^{\infty} \alpha(U_l, \Psi) < +\infty, \quad \sum_{p=-\infty}^{\infty} \alpha(G_p, \Psi) < +\infty \quad (13)$$

for every Lipschitzian mapping Ψ of C into E_n .

Proof. Since $\sum_l \|U_l\|_i \leq \sum_l \int_{E_1} N_i(\dot{U}_l, y) dy$, $\sum_p \|G_p\|_i \leq \sum_p \int_{E_1} N_i(\dot{G}_p, y) dy$, it is sufficient to prove that the functions

$$\sum_l N_i(\dot{U}_l, y), \quad \sum_p N_i(\dot{G}_p, y) \quad (i = 1, 2)$$

are integrable (with respect to the variable y) on E_1 . Clearly, we may consider the case $i = 2$ only. Let us keep the notation introduced in [2], section 15, pp. 589–591. From investigations described there we obtain for every $y \in E_1 - M$

$$N_2(\dot{U}_l, y) \leq \sum_{j=1}^n |s_y(U_l, u_j)|, \quad \sum_l \sum_{j=1}^n |s_y(U_l, u_j)| \leq \psi(y).$$

Noticing that M has measure zero and ψ is integrable on E_1 we see that integrability of $\sum_l N_2(\dot{U}_l, y)$ is checked. Similarly, investigations described in [2], p. 592, imply the inequality

$$\sum_p N_2(\dot{G}_p, y) \leq 2\psi(y) \quad (y \in E_1 - M)$$

showing that $\sum_p N_2(\dot{G}_p, y)$ is integrable on E_1 .

9,2. Theorem. Let $\Phi \in \mathbf{C}_n^{(0)}(C)$ and let Ψ be a Lipschitzian mapping of C into E_n . Then

$$\sum_{k=1}^m \int_{f^k} \Phi \, d\Psi = \sum_{l=1}^{\infty} \left\{ \sum_{p \geq l} [P(G_p, \Phi, \Psi) - P(G_{-p}, \Phi, \Psi)] \right\}. \quad (14)$$

Proof. We shall first prove

$$\sum_{k=1}^m \int_{f^k} \Phi \, d\Psi = \sum_{l=1}^{\infty} [P(U_l, \Phi, \Psi) + P(U_{1-l}, \Phi, \Psi)], \quad (15)$$

$$P(U_l, \Phi, \Psi) + P(U_{1-l}, \Phi, \Psi) = \sum_{p \geq l} [P(G_p, \Phi, \Psi) - P(G_{-p}, \Phi, \Psi)], \quad (16)$$

whence (14) follows at once. In view of (13) we may assume that $\Phi \in \mathbf{C}_n^{(1)}$ (cf. also 8,2). Define $\chi = [\chi_1, \chi_2]$ by (7). We obtain from 5,1

$$\sum_{k=1}^m \int_{f^k} \Phi \, d\Psi = - \sum_{k=1}^m \int_{f^k} \chi \, dI. \quad (18)$$

Keeping the notation introduced in [2], section 17, we derive from theorem 15 and remark 17 in [2]

$$\sum_{k=1}^m \int_{f^k} \chi \, dI = \sum_{l=1}^{\infty} [P_0(U_l, \chi) + P_0(U_{1-l}, \chi)]. \quad (15^{\text{bis}})$$

In a similar way we obtain from investigations on p. 592 in [2]

$$P_0(U_l, \chi) + P_0(U_{1-l}, \chi) = \sum_{p \geq l} [P_0(G_p, \chi) - P_0(G_{-p}, \chi)]. \quad (16^{\text{bis}})$$

Comparing the definition 6,1 of the present note with the remark 17 in [2] we see that

$$P_0(U_l, \chi) = - \int_{\dot{U}_l} \tilde{\chi} \, d\mathbf{P}^{U_l} = -P(U_l, \Phi, \Psi),$$

$$P_0(G_p, \chi) = -P(G_p, \Phi, \Psi).$$

Thus (15^{bis}), (16^{bis}) and (18) imply (15), (16).

10,1. Lemma. Let $A \subset E_2$ be a bounded set, $A_k \subset A$ ($k = 1, 2, \dots$) and suppose that

$$\lim_{k \rightarrow \infty} \mathbf{L}(A - A_k) = 0, \quad \limsup_{k \rightarrow \infty} \|A_k\| < +\infty.$$

Let Ψ be a Lipschitzian mapping of \bar{A} into E_n .

Then

$$\alpha(A, \Psi) < +\infty, \quad \limsup_{k \rightarrow \infty} \alpha(A_k, \Psi) < +\infty \quad (19)$$

and, for every $\Phi \in \mathbf{C}_n^{(0)}(\bar{A})$,

$$\lim_{k \rightarrow \infty} P(A_k, \Phi, \Psi) = P(A, \Phi, \Psi). \quad (20)$$

Proof. By lemma 4,2, (20) is true for any $\Phi \in \mathbf{C}_n^{(1)}$ (cf. 6,1). In view of 7,4 we obtain (19). Hence it follows easily that (20) can be extended, by continuity, to any $\Phi \in \mathbf{C}_n^{(0)}(\bar{A})$.

10,2. Definition. Let $A \subset E_2$ be a bounded set, $\mathbf{L}(A) = 0$. Let γ be a function defined almost everywhere on A such that the Lebesgue integral $\iint_K \gamma$ is available for every rectangle $K \subset A^\circ$. (Consequently, $\iint_B \gamma$ exists for every two-dimensional figure $B \subset A^\circ$ as well.) If

$$\lim_{k \rightarrow \infty} \iint_{A_k} \gamma \quad (21)$$

exists for every sequence of figures $A_k \nearrow A^\circ$ ($k \rightarrow \infty$) with

$$\sup_k H_1(A_k) < +\infty, \quad (22)$$

then the limit (21) is independent of the choice of figures $A_k \rightarrow A^\circ$ fulfilling (22) and its value will be denoted by $\mathbf{L}(A, \gamma)$.

10,3. Remark. Of course, $\mathbf{L}(A, \gamma) = \iint_A \gamma$ whenever γ happens to be Lebesgue integrable on A , so that $\mathbf{L}(A, \gamma)$ may be considered as an extension of the Lebesgue integral. For more general study of analogous extensions the reader may consult [9].

The articles [10], [11] reviewed in Ref. jour. 1959 which seem to deal with similar problems were not available to us.

10,4. Proposition. Let $A \subset E_2$ be a bounded set, $H_1(A) < +\infty$. Let $\Phi \in \mathbf{C}_n^{(0)}(\bar{A})$ and suppose that Ψ is a Lipschitzian mapping of \bar{A} into E_n .

If $\gamma = \text{rot}(\Phi, \Psi)$ in A° , then

$$P(A, \Phi, \Psi) = \mathbf{L}(A, \gamma).$$

This proposition follows easily from 10,1 and 10,2.

11. As an easy consequence of 9,2 and 10,4 we obtain the following theorem.

11,1. Theorem. Let us keep all the assumptions and notation of the theorem 1,1. Then

$$\sum_{k=1}^m \int_{J^k} \Phi \, d\Psi = \sum_{l=1}^m \left\{ \sum_{p \geq l} [\mathbf{L}(G_p, \gamma) - \mathbf{L}(G_{-p}, \gamma)] \right\}. \quad (23)$$

11,2. Remark. Theorem 1,1 is merely a corollary of 11,1.

11,3. Remark. The right hand side in (23) may be replaced by $\sum_{p=1}^n p[\mathbf{L}(G_p, \gamma) - \mathbf{L}(G_{-p}, \gamma)]$ if this series happens to converge.

11,4. Remark. Let $\Psi = [\Psi_1, \dots, \Psi_n]$ be a Lipschitzian mapping of G into E_n and let $\Gamma = [\Gamma_1, \dots, \Gamma_n]$ be a Lipschitzian mapping of $V = \Psi(G)$ into E_n . Suppose

that $M \subset V$, $\mathbf{L}\Psi^{-1}(M) = 0$ and that $\sum_{k=1}^n \tau_k^i(u^0) (u_k - u_k^0)$ is the differential of Γ^i with respect to V at any $u^0 = [u_1^0, \dots, u_n^0] \in V - M$. Put $\Phi(z) = \Gamma(\Psi(z))$,

$$\gamma(z) = \sum_{\substack{i, k=1 \\ i < k}}^n [\tau_i^k(\Psi(z)) - \tau_k^i(\Psi(z))] \cdot \begin{vmatrix} \frac{\partial \Psi_i(z)}{\partial x}, & \frac{\partial \Psi_i(z)}{\partial y} \\ \frac{\partial \Psi_k(z)}{\partial x}, & \frac{\partial \Psi_k(z)}{\partial y} \end{vmatrix}$$

as far as the symbols involved are meaningful. Then $\gamma = \text{rot}(\Phi, \Psi)$ in G .

This follows at once from theorem 12 in [12].

This assertion can be combined with 11,1.

REFERENCES

- [1] Radó T., *Length and area*, New York 1948.
- [2] Král J., *О криволинейных интегралах на плоскости*, Чех. мат. жур. т. 7 (82), 584—598.
- [3] Whitney H., *Geometric integration theory*, Princeton 1957 (русский перевод: *Геометрическая теория интегрирования*, Москва 1960).
- [4] Krickeberg K., *Über den Gausschen und den Stokesschen Integralsatz III*, Math. Nachrichten 12 (1954), 341—365.
- [5] Král J., *Poznámka o množinách, jejichž charakteristická funkce má za parciální derivaci zobecněnou míru*, Čas. pro pěst. mat. 86 (1961), 178—194.
- [6] Král J. und Mařík J., *Der Greensche Satz*, Čech. mat. žur. 7 (82), 1957, 235—247.
- [7] Mařík J., *The surface integral*, Čech. mat. žur. 6 (81), 1956, 522—557.
- [8] Radó T. and Reichelderfen P. V., *Continuous transformations in analysis*, Berlin-Göttingen-Heidelberg 1955.
- [9] Mařík J., *Uneigentliche mehrfache Integrale*, Wiss. Zeitschr. Humboldt-Univ. Berlin, Math.-Nat. Reihe X (1961) 3/4, 413—414.
- [10] Лепина Э. И., *Несобственные интегралы в n-мерном пространстве*, Уч. зап. Латв. ун-та, 1958, 20, 105—124 (см. Реф. жур. 1959, реф. 7052).
- [11] Лепина Э. И., *Обобщенная формула Грина*, *ibid.* 125—135 (см. Реф. жур. 1959, реф. 7053).
- [12] Král J., *On Lipschitzian mappings from the plane into Euclidean spaces*, Čech. mat. žur. 8 (83), 1958, 257—266.
- [13] Černý J.—Mařík J., *Integrální počet II*, Praha 1961.

Received May 17, 1962.

*Katedra matematické analýzy Karlovy university v Praze
Katedra matematiky Slovenskej vysokej školy technickej
v Bratislave*

ЗАМЕТКА К ТЕОРЕМЕ СТОКЕСА ДЛЯ ДВУМЕРНЫХ ИНТЕГРАЛОВ
В n -МЕРНОМ ПРОСТРАНСТВЕ

Иосеф Крал, Белослав Риечан

Выводы

Если f — непрерывное отображение отрезка $\langle a, b \rangle$ в плоскость E_2 , то символом $v(a, b, f)$ обозначим длину пути f , определенную обыкновенным образом. Если $v(a, b, f) < +\infty$, то мы будем говорить, что $f \in V(a, b)$. Подсистему всех $f \in V(a, b)$ удовлетворяющих условию $f(a) = f(b)$, обозначим символом $V_0(a, b)$. Если $f \in V(a, b)$ и $\Phi = [\Phi_1, \dots, \Phi_n]$, $\Psi = [\Psi_1, \dots, \Psi_n]$ — непрерывные отображения множества $f(\langle a, b \rangle)$ в пространство E_n , то полагаем по определению

$$\int_f \Phi d\Psi = \sum_{i=1}^n \int_a^b \Phi_i(f(t)) d\Psi_i(f(t))$$

в предложении, что существуют соответствующие интегралы Стильтьеса. Словом „интервал“ мы будем подразумевать двумерный компактный интервал. Если K — интервал, то обозначим f_K отображение из $V_0(a, b)$, которое даст параметрическое представление контура K (описываемого в положительном направлении при изменении параметра от a до b). Пусть теперь G — открытое множество в E_2 и пусть Φ, Ψ — непрерывные отображения множества G в пространство E_n . Пусть, далее, γ — функция определенная почти всюду на G и интегрируемая по Лебегу на каждом интервале $K \subset G$. Будем говорить, что $\gamma = \text{rot}(\Phi, \Psi)$ на G , если для каждого интервала $K \subset G$ справедливо равенство $\int_{f_K} \Phi d\Psi = \int_K \gamma$. Множество, которое является соединением конечного числа интервалов будем называть фигурой.

Символом H_1, L обозначим линейную меру Хаусдорфа и двумерную меру Лебега соответственно. Пусть A — ограниченное множество в E_2 , \dot{A} — его граница. $A^\circ = A - \dot{A}$. Пусть, далее, $L(A) = 0$ и пусть γ — функция, определенная почти всюду на A и интегрируемая на каждом интервале $K \subset A^\circ$. Если для каждой последовательности фигур $F_k \subset A^\circ$, удовлетворяющей требованию $\sup H_1(\dot{F}_k) < \infty$ существует предел $\lim_{k \rightarrow \infty} \int_{F_k} \gamma$, то этот предел не зависит от последовательности $\{F_k\}_{k=1}^\infty$ и мы его обозначим через $\mathbf{L}(A, \gamma)$; разумеется $\mathbf{L}(A, \gamma) = \int_A \gamma$ если γ интегрируемая на A .

Теорема. Пусть $f^j \in V_0(a_j, b_j)$ ($1 \leq j \leq m$), $C = \bigcup_{j=1}^m f^j(\langle a_j, b_j \rangle)$. Для $z \in E_2 - C$ положим $\omega(z) = \sum_{j=1}^m \text{ind}(z, f^j)$, где $\text{ind}(z, f^j)$ обозначает порядок точки z относительно пути f^j . Пусть $G_p = \{z; z \in E_2 - C, \omega(z) = p\}$, $G = \bigcup_{p \neq 0} G_p$ и пусть на $C \cup G$ определены непрерывные отображения Φ, Ψ в пространство E_n , причем Ψ удовлетворяет условию Липшица. Если $\gamma = \text{rot}(\Phi, \Psi)$ на G , тогда существуют несобственные интегралы $\mathbf{L}(G_p, \gamma)$ ($p \neq 0$) и имеет место формула

$$\sum_{j=1}^m \int_{f^j} \Phi d\Psi = \sum_{l=1}^l \left\{ \sum_{p \geq l} [\mathbf{L}(G_p, \gamma) - \mathbf{L}(G_{-p}, \gamma)] \right\}. \quad (*)$$

Правую часть равенства (*) можно заменить на $\sum_{p=1}^l p[\mathbf{L}(G_p, \gamma) - \mathbf{L}(G_{-p}, \gamma)]$ соотв. на $\int_G \gamma \omega$, если последние символы имеют смысл.