

Recep Şahin; Osman Bizim; I. N. Cangul

Commutator subgroups of the extended Hecke groups $\bar{H}(\lambda_q)$

Czechoslovak Mathematical Journal, Vol. 54 (2004), No. 1, 253–259

Persistent URL: <http://dml.cz/dmlcz/127882>

Terms of use:

© Institute of Mathematics AS CR, 2004

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

COMMUTATOR SUBGROUPS OF THE EXTENDED
HECKE GROUPS $\overline{H}(\lambda_q)$

R. SAHIN, Balikesir, O. BIZIM and I. N. CANGUL, Bursa

(Received July 9, 2001)

Abstract. Hecke groups $H(\lambda_q)$ are the discrete subgroups of $\text{PSL}(2, \mathbb{R})$ generated by $S(z) = -(z + \lambda_q)^{-1}$ and $T(z) = -1/z$. The commutator subgroup of $H(\lambda_q)$, denoted by $H'(\lambda_q)$, is studied in [2]. It was shown that $H'(\lambda_q)$ is a free group of rank $q - 1$.

Here the extended Hecke groups $\overline{H}(\lambda_q)$, obtained by adjoining $R_1(z) = 1/\bar{z}$ to the generators of $H(\lambda_q)$, are considered. The commutator subgroup of $\overline{H}(\lambda_q)$ is shown to be a free product of two finite cyclic groups. Also it is interesting to note that while in the $H(\lambda_q)$ case, the index of $H'(\lambda_q)$ is changed by q , in the case of $\overline{H}(\lambda_q)$, this number is either 4 for q odd or 8 for q even.

Keywords: Hecke group, extended Hecke group, commutator subgroup

MSC 2000: 11F06, 20H05, 20H10

1. INTRODUCTION

In [4], Erich Hecke introduced the groups $H(\lambda)$ generated by two linear fractional transformations

$$T(z) = -\frac{1}{z} \quad \text{and} \quad U(z) = z + \lambda,$$

where λ is a fixed positive real number. T and U have matrix representations

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & \lambda \\ 0 & 1 \end{pmatrix},$$

respectively. (In this work we identify each matrix A with $-A$, so that they each represent the same transformation). Let $S = T.U$, i.e.

$$S(z) = -\frac{1}{z + \lambda}.$$

E. Hecke showed that $H(\lambda)$ is Fuchsian if and only if $\lambda = \lambda_q = 2 \cos \frac{\pi}{q}$, where q is an integer $q \geq 3$ or $\lambda > 2$ is real. In these two cases $H(\lambda)$ is called a Hecke group. We consider the former case. Then the Hecke group $H(\lambda)$ is the discrete subgroup of $\text{PSL}(2, \mathbb{R})$ generated by S and U , where

$$U(z) = z + \lambda_q$$

and it has a presentation $H(\lambda) = \langle T, S \mid T^2 = S^q = I \rangle$.

The most important and studied Hecke group is the modular group $H(\lambda_3)$. In this case $\lambda_3 = 2 \cos \frac{\pi}{3} = 1$, i.e. all coefficients of the elements of $H(\lambda_3)$ are rational integers. In the literature, the symbols Γ and $\Gamma(1)$ are used to denote the modular group. In this paper we shall use $H(\lambda_3)$ for this purpose. The next two most important Hecke groups are those for $q = 4$ and $q = 6$, in which cases $\lambda_q = \sqrt{2}$ and $\sqrt{3}$, respectively.

The extended modular group $\overline{H}(\lambda_3)$ has a presentation

$$\overline{H}(\lambda_3) = \langle R_1, R_2, R_3 \mid R_1^2 = R_2^2 = R_3^2 = (R_1 R_2)^3 = (R_3 R_1)^2 = I \rangle$$

where

$$R_1(z) = \frac{1}{\bar{z}}, \quad R_2(z) = \frac{-1}{\bar{z} + 1}, \quad R_3(z) = -\bar{z}.$$

The modular group is a subgroup of index 2 in $\overline{H}(\lambda_3)$ (see [3]). It has a presentation

$$H(\lambda_3) = \langle T, S \mid T^2 = S^3 = I \rangle \cong C_2 * C_3,$$

where

$$T = R_3 R_1 = R_1 R_3, \quad S = R_1 R_2.$$

Putting $R = R_1$, we have

$$\overline{H}(\lambda_3) = \langle T, S, R \mid T^2 = S^3 = R^2 = I, RT = TR, RS = S^{-1}R \rangle.$$

Similarly the extended Hecke group $\overline{H}(\lambda_q)$ has a presentation

$$\overline{H}(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, RT = TR, RS = S^{-1}R \rangle$$

and Hecke group $H(\lambda_q)$ is a subgroup of index 2 in $\overline{H}(\lambda_q)$.

The commutator subgroup of G is denoted by G' and defined by

$$\langle [g, h] \mid g, h \in G \rangle$$

where $[g, h] = ghg^{-1}h^{-1}$. Since G' is a normal subgroup of G , we can form the factor-group G/G' which is the largest abelian quotient group of G .

In this work we obtain some results concerning commutator subgroups of the extended Hecke group $\overline{H}(\lambda_q)$.

2. COMMUTATOR SUBGROUPS OF THE EXTENDED HECKE GROUP $\overline{H}(\lambda_q)$

The commutator subgroup of the Hecke group $H(\lambda_q)$ is denoted by $H'(\lambda_q)$. We have

$$T^2 = S^q = I, \quad TS = ST$$

in $H(\lambda_q)/H'(\lambda_q)$. So one can find

$$H(\lambda_q)/H'(\lambda_q) \cong C_2 \times C_q$$

and hence it is isomorphic to C_{2q} if q is odd. Therefore

$$|H(\lambda_q) : H'(\lambda_q)| = 2q.$$

If q is even, $(TS)^q = 1$ while if q is odd, $(TS)^{2q} = 1$. In particular, $H'(\lambda_q)$ is a free group of rank $q - 1$ (see [1]).

By [5], the Reidemeister-Schreier method gives the generators of $H'(\lambda_q)$ as

$$a_1 = TSTS^{q-1}, \quad a_2 = TS^2TS^{q-2}, \quad \dots, \quad a_{q-1} = TS^{q-1}TS.$$

Similarly for the extended Hecke group $\overline{H}(\lambda_q)$ we have

$$T^2 = S^q = R^2 = I, \quad RT = TR, \quad RS = S^{-1}R, \quad RS = SR, \quad TS = ST$$

in $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q)$.

Theorem 1. *Let q be odd, then*

- (i) $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong V_4 \cong C_2 \times C_2$
- (ii) $\overline{H}'(\lambda_q) = \langle S, TST \mid S^q = (TST)^q = I \rangle \cong C_q * C_q$.

Proof. (i) Since the extended Hecke group $\overline{H}(\lambda_q)$ has a presentation

$$\overline{H}(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, \quad RT = TR, \quad RS = S^{-1}R \rangle$$

and

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, S, R \mid T^2 = S^q = R^2 = I, \quad RT = TR, \quad RS = S^{-1}R, \\ RS = SR, \quad TS = ST \rangle$$

one has $RS = S^{-1}R$ and $RS = SR$, and thus

$$S^{q-2} = S^q = S^2 = I.$$

This shows that $S = I$, as q is odd. Thus

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, R \mid T^2 = R^2 = (TR)^2 = I \rangle$$

and finally

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong V_4 \cong C_2 \times C_2.$$

(ii) Now we determine the set of generators for $\overline{H}'(\lambda_q)$. We choose a Schreier transversal for $\overline{H}'(\lambda_q)$ as

$$I, T, R, TR.$$

According to the Reidemeister-Schreier method, we can form all possible products

$$\begin{array}{lll} I \cdot T \cdot (T)^{-1} = I, & I \cdot S \cdot (I)^{-1} = S, & I \cdot R \cdot (R)^{-1} = I, \\ T \cdot T \cdot (I)^{-1} = I, & T \cdot S \cdot (T)^{-1} = TST, & T \cdot R \cdot (TR)^{-1} = I, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & R \cdot S \cdot (R)^{-1} = RSR, & R \cdot R \cdot (I)^{-1} = I, \\ TR \cdot T \cdot (R)^{-1} = TRTR, & TR \cdot S \cdot (TR)^{-1} = TRSRT, & TR \cdot R \cdot (T)^{-1} = I. \end{array}$$

Since

$$\begin{array}{l} RTRT = I, \\ TRTR = I, \\ RSR = S^{-1}, \\ TRSRT = TS^{-1}T = (TST)^{-1}, \end{array}$$

the generators are S and TST . Thus $\overline{H}'(\lambda_q)$ has a presentation

$$\overline{H}'(\lambda_q) = \langle S, TST \mid S^q = (TST)^q = I \rangle \cong C_q * C_q.$$

□

Theorem 2. *Let q be even, then*

- (i) $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong C_2 \times C_2 \times C_2$
- (ii) $\overline{H}'(\lambda_q) = \langle S^2, TS^2T, TSTS^{q-1} \mid (S^2)^{q/2} = (TS^2T)^{q/2} = (TSTS^{q-1})^\infty = I \rangle$.

Proof. (i) If the representations of $\overline{H}(\lambda_q)$ and $\overline{H}(\lambda_q)/\overline{H}'(\lambda_q)$ are considered, we obtain $S^2 = I$ as $RS = S^{-1}R$ and $RS = SR$, $S^{q-2} = S^q = S^2 = I$ as q is odd. Therefore

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) = \langle T, S, R \mid T^2 = S^2 = R^2 = (RT)^2 = (RS)^2 = (TS)^2 = I \rangle$$

and so

$$\overline{H}(\lambda_q)/\overline{H}'(\lambda_q) \cong C_2 \times C_2 \times C_2.$$

(ii) Again we choose a Schreier transversal for $\overline{H}'(\lambda_q)$ as

$$I, T, R, S, TR, SR, TS, TSR.$$

Hence, all possible products are

$$\begin{array}{ll} I \cdot T \cdot (T)^{-1} = I, & TR \cdot T \cdot (R)^{-1} = TRTR, \\ T \cdot T \cdot (I)^{-1} = I, & SR \cdot T \cdot (TSR)^{-1} = SRTRS^{-1}T, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & TS \cdot T \cdot (S)^{-1} = TSTS^{-1}, \\ S \cdot T \cdot (TS)^{-1} = STS^{-1}T, & TSR \cdot T \cdot (SR)^{-1} = TSRTRS^{-1}, \\ I \cdot S \cdot (S)^{-1} = I, & TR \cdot S \cdot (TSR)^{-1} = TRSRS^{-1}T, \\ T \cdot S \cdot (TS)^{-1} = I, & SR \cdot S \cdot (R)^{-1} = SRSR, \\ R \cdot S \cdot (SR)^{-1} = RSRS^{-1}, & TS \cdot S \cdot (T)^{-1} = TS^2T, \\ S \cdot S \cdot (I)^{-1} = S^2, & TSR \cdot S \cdot (TR)^{-1} = TSRSRT, \\ I \cdot R \cdot (R)^{-1} = I, & TR \cdot R \cdot (T)^{-1} = I, \\ T \cdot R \cdot (TR)^{-1} = I, & SR \cdot R \cdot (S)^{-1} = I, \\ R \cdot R \cdot (I)^{-1} = I, & TS \cdot R \cdot (TSR)^{-1} = I, \\ S \cdot R \cdot (SR)^{-1} = I, & TSR \cdot R \cdot (TS)^{-1} = I. \end{array}$$

Since $(STS^{-1}T)^{-1} = TSTS^{-1}$, $(TRTR)^{-1} = RTRT = I$, $(RSRS^{-1}) = (S^2)^{-1}$, $SRSR = I$, $(SRTRS^{-1}T)^{-1} = TSRTRS^{-1} = TSTS^{-1}$, $TRSRS^{-1}T = (TS^2T)^{-1}$, $TSRSRT = I$, the generators of $\overline{H}'(\lambda_q)$ are S^2 , TS^2T , $TSTS^{q-1}$. Thus $\overline{H}'(\lambda_q)$ has a presentation

$$\overline{H}'(\lambda_q) = \langle S^2 \rangle * \langle TS^2T \rangle * \langle TSTS^{q-1} \rangle.$$

Example 1. Let $q = 3$. Then $\overline{H}(\lambda_3)$ is the extended modular group. In this case

$$\overline{H}(\lambda_3)/\overline{H}'(\lambda_3) = \langle T, R \mid T^2 = R^2 = (TR)^2 = I \rangle$$

and a Schreier transversal is

$$I, T, R, TR.$$

Hence,

$$\begin{array}{lll} I \cdot T \cdot (T)^{-1} = I, & I \cdot S \cdot (I)^{-1} = S, & I \cdot R \cdot (R)^{-1} = I, \\ T \cdot T \cdot (I)^{-1} = I, & T \cdot S \cdot (T)^{-1} = TST, & T \cdot R \cdot (TR)^{-1} = I, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & R \cdot S \cdot (R)^{-1} = RSR, & R \cdot R \cdot (I)^{-1} = I, \\ TR \cdot T \cdot (R)^{-1} = TRTR, & TR \cdot S \cdot (TR)^{-1} = TRSRT, & TR \cdot R \cdot (T)^{-1} = I \end{array}$$

and since $RTRT = I$, $TRTR = I$, $RSR = S^{-1}$, $TRSRT = TS^{-1}T = (TST)^{-1}$, the generators of $\overline{H}(\lambda_3)$ are S and TST . Thus $\overline{H}'(\lambda_3)$ has a presentation

$$\overline{H}'(\lambda_3) = \langle S, TST \mid S^3 = (TST)^3 = I \rangle \cong C_3 * C_3.$$

Notice that this result coincides with the ones given in [5] for the extended modular group.

Example 2. Let $q = 6$. Then $\overline{H}(\lambda_6)$ and $\overline{H}(\lambda_6)/\overline{H}'(\lambda_6)$ have presentations

$$\overline{H}(\lambda_6) = \langle T, S, R \mid T^2 = S^6 = R^2 = I, TR = RT, RS = S^{-1}R \rangle$$

and

$$\begin{aligned} \overline{H}(\lambda_6)/\overline{H}'(\lambda_6) = \langle T, S, R \mid T^2 = S^6 = R^2 = I, RT = TR, RS = S^{-1}R, \\ RS = SR, TS = ST \rangle. \end{aligned}$$

Since $RS = S^{-1}R$ and $RS = SR$, $S^{-1} = S^5$ and so $S^4 = S^6 = I$, $S^2 = I$. Hence

$$\overline{H}(\lambda_6)/\overline{H}'(\lambda_6) = \langle T, S, R \mid T^2 = S^2 = R^2 = (RT)^2 = (RS)^2 = (TS)^2 = I \rangle.$$

We can choose a Schreier transversal as

$$I, T, R, S, TR, SR, TS, TSR.$$

In this case all the possibilities are

$$\begin{array}{ll} I \cdot T \cdot (T)^{-1} = I, & TR \cdot T \cdot (R)^{-1} = TRTR, \\ T \cdot T \cdot (I)^{-1} = I, & SR \cdot T \cdot (TSR)^{-1} = SRTRS^5T, \\ R \cdot T \cdot (TR)^{-1} = RTRT, & TS \cdot T \cdot (S)^{-1} = TSTS^5, \\ S \cdot T \cdot (TS)^{-1} = STS^5T, & TSR \cdot T \cdot (SR)^{-1} = TSRTRS^5, \\ I \cdot S \cdot (S)^{-1} = I, & TR \cdot S \cdot (TSR)^{-1} = TRSRS^5T, \\ T \cdot S \cdot (TS)^{-1} = I, & SR \cdot S \cdot (R)^{-1} = SRSR, \\ R \cdot S \cdot (SR)^{-1} = RSRS^5, & TS \cdot S \cdot (T)^{-1} = TS^2T, \\ S \cdot S \cdot (I)^{-1} = S^2, & TSR \cdot S \cdot (TR)^{-1} = TSRSRT, \\ I \cdot R \cdot (R)^{-1} = I, & TR \cdot R \cdot (T)^{-1} = I, \\ T \cdot R \cdot (TR)^{-1} = I, & SR \cdot R \cdot (S)^{-1} = I, \\ R \cdot R \cdot (I)^{-1} = I, & TS \cdot R \cdot (TSR)^{-1} = I, \\ S \cdot R \cdot (SR)^{-1} = I, & TSR \cdot R \cdot (TS)^{-1} = I. \end{array}$$

Since $(STS^5T)^{-1} = TSTS^5$, $(TRTR)^{-1} = RTRT = I$, $(RSRS^5) = (S^2)^{-1}$, $SRSR = I$, $(SRTRS^5T)^{-1} = TSRTRS^5 = TSTS^5$, $TRSR S^5T = (TS^2T)^{-1}$, $TSRSRT = I$, the generators of $\overline{H}'(\lambda_q)$ are S^2 , TS^2T , $TSTS^5$. Thus $\overline{H}'(\lambda_6)$ has a presentation

$$\overline{H}'(\lambda_6) = \langle S^2 \rangle * \langle TS^2T \rangle * \langle TSTS^5 \rangle.$$

References

- [1] *R. B. J. T. Allenby*: Rings, Fields and Groups. Second Edition. Edward Arnold, London-New York-Melbourne-Auckland, 1991.
- [2] *I. N. Cangül and D. Singerman*: Normal subgroups of Hecke groups and regular maps. *Math. Proc. Camb. Phil. Soc.* *123* (1998), 59–74.
- [3] *H. S. M. Coxeter and W. O. J. Moser*: Generators and Relations for Discrete Groups. Springer, Berlin, 1957.
- [4] *E. Hecke*: Über die Bestimmung Dirichletscher Reihen durch ihre Funktionalgleichungen. *Math. Ann.* *112* (1936), 664–699.
- [5] *D. L. Johnson*: Topics in the Theory of Group Presentations. L.M.S. Lecture Note Series 42. Cambridge Univ. Press, Cambridge, 1980.
- [6] *G. A. Jones and J. S. Thornton*: Automorphisms and congruence subgroups of the extended modular group. *J. London Math. Soc.* *34* (1986), 26–40.

Authors' addresses: R. Sahin, Balikesir Universitesi Fen-Edebiyat Fakültesi Matematik Bölümü 10100, Balikesir, Turkey, e-mail: rsahin@balikesir.edu.tr; O. Bizim, I. N. Cangül, Uludag Universitesi Fen-Edebiyat Fakültesi Matematik Bölümü Görükle 16059, Bursa, Turkey, e-mails: obizim@uludag.edu.tr, cangul@uludag.edu.tr.