

Jaroslav Ježek; M. Maróti; R. McKenzie
Quasiequational theories of flat algebras

Czechoslovak Mathematical Journal, Vol. 55 (2005), No. 3, 665–675

Persistent URL: <http://dml.cz/dmlcz/128010>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

QUASIEQUATIONAL THEORIES OF FLAT ALGEBRAS

J. JEŽEK, Praha, M. MARÓTI and R. MCKENZIE, Nashville

(Received October 16, 2002)

Abstract. We prove that finite flat digraph algebras and, more generally, finite compatible flat algebras satisfying a certain condition are finitely q -based (possess a finite basis for their quasiequations). We also exhibit an example of a twelve-element compatible flat algebra that is not finitely q -based.

Keywords: quasiequation, flat algebra

MSC 2000: 08C15, 08B05

1. INTRODUCTION

For a finite directed graph (V, E) one can define an algebra with the underlying set $V \cup E \cup \{0\}$, one constant 0 and two binary operations \wedge, \cdot in this way: $a \wedge a = a$ and $a \wedge b = 0$ whenever $a \neq b$; $ab = c$ whenever $a, c \in V$ and $b = (a, c) \in E$; $ab = 0$ in all other cases. Algebras obtained from finite directed graphs in this way are called finite *flat digraph algebras*. One particular six-element flat digraph algebra (inherently non-finitely based for equations) played a significant role in the proof of undecidability of the existence of a finite basis for the equational theory of a finite algebra ([2], [3] and [4]). It was plausible to expect that it could serve a similar purpose in an attempt to prove that also the existence of a finite basis for the quasiequations of a finite algebra is undecidable. However, in this paper we are going to show that all finite flat digraph algebras are finitely q -based (possess a finite basis for their uasiequations), which makes them unsuitable. We will investigate a more

The first author was partially supported by the grant # 201/02/0594 of the Grant Agency of the Czech Republic, and by the Institutional grant MSM0021620839; the second author was partially supported by the grant No. T\37877 of the Hungarian National Foundation for Scientific Research (OTHA); the third author was supported by the NSF grant # DMS-9971352.

general class of finite compatible flat algebras, in which (under a modest assumption on the signature) every algebra can be embedded both into a finitely q -based and into a non-finitely q -based algebra.

For the terminology and basic concepts of universal algebra the reader is referred to the monograph [5]. For the literature on quasiequational theories see, e.g., [1] and [6].

2. COMPATIBLE 0-SEMILATTICE ALGEBRAS

Let σ be a finite signature containing (among other symbols) a binary symbol \wedge (the meet) and a nullary symbol 0 .

By a *0-semilattice algebra* we mean a σ -algebra satisfying the equations

- (1) $x \wedge (y \wedge z) = (x \wedge y) \wedge z$,
- (2) $x \wedge y = y \wedge x$,
- (3) $x \wedge x = x$,
- (4) $f(x_1, \dots, x_{i-1}, 0, x_{i+1}, \dots, x_n) = 0$ for every n -ary operation f of σ and every $i \in \{1, \dots, n\}$.

A 0-semilattice algebra is said to be *compatible* if it satisfies the equations

- (5) $f(z_1, \dots, z_{i-1}, x \wedge y, z_{i+1}, \dots, z_n) = f(z_1, \dots, z_{i-1}, x, z_{i+1}, \dots, z_n) \wedge f(z_1, \dots, z_{i-1}, y, z_{i+1}, \dots, z_n)$ for every n -ary operation f of σ and every $i \in \{1, \dots, n\}$.

So, the class of compatible 0-semilattice σ -algebras is a variety.

For a variable x , *basic x -terms* of depth n are defined as follows. The term x is the only basic x -term of depth 0. For $n > 0$, basic x -terms of depth n are the terms $f(x_1, \dots, x_{i-1}, t, x_{i+1}, \dots, x_n)$ such that f is an n -ary operation of σ , $1 \leq i \leq n$, t is a basic x -term of depth $n - 1$ and x_1, \dots are variables different from x . A basic x -term t will be usually denoted by $t(x)$, in which case $t(u)$ stands for the term resulting from t by substituting u for x (where u is any term).

For a σ -algebra B and a basic x -term t of depth n , any interpretation of the variables different from x by elements of B gives rise to a unary polynomial of B . The unary polynomials obtained in this way will be called the *basic polynomials* of B of depth n .

Lemma 2.1. *Let A be a compatible 0-semilattice algebra. Then $p(a \wedge b) = p(a) \wedge p(b)$ for all basic polynomials p of A and all elements $a, b \in A$.*

Proof. It is easy. (Observe that the statement is not true for all unary polynomials p .) □

Lemma 2.2. *Let A be a compatible 0-semilattice algebra and F be a proper filter of A (i.e., a nonempty subset closed under meet, not containing 0 and such that $b \in F$ whenever $a \in F$ and $a \leq b$). Then for every basic polynomial p of A , $p^{-1}(F)$ is either empty or a proper filter of A .*

Proof. It follows easily from Lemma 2.1. □

By a *flat* algebra we mean a 0-semilattice algebra A such that $a \wedge b = 0$ for all pairs of distinct elements $a, b \in A$. Observe that a flat algebra is monotonic, i.e., satisfies $x \leq y \rightarrow f(z_1, \dots, z_{i-1}, x, z_{i+1}, \dots, z_n) \leq f(z_1, \dots, z_{i-1}, y, z_{i+1}, \dots, z_n)$ for every n -ary operation f of σ and every $i \in \{1, \dots, n\}$.

One can easily see that a flat algebra is compatible if and only if

$$(5') \quad f(c_1, \dots, c_{i-1}, a, c_{i+1}, \dots, c_n) = f(c_1, \dots, c_{i-1}, b, c_{i+1}, \dots, c_n) \neq 0 \text{ implies } a = b$$

for every n -ary operation f of σ and every $i \in \{1, \dots, n\}$.

For every partial algebra G of a signature τ not containing \wedge and 0 we can define a flat $\tau \cup \{\wedge, 0\}$ -algebra A , called the *flat algebra over G* , by $A = G \cup \{0\}$, $f(a_1, \dots, a_n) = a$ in A whenever $f(a_1, \dots, a_n) = a$ in G , and $f(a_1, \dots, a_n) = 0$ otherwise. This flat algebra is not necessarily compatible. For example, if G is a finite groupoid, then the flat algebra over G is compatible if and only if G is a quasigroup. Finite flat digraph algebras are all compatible.

Observation 2.3. *For every finite compatible flat algebra A there exists a first-order sentence Φ such that the finite models of Φ are precisely the finite algebras belonging to the quasivariety generated by A .*

Proof. Put $K = |A|$. It is easy to see that the following are equivalent for a finite compatible 0-semilattice algebra B :

- (e1) B belongs to the quasivariety generated by A ;
- (e2) every two elements b_0, b_1 of B such that $b_0 < b_1$ can be separated by a congruence of B , the factor by which is isomorphic to a subalgebra of A ;
- (e3) for every $b_0, b_1 \in B$ with $b_0 < b_1$ there exist elements $c_1, \dots, c_r \in B$ for some $r < K$ such that the principal filters F_1, \dots, F_r generated by c_1, \dots, c_r are pairwise disjoint, $b_1 \in F_1$, b_0 belongs to the complement O of $F_1 \cup \dots \cup F_r$ in B , the equivalence R with blocks O, F_1, \dots, F_r is a congruence of B and the factor B/R is isomorphic to a subalgebra of A .

Clearly, the condition (e3) can be rewritten as a first-order sentence. □

3. THE QUASIVARIETY Q'_A

In the following let A be a finite compatible, flat algebra. Put $K = |A|$.

Denote by Q'_A the quasivariety determined by the equations (1)–(5) and the following quasiequations:

- (6) $x_0 \leq x_1 \ \& \ t(x) \geq x_1 \ \& \ u(x) \geq x_1 \ \& \ t(y) \geq x_1 \ \& \ u(y) \wedge x_1 \leq x_0 \rightarrow x_0 = x_1$ for every pair of basic x -terms t, u of depth $\leq K$;
- (7) $x_0 \leq x_1 \ \& \ H_{t_1, \dots, t_K} \rightarrow x_0 = x_1$ for every K -tuple of basic x -terms t_1, \dots, t_K of depth $\leq K$, where H_{t_1, \dots, t_K} is the conjunction of the following equations:

$$t_i(x_i) \geq x_1 \quad (i = 1, \dots, K),$$

$$t_i(x_j) \wedge x_1 \leq x_0 \quad (i, j = 1, \dots, K \text{ and } i \neq j).$$

Lemma 3.1. Q'_A is a finitely q -based quasivariety containing A .

Proof. The set of quasiequations (6)–(7) is essentially finite, as it contains only finitely many quasiequations that differ by not only renaming their variables. Consequently, Q'_A is finitely q -based. It remains to prove that (6) and (7) are satisfied in A . Suppose that (6) fails in A by some interpretation $v \mapsto v'$ of variables. Then $x'_0 < x'_1$, so that $x'_0 = 0$; now $t(x') \geq x'_1$ implies $t(x') = x'_1$. Similarly we get $u(x') = x'_1$ and $t(y') = x'_1$. But A satisfies (5'), so $t(x') = t(y') \neq 0$ implies $x' = y'$; hence $x'_1 = u(x') \wedge x'_1 = u(y') \wedge x'_1 = 0$, a contradiction. Using the fact that A cannot contain K nonzero, pairwise distinct elements, one can similarly prove that A satisfies the quasiequations (7). □

Lemma 3.2. Let $B \in Q'(A)$ and $b_0, b_1 \in B$ be two elements such that $b_1 \not\leq b_0$; let F be a maximal filter of B such that $b_1 \in F$ and $b_0 \notin F$. For any two basic polynomials p, q of B of depth $\leq K$, the sets $p^{-1}(F)$ and $q^{-1}(F)$ are either disjoint or equal.

Proof. The two basic polynomials p and q correspond to two basic terms t and u of depth $\leq K$. Suppose that there exist elements x', y' such that $p(x') \in F$, $p(y') \in F$, $q(x') \in F$ and $q(y') \notin F$. It follows from the maximality of F that there exists an element $e \in F$ with $q(y') \wedge e \leq b_0$. Put $x'_1 = p(x') \wedge p(y') \wedge q(x') \wedge e$, so that $x'_1 \in F$. Put $x'_0 = b_0 \wedge x'_1$, so that $x'_0 < x'_1$. But the quasiequation (e6) interpreted by $x \mapsto x', y \mapsto y', x_0 \mapsto x'_0, x_1 \mapsto x'_1$ gives $x'_0 = x'_1$, a contradiction. □

Lemma 3.3. Let $B \in Q'(A)$ and $b_0, b_1 \in B$ be two elements such that $b_1 \notin b_0$; let F be a maximal filter of B such that $b_1 \in F$ and $b_0 \notin F$. There are at most $K - 1$ nonempty subsets of B that can be expressed as $q^{-1}(F)$ for a basic polynomial q of B , and they can be arranged into a sequence F_1, \dots, F_r (for some $r < K$) in such a way that $F_1 = F$ and for every $i \in \{2, \dots, r\}$ there are an index $j \in \{1, \dots, i-1\}$ and a basic polynomial p_i of B of depth 1 with $F_i = p_i^{-1}(F_j)$. The collection F_1, \dots, F_r , together with the complement of their union, is a partition and the corresponding equivalence is a congruence of B .

Proof. Let us define a (finite or infinite) sequence $F_1, p_1, F_2, p_2, \dots$ of filters F_i and basic polynomials p_i of depth ≤ 1 by induction in this way: $F_1 = F$ and p_1 is the identity on B ; if F_i, p_i have been defined and if there exist an element $a \notin F_1 \cup \dots \cup F_i$ and a basic polynomial p of depth 1 such that $p(a) \in F_j$ for some $j \leq i$, take one such pair a, p and put $p_{i+1} = p$ and $F_{i+1} = p_{i+1}^{-1}(F_j)$; if there is no such pair a, p , the sequence already constructed will have no continuation. Clearly (by induction on i), $F_i = q_i^{-1}(F)$ for a basic polynomial q_i of B of depth $< i$. The sets F_i are pairwise disjoint filters according to Lemmas 2.2 and 3.2.

Suppose that the sequence has at least K members F_1, \dots, F_K . For any $i = 1, \dots, K$ take an element $x'_i \in F_i$, so that $q_i(x'_i) \in F$. For every $i \neq j$ we have $x'_j \notin F_i$, i.e., $q_i(x'_j) \notin F$, so that there exists an element $e_{i,j} \in F$ with $q_i(x'_j) \wedge e_{i,j} \leq b_0$. There is an element $x'_1 \in F$ such that $x'_1 \leq q_i(x'_i)$ for all i and $x'_1 \leq e_{i,j}$ for all $i \neq j$. Put $x'_0 = b_0 \wedge x'_1$, so that $x'_0 < x'_1$. But the quasiequation (e7), interpreted in the obvious way, says that $x'_0 = x'_1$, a contradiction.

So, the sequence F_1, p_1, \dots ends with F_r, p_r for some $r \leq K - 1$. Clearly, every subset of the form $q^{-1}(F)$ for a basic polynomial q can be found among F_1, \dots, F_r . Put $O = B - (F_1 \cup \dots \cup F_r)$, so that $0 \in O$ and F_1, \dots, F_r, O is a partition of B . It remains to prove that the corresponding equivalence is a congruence of B .

Suppose that there exist an n -ary operation f in σ and an n -tuple a_1, \dots, a_n of elements of B such that $a_j \in O$ for some j but $f(a_1, \dots, a_n) \in F_i$ for some i . Then $p(a_j) \in F_i$ where $p(x) = f(a_1, \dots, a_{j-1}, x, a_{j+1}, \dots, a_n)$ is a basic polynomial of depth 1 and $a_j \notin F_1 \cup \dots \cup F_r$, so that $(q_i p)^{-1}(F)$ is nonempty and different from all F_1, \dots, F_r , a contradiction. We have proved that if at least one of the elements a_1, \dots, a_n belongs to O , then $f(a_1, \dots, a_n) \in O$.

Now it remains to show that if f is n -ary, $f(a_1, \dots, a_n) \in F_j$ and $a_i, a'_i \in F_k$ for some $j, k \in \{1, \dots, r\}$ and $i \in \{1, \dots, n\}$, then $f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_n) \in F_j$. Put $q(x) = q_j(f(a_1, \dots, a_{i-1}, x, a_{i+1}, \dots, a_n))$, so that q is a basic polynomial of B of depth at most K . We have $q(a_i) \in F$ and $q_k(a_i) \in F$, so that $q^{-1}(F) = q_k^{-1}(F)$. Since a'_i belongs to this set, we get $q(a'_i) \in F$, i.e., $f(a_1, \dots, a_{i-1}, a'_i, a_{i+1}, \dots, a_n) \in F_j$. \square

Theorem 3.4. *Let A be a finite compatible, flat algebra with K elements. Then Q'_A is a finitely q -based and locally finite quasivariety containing A ; every algebra in Q'_A is isomorphic to a subdirect product of algebras of cardinality at most K . Consequently, A is not inherently nonfinitely q -based.*

Proof. Let $B \in Q'_A$. For every pair b_0, b_1 of distinct elements of B (we can assume that $b_1 \not\leq b_0$) there exists a maximal filter of B containing b_1 but not b_0 , so that by Lemma 3.3 these two elements can be separated by a congruence with at most K blocks. It follows that every algebra from B is isomorphic to a subdirect product of algebras of cardinality at most K . Thus Q'_A is contained in a finitely generated variety and hence it is locally finite. According to Lemma 3.1, Q'_A is finitely q -based and contains A . \square

4. FINITELY q -BASED COMPATIBLE FLAT ALGEBRAS

Let A be a finite compatible flat algebra. By a *segment* of A we will mean a nonempty subset of A , the elements of which can be arranged into a finite sequence $0, c_1, \dots, c_r$ in such a way that $c_1 \neq 0$ and for every $i = 2, \dots, r$ there exists a basic polynomial p of A of depth 1 with $p(c_i) = c_j$ for some $j \in \{1, \dots, i - 1\}$.

Let S be a segment of A . The algebra obtained from S , considered as a partial subalgebra of A , by setting all the undefined operations to 0 will be called the *0-completion* of S .

Let S be a segment of A and S' be the subalgebra of A generated by S . The segment S is said to be *regular* if the equivalence on S' with the only non-singleton block $\{0\} \cup (S' - S)$ is a congruence of S' . In that case, the factor of S' by this congruence is isomorphic to the 0-completion of S .

Theorem 4.1. *Let A be a finite compatible flat algebra such that the 0-completion of every regular segment of A belongs to the quasivariety generated by A . Then A is finitely q -based.*

Proof. Denote by Q''_A the subquasivariety of Q'_A determined by the quasi-equations (1)–(7) and all quasiequations in at most K variables that are satisfied in A . (Here $K = |A|$.) Since Q'_A is locally finite by Theorem 3.4, Q''_A is locally finite. Since only finitely many equations are needed to reduce the terms in at most K variables to a finite set T_0 of such terms, and then quasiequations in at most K variables correspond to subsets of T_0^2 with distinguished elements, Q''_A is finitely q -based. Of course, $A \in Q''_A$. We are going to prove that Q''_A is the quasivariety generated by A . It is sufficient to show that every finite algebra from Q''_A belongs to the quasivariety generated by A .

Let B be a finite algebra from Q''_A ; let $b_0, b_1 \in B$ be such that $b_1 \not\leq b_0$. By 3.3 there is a congruence with at most K blocks O, F_1, \dots, F_r , yielding a quotient algebra C , such that F_1, \dots, F_r are filters (now they are principal filters), $F_1 = F$, $b_1 \in F_1$, $b_0 \in O$, and for every $i \in \{2, \dots, r\}$ there exist an index $j < i$ and a basic polynomial p_i of length 1 with $F_i = p_i^{-1}(F_j)$. But all the coefficients occurring in p_i belong to $F_1 \cup \dots \cup F_r$, so there exists a basic x -term $u_i(x, x_1, \dots, x_r)$ of depth 1 such that $u_i(F_i, F_1, \dots, F_r) \subseteq F_j$. Now we can combine these terms u_i together to obtain, for each i , a basic x -term $t_i(x, x_1, \dots, x_r)$ such that $t_i(F_i, F_1, \dots, F_r) \subseteq F$, i.e., $t_i^C(F_i, F_1, \dots, F_r) = F_1$. (We take $t_1 = x$.) For any term u denote by $t_i(u)$ the term obtained from t_i by replacing the only occurrence of x with u . Now consider the quasiequation

$$x_0 \leq x_1 \ \& \ D \rightarrow x_0 = x_1$$

where D is the conjunction of all these equations:

- (i) $t_i(x_i) \geq x_1$, for any $i = 1, \dots, r$;
- (ii) $t_i(x_j) \wedge x_1 \leq x_0$, for any $i, j \in \{1, \dots, r\}$ with $i \neq j$;
- (iii) $t_i(f(x_{i_1}, \dots, x_{i_n})) \geq x_1$, for any n -ary operation f of σ and any i, i_1, \dots, i_n with $f^C(F_{i_1}, \dots, F_{i_n}) = F_i$;
- (iv) $t_i(u) \wedge x_1 \leq x_0$, for any $i = 1, \dots, r$ and any term u in variables x_1, \dots, x_r containing a subterm $f(x_{i_1}, \dots, x_{i_n})$ with $f^C(F_{i_1}, \dots, F_{i_n}) = O$ (it is possible to consider only finitely many such terms u).

Clearly, this quasiequation fails in B ; since it is a quasiequation in at most K variables x_0, \dots, x_r , it must fail in A by some elements a_0, a_1, \dots, a_r . But then the subset $\{a_0, a_1, \dots, a_r\}$ is a regular segment of A , and the 0-completion of this subset is isomorphic to C . Since C belongs to the quasivariety generated by A , the elements b_0, b_1 were separated by a congruence, the factor by which belongs to the quasivariety. □

Corollary 4.2. *Every finite flat digraph algebra is finitely q -based.*

Proof. In this case, all segments are subalgebras. □

Corollary 4.3. *The flat algebra over any finite quasigroup (considered as a groupoid) is finitely q -based.*

Proof. In this case, all regular segments are subalgebras. □

Corollary 4.4. *If σ is the signature containing only one unary symbol in addition to \wedge and 0, then every finite compatible flat σ -algebra is finitely q -based.*

Proof. In this case, the 0-completion of every segment is isomorphic to a subalgebra. □

5. THE EMBEDDING THEOREM

Theorem 5.1. *Let σ be a finite signature containing, in addition to \wedge and 0 , at least two unary symbols f and g (and, possibly, some other operation symbols). Then every finite compatible flat σ -algebra can be embedded into two finite compatible flat σ -algebras, one finitely q -based and the other one not finitely q -based.*

Proof. Let G be a finite compatible flat algebra.

Denote by S_1, \dots, S_r all the segments of G . (It would be sufficient to take just those with the 0 -completions not belonging to the quasivariety generated by G .) For every $i = 1, \dots, r$ let us take an isomorphic copy T_i of the partial algebra $S_i - \{0\}$, in such a way that the sets G, T_1, \dots, T_r are pairwise disjoint. Denote by G' the flat algebra with the underlying set $G \cup T_1 \cup \dots \cup T_r$, with the operations evaluated to 0 in all cases except when needed to define them in such a way that G is a subalgebra and T_i are partial subalgebras. It follows from Theorem 4.1 that G' is finitely q -based.

Next we are going to construct a non-finitely q -based extension of G . Let us take one fixed positive integer k such that $k \geq 2$ and there is no sequence u_0, u_1, \dots, u_k of pairwise distinct elements of $G - \{0\}$ such that $g(u_{i-1}) = u_i$ for $i = 1, \dots, k$. Denote by A the flat algebra, with G as a subalgebra, containing $k + 10$ additional elements $u_0, u_1, \dots, u_k, a, b, c, v_2, a_2, b_2, v_3, a_3, c_3$ with all operations not inside G evaluated to 0 except for

$$\begin{aligned}
 &g(u_{i-1}) = u_i \quad \text{for } i = 1, \dots, k, \\
 &f(u_0) = a, \quad f(a) = b, \quad g(a) = c, \\
 &f(v_2) = a_2, \quad f(a_2) = b_2, \quad f(v_3) = a_3, \quad g(a_3) = b_3.
 \end{aligned}$$

(Fig. 1, in which the elements not belonging to G are pictured for $k = 2$, may help to understand this definition.)




Fig. 1

Denote by Q the quasivariety generated by A . A σ -algebra B belongs to Q if and only if every two distinct elements of B can be separated by a homomorphism of B into A .

For every positive integer n let A_n be the σ -algebra with elements $0, u_0, \dots, u_k, \alpha_{i,j}, \beta_i, \gamma_j$ ($0 \leq i \leq n, 0 \leq j \leq n-1, i-1 \leq j \leq i$) and with operations defined in this way: A_n is a semilattice with the only comparabilities $0 < u_i$ ($i = 0, \dots, k$), $0 < \beta_n < \beta_{n-1} < \dots < \beta_0$, $0 < \gamma_{n-1} < \gamma_{n-2} < \dots < \gamma_0$, $0 < \alpha_{n,n-1} < \alpha_{n-1,n-1} < \alpha_{n-1,n-2} < \dots < \alpha_{1,0} < \alpha_{0,0}$; the other operations evaluate to 0 except that $g(u_{i-1}) = u_i$ ($i = 1, \dots, k$), $f(u_0) = \alpha_{0,0}$, $f(\alpha_{i,j}) = \beta_i$, $g(\alpha_{i,j}) = \gamma_j$. (Fig. 2, in which the situation is illustrated for $k = 2$ and $n = 3$, may help to understand this definition. In the picture lines with arrows indicate unary operations, while the other lines represent coverings but the covers between 0 and the elements u_i are not indicated.)




Fig. 2

Denote by r_n the equivalence on A_n with the only non-singleton block $\{0, \beta_n\}$. Clearly, r_n is a congruence of A_n . Denote the factor A_n/r_n by B_n . For $a \in A_n - \{0, \beta_n\}$, the element a/r_n will be identified with a .

Suppose that there exists a homomorphism $H: B_n \rightarrow A$ such that $H(u_k) \neq H(0/r_n)$, i.e., $H(u_k) \neq 0$. Since $g^k(u_0) = u_k$ in B_n and there is no other element e

in A with $g^k(e) \neq 0$ and $g^{k+1}(e) = 0$ other than u_0 , we get $H(u_0) = u_0$ and then $H(u_i) = H(g^i(u_0)) = g^i(H(u_0)) = g^i(u_0) = u_i$ for all i . Now $H(\alpha_{0,0}) = H(f(u_0)) = f(H(u_0)) = f(u_0) = a$. Consequently, $H(\beta_0) = b$ and $H(\gamma_0) = c$. Since $g(\alpha_{1,0}) = \gamma_0$ and a is the only element of A with $g(a) = c$, it follows that $H(\alpha_{1,0}) = a$. If $H(\alpha_{i,i-1}) = a$ for some $i < n$, then using f in a similar way we can show that $H(\alpha_{i,i}) = a$, and then using g to show that $H(\alpha_{i+1,i}) = a$. By induction we get $H(\alpha_{n,n-1}) = a$. But then $H(0/r_n) = H(\beta_n/r_n) = H(f(\alpha_{n,n-1})) = f(a) = b$, a contradiction.

Since the element u_k cannot be separated from $0/r_n$ by a homomorphism of B_n into A , we conclude that B_n does not belong to Q .

Let $\alpha_{m,m'}$ be an element of B_n such that $0 < m < n$. Clearly, the set $C = B_n - \{\alpha_{m,m'}\}$ is a subalgebra of B_n . We are going to prove that C belongs to Q . For this purpose, it is sufficient to show that whenever e, e' are two elements of C such that e is covered by e' , then e, e' can be separated by a homomorphism of C into A .

For every $i \leq n - 1$ define a mapping ψ_i of B_n into A by $\psi_i(u_0) = v_2, \psi_i(e) = a_2$ for $e \geq \alpha_{i,i}, \psi_i(e) = b_2$ for $e \geq \beta_i$ and $\psi_i(e) = 0$ for all other elements e . Also, for every $i \leq n - 1$ define a mapping χ_i of B_n into A by $\chi_i(u_0) = v_3, \chi_i(e) = a_3$ for $e \geq \alpha_{i+1,i}, \chi_i(e) = c_3$ for $e \geq \gamma_i$ and $\chi_i(e) = 0$ for all other elements e . It is easy to check that both ψ_i and χ_i are homomorphisms. Consequently, their restrictions to C are homomorphisms of C into A . The only pairs of covers not separated by any of these homomorphisms are the pairs $(0, u_1), \dots, (0, u_k)$. So, it remains to separate these pairs of elements.

If $m = m'$, then these pairs are separated by the homomorphism φ defined in this way: $\varphi(u_0) = u_0, \dots, \varphi(u_k) = u_k, \varphi(e) = a$ for $e \geq \alpha_{m,m-1}, \varphi(e) = b$ for $e \geq \beta_m, \varphi(e) = c$ for $e \geq \gamma_{m-1}$ and $\varphi(e) = 0$ for all other elements e . If $m' = m - 1$, then they are separated by the homomorphism φ' defined in this way: $\varphi'(u_0) = u_0, \dots, \varphi'(u_k) = u_k, \varphi'(e) = a$ for $e \geq \alpha_{m',m'}, \varphi'(e) = b$ for $e \geq \beta_{m'}, \varphi'(e) = c$ for $e \geq \gamma_{m'}$ and $\varphi'(e) = 0$ for all other elements e .

We have proved that C belongs to Q . Since every subalgebra of B_n generated by at most $n - k$ elements is contained in at least one such C , it follows that every subalgebra generated by at most $n - k$ elements belongs to Q . Consequently, there is no base for the quasiequations of Q that would contain only quasiequations in at most $n - k$ variables. Since k was fixed while n was arbitrary, there is no finite base at all. \square

Remark 5.2. In the above construction of the algebra A it was not essential that the elements b_2 and c_3 are distinct.

References

- [1] *J. Lawrence and R. Willard*: On finitely based groups and nonfinitely based quasivarieties. *J. Algebra* 203 (1998), 1–11.
- [2] *R. McKenzie*: The residual bounds of finite algebras. *Internat. J. Alg. Comput.* 6 (1996), 1–28.
- [3] *R. McKenzie*: The residual bound of a finite algebra is not computable. *Internat. J. Alg. Comput.* 6 (1996), 29–48.
- [4] *R. McKenzie*: Tarski’s finite basis problem is undecidable. *Internat. J. Alg. Comput.* 6 (1996), 49–104.
- [5] *R. McKenzie, G. McNulty and W. Taylor*: *Algebras, Lattices, Varieties, Vol. I*. Wadsworth & Brooks/Cole, Monterey, 1987.
- [6] *D. Pigozzi*: Finite basis theorems for relatively congruence-distributive quasivarieties. *Transactions of the AMS* 310 (1988), 499–533.

Authors’ addresses: J. Ježek, MFF UK, Sokolovská 83, 186 00 Praha 8, Czech Republic, e-mail: jezek@karlin.mff.cuni.cz; M. Maróti and R. McKenzie, Department of Mathematics, Vanderbilt University, 1326 Stevenson Center, Nashville, TN 37240, e-mails: miklos.maroti@vanderbilt.edu, ralph.n.mckenzie@vanderbilt.edu.