

Silvestru Sever Dragomir

Bounds for Convex Functions of Čebyšev Functional Via Sonin's Identity with Applications

Communications in Mathematics, Vol. 22 (2014), No. 2, 107–132

Persistent URL: <http://dml.cz/dmlcz/144124>

Terms of use:

© University of Ostrava, 2014

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Bounds for Convex Functions of Čebyšev Functional Via Sonin's Identity with Applications

Silvestru Sever Dragomir

Abstract. Some new bounds for the Čebyšev functional in terms of the Lebesgue norms

$$\left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p}$$

and the Δ -seminorms

$$\|f\|_p^\Delta := \left(\int_a^b \int_a^b |f(t) - f(s)|^p dt ds \right)^{\frac{1}{p}}$$

are established. Applications for mid-point and trapezoid inequalities are provided as well.

1 Introduction

For two Lebesgue integrable functions $f, g: [a, b] \rightarrow \mathbb{R}$, consider the Čebyšev functional

$$C(f, g) := \frac{1}{b-a} \int_a^b f(t)g(t) dt - \frac{1}{(b-a)^2} \int_a^b f(t) dt \int_a^b g(t) dt.$$

In 1935, Grüss [7] showed that

$$|C(f, g)| \leq \frac{1}{4}(M-m)(N-n), \tag{1}$$

provided that there exists the real numbers m, M, n, N such that

$$m \leq f(t) \leq M \quad \text{and} \quad n \leq g(t) \leq N \quad \text{for a.e. } t \in [a, b]. \tag{2}$$

2010 MSC: 26D15, 25D10

Key words: Absolutely continuous functions, Convex functions, Integral inequalities, Čebyšev functional, Jensen's inequality, Lebesgue norms, Mid-point inequalities, Trapezoid inequalities

The constant $\frac{1}{4}$ is best possible in (1) in the sense that it cannot be replaced by a smaller quantity.

Another, however less known result, even though it was obtained by Čebyšev in 1882 [5], states that

$$|C(f, g)| \leq \frac{1}{12} \|f'\|_\infty \|g'\|_\infty (b-a)^2, \quad (3)$$

provided that f', g' exist and are continuous on $[a, b]$ and $\|f'\|_\infty = \sup_{t \in [a, b]} |f'(t)|$. The constant $\frac{1}{12}$ cannot be improved in the general case.

Čebyšev inequality (3) also holds if $f, g: [a, b] \rightarrow \mathbb{R}$ are assumed to be absolutely continuous and $f', g' \in L_\infty[a, b]$ while $\|f'\|_\infty = \text{ess sup}_{t \in [a, b]} |f'(t)|$.

A mixture between Grüss' result (1) and Čebyšev's one (3) is the following inequality obtained by Ostrowski in 1970 [12]:

$$|C(f, g)| \leq \frac{1}{8} (b-a)(M-m) \|g'\|_\infty, \quad (4)$$

provided that f is *Lebesgue integrable* and satisfies (2) while g is absolutely continuous and $g' \in L_\infty[a, b]$. The constant $\frac{1}{8}$ is best possible in (4).

The case of *euclidean norms* of the derivative was considered by A. Lupaş in [9] in which he proved that

$$|C(f, g)| \leq \frac{1}{\pi^2} \|f'\|_2 \|g'\|_2 (b-a), \quad (5)$$

provided that f, g are absolutely continuous and $f', g' \in L_2[a, b]$. The constant $\frac{1}{\pi^2}$ is the best possible.

Recently, Cerone and Dragomir [2] have proved the following results:

$$|C(f, g)| \leq \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_q \cdot \frac{1}{b-a} \left(\int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right|^p dt \right)^{\frac{1}{p}}, \quad (6)$$

where $p > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$ or $p = 1$ and $q = \infty$, and

$$|C(f, g)| \leq \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_1 \cdot \frac{1}{b-a} \text{ess sup}_{t \in [a, b]} \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right|,$$

provided that $f \in L_p[a, b]$ and $g \in L_q[a, b]$ ($p > 1, \frac{1}{p} + \frac{1}{q} = 1; p = 1, q = \infty$ or $p = \infty, q = 1$).

Notice that for $q = \infty, p = 1$ in (6) we obtain

$$\begin{aligned} |C(f, g)| &\leq \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_\infty \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \\ &\leq \|g\|_\infty \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \end{aligned}$$

and if g satisfies (2), then

$$\begin{aligned}
 |C(f, g)| &\leq \inf_{\gamma \in \mathbb{R}} \|g - \gamma\|_{\infty} \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \\
 &\leq \left\| g - \frac{n+N}{2} \right\|_{\infty} \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt \\
 &\leq \frac{1}{2}(N-n) \cdot \frac{1}{b-a} \int_a^b \left| f(t) - \frac{1}{b-a} \int_a^b f(s) ds \right| dt. \tag{7}
 \end{aligned}$$

The inequality between the first and the last term in (7) has been obtained by Cheng and Sun in [6]. However, the sharpness of the constant $\frac{1}{2}$, a generalization for the abstract Lebesgue integral and the discrete version of it have been obtained in [3].

For other recent results on the Grüss inequality, see [8], [10] and [13] and the references therein.

In this paper, some new bounds for the Čebyšev functional in terms of the Lebesgue norms $\|f - \frac{1}{b-a} \int_a^b f(t) dt\|_{[a,b],p}$ and the Δ -seminorms are established. Applications for mid-point and trapezoid inequalities are provided as well.

2 Some Results Via Sonin’s Identity

The following result for convex functions of Čebyšev functional holds.

Theorem 1. *Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue integrable functions on $[a, b]$. If $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ is convex on \mathbb{R} then we have the inequality*

$$\begin{aligned}
 \Phi[C(f, g)] &\leq \frac{1}{b-a} \inf_{\lambda \in \mathbb{R}} \int_a^b \Phi \left[\left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) (g(x) - \lambda) \right] dx \\
 &\leq \frac{1}{(b-a)^2} \inf_{\lambda \in \mathbb{R}} \int_a^b \int_a^b \Phi \left[(f(x) - f(t))(g(x) - \lambda) \right] dt dx. \tag{8}
 \end{aligned}$$

Proof. Start with Sonin’s identity [11, p. 246]

$$C(f, g) = \frac{1}{b-a} \int_a^b \left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) (g(x) - \lambda) dx$$

that holds for any $\lambda \in \mathbb{R}$.

If we use Jensen’s inequality we have for any $\lambda \in \mathbb{R}$

$$\begin{aligned}
 \Phi[C(f, g)] &= \Phi \left[\frac{1}{b-a} \int_a^b \left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) (g(x) - \lambda) dx \right] \\
 &\leq \frac{1}{b-a} \int_a^b \Phi \left[\left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) (g(x) - \lambda) \right] dx \\
 &= \frac{1}{b-a} \int_a^b \Phi \left[\frac{1}{b-a} \int_a^b \left[(f(x) - f(t))(g(x) - \lambda) \right] dt \right] dx \\
 &\leq \frac{1}{(b-a)^2} \int_a^b \int_a^b \Phi \left[(f(x) - f(t))(g(x) - \lambda) \right] dt dx.
 \end{aligned}$$

Taking the infimum over $\lambda \in \mathbb{R}$ we deduce the desired inequalities (8). □

Remark 1. If we write inequality (8) for the convex function $\Phi(x) = |x|^p$, $p \geq 1$, then we get the inequality

$$\begin{aligned} |C(f, g)| &\leq \frac{1}{(b-a)^{1/p}} \inf_{\lambda \in \mathbb{R}} \left\{ \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p |g(x) - \lambda|^p dx \right\}^{1/p} \\ &\leq \frac{1}{(b-a)^{2/p}} \inf_{\lambda \in \mathbb{R}} \left\{ \int_a^b \int_a^b |f(x) - f(t)|^p |g(x) - \lambda|^p dt dx \right\}^{1/p}. \end{aligned} \quad (9)$$

Utilising Hölder's integral inequality we have

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$\begin{aligned} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p |g(x) - \lambda|^p dx \\ \leq \operatorname{ess\,sup}_{x \in [a, b]} \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p \int_a^b |g(x) - \lambda|^p dx \\ = \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a, b], \infty}^p \|g - \lambda\|_{[a, b], p}^p, \end{aligned}$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\begin{aligned} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p |g(x) - \lambda|^p dx \\ \leq \left(\int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^{p\beta} dx \right)^{1/\beta} \left(\int_a^b |g(x) - \lambda|^{p\alpha} dx \right)^{1/\alpha} \\ = \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a, b], p\beta}^p \|g - \lambda\|_{[a, b], p\alpha}^p, \end{aligned}$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$\begin{aligned} \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p |g(x) - \lambda|^p dx \\ \leq \operatorname{ess\,sup}_{x \in [a, b]} |g(x) - \lambda|^p \int_a^b \left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right|^p dx \\ = \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a, b], p}^p \|g - \lambda\|_{[a, b], \infty}^p. \end{aligned}$$

Utilising (9) we can state the following result.

Theorem 2. Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue measurable functions on $[a, b]$. Then

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a, b], p} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a, b], \infty},$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a,b], p\alpha} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], p\beta},$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a,b], \infty} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], p}.$$

We have the following particular cases of interest.

Corollary 1. Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue measurable functions on $[a, b]$. Then

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], p} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], \infty},$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], p\alpha} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], p\beta},$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], \infty} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], p}.$$

If one function is bounded, then we can state the following result.

Corollary 2. Assume that $f, g: [a, b] \rightarrow \mathbb{R}$ are Lebesgue measurable functions on $[a, b]$. If there exist constants n, N such that $n \leq g(t) \leq N$ for a.e. $t \in [a, b]$, then

a) for $f \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{2}(N-n) \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], \infty},$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2}(N-n) \frac{1}{(b-a)^{1/p\beta}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b], p\beta},$$

c) for $f \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2}(N - n) \frac{1}{(b - a)^{1/p}} \left\| f - \frac{1}{b - a} \int_a^b f(t) dt \right\|_{[a, b], p}.$$

Proof. We observe that

$$\begin{aligned} \left\| g - \frac{n + N}{2} \right\|_{[a, b], p} &= \left(\int_a^b \left| g(t) - \frac{n + N}{2} \right|^p dt \right)^{1/p} \\ &\leq \left(\int_a^b \left(\frac{N - n}{2} \right)^p dt \right)^{1/p} = \frac{N - n}{2} (b - a)^{1/p}, \\ \left\| g - \frac{n + N}{2} \right\|_{[a, b], p\alpha} &= \left(\int_a^b \left| g(t) - \frac{n + N}{2} \right|^{p\alpha} dt \right)^{1/p\alpha} \\ &\leq \frac{N - n}{2} (b - a)^{1/p\alpha} \end{aligned}$$

and

$$\left\| g - \frac{n + N}{2} \right\|_{[a, b], \infty} \leq \frac{N - n}{2}.$$

Utilising Theorem 2 we deduce the desired result of Corollary 2. \square

When one function is of bounded variation, then we can state the following result.

Corollary 3. *If $f: [a, b] \rightarrow \mathbb{R}$ is Lebesgue integrable and $g: [a, b] \rightarrow \mathbb{R}$ is of bounded variation, then*

a) for $f \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \left\| f - \frac{1}{b - a} \int_a^b f(t) dt \right\|_{[a, b], \infty},$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \frac{1}{(b - a)^{1/p\beta}} \left\| f - \frac{1}{b - a} \int_a^b f(t) dt \right\|_{[a, b], p\beta},$$

c) for $f \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \frac{1}{(b - a)^{1/p}} \left\| f - \frac{1}{b - a} \int_a^b f(t) dt \right\|_{[a, b], p},$$

where $\bigvee_a^b(g)$ is the total variation of the function g on the interval $[a, b]$.

Proof. Since $g: [a, b] \rightarrow \mathbb{R}$ is of bounded variation, then for any $t \in [a, b]$ we have

$$\begin{aligned} \left| g(t) - \frac{g(a) + g(b)}{2} \right| &= \left| \frac{g(t) - g(a) + g(t) - g(b)}{2} \right| \\ &\leq \frac{1}{2} [|g(t) - g(a)| + |g(b) - g(t)|] \leq \frac{1}{2} \bigvee_a^b(g). \end{aligned}$$

Then

$$\begin{aligned} \left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b],p} &= \left(\int_a^b \left| g(t) - \frac{g(a) + g(b)}{2} \right|^p dt \right)^{1/p} \\ &\leq \left(\int_a^b \left(\frac{1}{2} \bigvee_a^b(g) \right)^p dt \right)^{1/p} = \frac{1}{2} \bigvee_a^b(g) (b-a)^{1/p}, \\ \left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b],p\alpha} &\leq \frac{1}{2} \bigvee_a^b(g) (b-a)^{1/p\alpha}, \end{aligned}$$

and

$$\left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b],\infty} \leq \frac{1}{2} \bigvee_a^b(g).$$

Utilising Theorem 2 we deduce the desired result of Corollary 3. \square

For functions h that are *Lipschitzian in the middle point* with the constant $L_{\frac{a+b}{2}}$ and the exponent $q > 0$, i.e. satisfying the condition

$$\left| h(t) - h\left(\frac{a+b}{2}\right) \right| \leq L_{\frac{a+b}{2}} \left| t - \frac{a+b}{2} \right|^q$$

for any $t \in [a, b]$, we have the following result as well.

Corollary 4. *If $f: [a, b] \rightarrow \mathbb{R}$ is Lebesgue integrable and $g: [a, b] \rightarrow \mathbb{R}$ is Lipschitzian in the middle point with the constant $L_{\frac{a+b}{2}}$ and the exponent $q > 0$, then*

a) for $f \in L_\infty[a, b]$

$$|C(f, g)| \leq L_{\frac{a+b}{2}} \frac{(b-a)^q}{2^q (qp+1)^{1/p}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],\infty}, \quad (10)$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q-1/p\beta}}{2^q (qp\alpha+1)^{1/p\alpha}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p\beta}, \quad (11)$$

c) for $f \in L_p[a, b]$

$$|C(f, g)| \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q-1/p}}{2^q} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p}. \quad (12)$$

Proof. We have

$$\begin{aligned} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p} &= \left(\int_a^b \left| g(t) - g\left(\frac{a+b}{2}\right) \right|^p dt \right)^{1/p} \\ &\leq \left(\int_a^b L_{\frac{a+b}{2}}^p \left| t - \frac{a+b}{2} \right|^{qp} dt \right)^{1/p} \\ &= L_{\frac{a+b}{2}} \left(\int_a^b \left| t - \frac{a+b}{2} \right|^{qp} dt \right)^{1/p}. \end{aligned} \quad (13)$$

Observe that

$$\begin{aligned} &\left(\int_a^b \left| t - \frac{a+b}{2} \right|^{qp} dt \right)^{1/p} \\ &= \left(\int_a^{\frac{a+b}{2}} \left(\frac{a+b}{2} - t \right)^{qp} dt + \int_{\frac{a+b}{2}}^b \left(t - \frac{a+b}{2} \right)^{qp} dt \right)^{1/p} \\ &= \left(2 \int_{\frac{a+b}{2}}^b \left(t - \frac{a+b}{2} \right)^{qp} dt \right)^{1/p} = \left(2 \left(t - \frac{a+b}{2} \right)^{qp+1} \Big|_{\frac{a+b}{2}}^b \right)^{1/p} \\ &= \left(2 \frac{\left(\frac{b-a}{2} \right)^{qp+1}}{qp+1} \right)^{1/p} = \left(\frac{(b-a)^{qp+1}}{2^{qp}(qp+1)} \right)^{1/p} = \frac{(b-a)^{q+1/p}}{2^q(qp+1)^{1/p}}. \end{aligned}$$

Then by (13) we have

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p} \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p}}{2^q(qp+1)^{1/p}}.$$

Also

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p\alpha} \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p\alpha}}{2^q(qp\alpha+1)^{1/p\alpha}}$$

and

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],\infty} \leq L_{\frac{a+b}{2}} \frac{(b-a)^q}{2^q}.$$

Utilising Theorem 2 we obtain

a) for $f \in L_\infty[a, b]$

$$\begin{aligned} |C(f, g)| &\leq \frac{1}{(b-a)^{1/p}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],\infty} \\ &\leq \frac{1}{(b-a)^{1/p}} L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p}}{2^q (qp+1)^{1/p}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],\infty} \\ &= L_{\frac{a+b}{2}} \frac{(b-a)^q}{2^q (qp+1)^{1/p}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],\infty}, \end{aligned}$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\begin{aligned} |C(f, g)| &\leq \frac{1}{(b-a)^{1/p}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p\alpha} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p\beta} \\ &\leq \frac{1}{(b-a)^{1/p}} L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p\alpha}}{2^q (qp\alpha+1)^{1/p\alpha}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p\beta} \\ &= L_{\frac{a+b}{2}} \frac{(b-a)^{q-1/p\beta}}{2^q (qp\alpha+1)^{1/p\alpha}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p\beta}, \end{aligned}$$

c) and for $f \in L_p[a, b]$

$$\begin{aligned} |C(f, g)| &\leq \frac{1}{(b-a)^{1/p}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],\infty} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p} \\ &\leq \frac{1}{(b-a)^{1/p}} L_{\frac{a+b}{2}} \frac{(b-a)^q}{2^q} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p} \\ &= L_{\frac{a+b}{2}} \frac{(b-a)^{q-1/p}}{2^q} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p}. \end{aligned}$$

Thus the inequalities (10)–(12) are proved. □

Remark 2. If the function g is Lipschitzian with the constant $L > 0$, then

a) for $f \in L_\infty[a, b]$

$$|C(f, g)| \leq L \frac{b-a}{2(p+1)^{1/p}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],\infty}, \tag{14}$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq L \frac{(b-a)^{1-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p\beta}, \tag{15}$$

c) for $f \in L_p[a, b]$

$$|C(f, g)| \leq L \frac{(b-a)^{1-1/p}}{2} \left\| f - \frac{1}{b-a} \int_a^b f(t) dt \right\|_{[a,b],p}. \quad (16)$$

3 Δ -Seminorms and Related Inequalities

For $f \in L_p[a, b]$, $p \in [1, \infty)$, we can define the functional (see [1] and [4])

$$\|f\|_p^\Delta := \left(\int_a^b \int_a^b |f(t) - f(s)|^p dt ds \right)^{\frac{1}{p}}$$

and for $f \in L_\infty[a, b]$, we can define

$$\|f\|_\infty^\Delta := \operatorname{ess\,sup}_{(t,s) \in [a,b]^2} |f(t) - f(s)|.$$

If we consider $f_\Delta : [a, b]^2 \rightarrow \mathbb{R}$,

$$f_\Delta(t, s) = f(t) - f(s),$$

then obviously

$$\|f\|_p^\Delta = \|f_\Delta\|_p, \quad p \in [1, \infty),$$

where $\|\cdot\|_p$ are the usual Lebesgue p -norms on $[a, b]^2$.

Using the properties of the Lebesgue p -norms, we may deduce the following seminorm properties for $\|\cdot\|_p^\Delta$:

- (i) $\|f\|_p^\Delta \geq 0$ for $f \in L_p[a, b]$ and $\|f\|_p^\Delta = 0$ implies that $f = c$ (c is a constant) a.e. in $[a, b]$,
- (ii) $\|f + g\|_p^\Delta \leq \|f\|_p^\Delta + \|g\|_p^\Delta$ if $f, g \in L_p[a, b]$,
- (iii) $\|\alpha f\|_p^\Delta = |\alpha| \|f\|_p^\Delta$.

We call $\|\cdot\|_p^\Delta$ as Δ -seminorms.

We note that if $p = 2$, then

$$\begin{aligned} \|f\|_2^\Delta &= \left(\int_a^b \int_a^b (f(t) - f(s))^2 dt ds \right)^{\frac{1}{2}} \\ &= \sqrt{2} \left((b-a) \|f\|_2^2 - \left(\int_a^b f(t) dt \right)^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Using the inequalities (1), (3) and (5), we obtain the following estimate for $\|\cdot\|_2^\Delta$:

a) for $m \leq f \leq M$

$$\|f\|_2^\Delta \leq \frac{\sqrt{2}}{2} (M - m)(b - a),$$

b) for $f' \in L_\infty[a, b]$

$$\|f\|_2^\Delta \leq \frac{\sqrt{2}}{2\sqrt{3}} \|f'\|_\infty (b-a)^2,$$

c) for $f' \in L_2[a, b]$

$$\|f\|_2^\Delta \leq \frac{\sqrt{2}}{\pi} \|f'\|_2 (b-a)^{\frac{3}{2}},$$

since

$$\|f\|_2^\Delta = \sqrt{2}(b-a)[C(f, f)]^{\frac{1}{2}}.$$

If $f: [a, b] \rightarrow \mathbb{R}$ is absolutely continuous on $[a, b]$, then we can point out the following bounds for $\|f\|_p^\Delta$ in terms of $\|f'\|_p$.

Theorem 3. *Assume that $f: [a, b] \rightarrow \mathbb{R}$ is absolutely continuous on $[a, b]$.*

(i) *If $p \in [1, \infty)$, then we have the inequality*

a) *for $f' \in L_\infty[a, b]$*

$$\|f\|_p^\Delta \leq \frac{2^{\frac{1}{p}}(b-a)^{1+\frac{2}{p}}}{[(p+1)(p+2)]^{\frac{1}{p}}} \|f'\|_\infty,$$

b) *for $f' \in L_\alpha[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$*

$$\|f\|_p^\Delta \leq \frac{(2\beta^2)^{\frac{1}{p}}(b-a)^{\frac{1}{\beta}+\frac{2}{p}}}{[(p+\beta)(p+2\beta)]^{\frac{1}{p}}} \|f'\|_\alpha,$$

c) *for $f' \in L_1[a, b]$*

$$\|f\|_p^\Delta \leq (b-a)^{\frac{2}{p}} \|f'\|_1.$$

(ii) *If $p = \infty$, then we have the inequality*

a) *for $f' \in L_\infty[a, b]$*

$$\|f\|_\infty^\Delta \leq (b-a) \|f'\|_\infty,$$

b) *for $f' \in L_\alpha[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$*

$$\|f\|_\infty^\Delta \leq (b-a)^{\frac{1}{\beta}} \|f'\|_\alpha,$$

c) *for $f' \in L_1[a, b]$*

$$\|f\|_\infty^\Delta \leq \|f'\|_1.$$

The following result of Grüss type holds, see [4].

Theorem 4. Let $f, g: [a, b] \rightarrow \mathbb{R}$ be measurable on $[a, b]$. Then we have the inequality

$$|C(f, g)| \leq \frac{1}{2(b-a)^2} \|f\|_p^\Delta \|g\|_q^\Delta,$$

where $p = 1, q = \infty$, or $p > 1, \frac{1}{p} + \frac{1}{q} = 1$, or $q = 1, p = \infty$, provided all integrals involved exist.

The inequality is sharp in the sense that if we take $f(x) = g(x) = \operatorname{sgn}(x - \alpha)$ with $\alpha = \frac{a+b}{2}$, then the equality results.

Making use of the double integral inequality

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \inf_{\lambda \in \mathbb{R}} \left\{ \int_a^b \int_a^b |f(x) - f(t)|^p |g(x) - \lambda|^p dt dx \right\}^{1/p},$$

obtained in (9) we can state the following result as well.

Theorem 5. Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue measurable functions on $[a, b]$. Then

a) for $f \in L_\infty[a, b], g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a,b],p} \|f\|_\infty^\Delta, \quad (17)$$

b) for $f \in L_{p\beta}[a, b], g \in L_{p\alpha}[a, b], \alpha > 1, \frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p+1/p\beta}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a,b],p\alpha} \|f\|_{p\beta}^\Delta, \quad (18)$$

c) for $f \in L_p[a, b], g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \inf_{\lambda \in \mathbb{R}} \|g - \lambda\|_{[a,b],\infty} \|f\|_p^\Delta. \quad (19)$$

Proof. Utilising Hölder's inequality for double integrals, we have

a) for $f \in L_\infty[a, b], g \in L_p[a, b]$

$$\begin{aligned} \int_a^b \int_a^b |f(x) - f(t)|^p |g(x) - \lambda|^p dt dx &\leq \operatorname{ess\,sup}_{(x,t) \in [a,b]^2} |f(x) - f(t)|^p \\ &\quad \times \int_a^b \int_a^b |g(x) - \lambda|^p dt dx \\ &= (\|f\|_\infty^\Delta)^p (b-a) \|g - \lambda\|_{[a,b],p}^p. \end{aligned}$$

Then

$$\begin{aligned} |C(f, g)|^p &\leq \frac{1}{(b-a)^2} (\|f\|_\infty^\Delta)^p (b-a) \|g - \lambda\|_{[a,b],p}^p \\ &= \frac{1}{b-a} (\|f\|_\infty^\Delta)^p \|g - \lambda\|_{[a,b],p}^p. \end{aligned}$$

b) For $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, we have

$$\begin{aligned} \int_a^b \int_a^b |f(x) - f(t)|^p |g(x) - \lambda|^p dt dx &\leq \left(\int_a^b \int_a^b |f(x) - f(t)|^{p\beta} dt dx \right)^{1/\beta} \\ &\quad \times \left(\int_a^b \int_a^b |g(x) - \lambda|^{p\alpha} dt dx \right)^{1/\alpha} \\ &= (\|f\|_{p\beta}^\Delta)^p (b-a)^{1/\alpha} \|g - \lambda\|_{[a,b], p\alpha}^p. \end{aligned}$$

Then

$$\begin{aligned} |C(f, g)|^p &\leq \frac{1}{(b-a)^2} (\|f\|_{p\beta}^\Delta)^p (b-a)^{1/\alpha} \|g - \lambda\|_{[a,b], p\alpha}^p \\ &= \frac{1}{(b-a)^{1+1/\beta}} (\|f\|_{p\beta}^\Delta)^p \|g - \lambda\|_{[a,b], p\alpha}^p. \end{aligned}$$

c) For $f \in L_p[a, b]$, $g \in L_\infty[a, b]$ we have

$$\begin{aligned} \int_a^b \int_a^b |f(x) - f(t)|^p |g(x) - \lambda|^p dt dx &\leq \operatorname{ess\,sup}_{x \in [a,b]} |g(x) - \lambda|^p \\ &\quad \times \int_a^b \int_a^b |f(x) - f(t)|^p dt dx \\ &= \|g - \lambda\|_{[a,b], \infty}^p (\|f\|_p^\Delta)^p. \end{aligned}$$

Then

$$|C(f, g)|^p \leq \frac{1}{(b-a)^2} \|g - \lambda\|_{[a,b], \infty}^p (\|f\|_p^\Delta)^p.$$

Taking the power $\frac{1}{p}$ and then the infimum over $\lambda \in \mathbb{R}$, we get the desired results. \square

Some particular cases of interest are as follows.

Corollary 5. Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue measurable functions on $[a, b]$. Then

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], p} \|f\|_\infty^\Delta,$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p+1/p\beta}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], p\alpha} \|f\|_{p\beta}^\Delta$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \left\| g - \frac{1}{b-a} \int_a^b g(t) dt \right\|_{[a,b], \infty} \|f\|_p^\Delta.$$

The case when one function is bounded is as follows.

Corollary 6. Assume that $f, g: [a, b] \rightarrow \mathbb{R}$ are Lebesgue integrable functions on $[a, b]$. If there exist constants n, N such that $n \leq g(t) \leq N$ for a.e. $t \in [a, b]$, then

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2}(N-n)\|f\|_\infty^\Delta, \quad (20)$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2}(N-n) \frac{1}{(b-a)^{2/p\beta}} \|f\|_{p\beta}^\Delta \quad (21)$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{2}(N-n) \frac{1}{(b-a)^{2/p}} \|f\|_p^\Delta. \quad (22)$$

Proof. From (17)–(19) we have

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{N+n}{2} \right\|_{[a,b], p} \|f\|_\infty^\Delta. \quad (23)$$

Since

$$\left\| g - \frac{n+N}{2} \right\|_{[a,b], p} \leq \frac{N-n}{2} (b-a)^{1/p}$$

then by (23) we get (20).

b) For $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p+1/p\beta}} \left\| g - \frac{N+n}{2} \right\|_{[a,b], p\alpha} \|f\|_{p\beta}^\Delta. \quad (24)$$

Since

$$\left\| g - \frac{n+N}{2} \right\|_{[a,b], p\alpha} \leq \frac{N-n}{2} (b-a)^{1/p\alpha}$$

then by (24) we get (21).

c) For $f \in L_p[a, b]$, $g \in L_\infty[a, b]$ we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \left\| g - \frac{N+n}{2} \right\|_{[a,b],\infty} \|f\|_p^\Delta. \quad (25)$$

Since

$$\left\| g - \frac{n+N}{2} \right\|_{[a,b],\infty} \leq \frac{N-n}{2},$$

then by (25) we get (22). □

The case when one function is of bounded variation, is as follows.

Corollary 7. *If $f: [a, b] \rightarrow \mathbb{R}$ is Lebesgue integrable and $g: [a, b] \rightarrow \mathbb{R}$ is of bounded variation, then*

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \|f\|_\infty^\Delta,$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \frac{1}{(b-a)^{2/p\beta}} \|f\|_{p\beta}^\Delta,$$

c) for $f \in L_p[a, b]$, $g \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{2} \bigvee_a^b(g) \frac{1}{(b-a)^{2/p}} \|f\|_p^\Delta.$$

Proof. From (17)–(19) we have

a) for $f \in L_\infty[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - \frac{g(a)+g(b)}{2} \right\|_{[a,b],p} \|f\|_\infty^\Delta. \quad (26)$$

Since

$$\left\| g - \frac{g(a)+g(b)}{2} \right\|_{[a,b],p} \leq \frac{1}{2} \bigvee_a^b(g) (b-a)^{1/p},$$

then by (26) we get the desired result.

b) For $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p+1/p\beta}} \left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b], p\alpha} \|f\|_{p\beta}^{\Delta}. \quad (27)$$

Since

$$\left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b], p\alpha} \leq \frac{1}{2} \bigvee_a^b(g) (b-a)^{1/p\alpha},$$

then by (27) we get the desired result.

c) For $f \in L_p[a, b]$, $g \in L_{\infty}[a, b]$ we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b], \infty} \|f\|_p^{\Delta}. \quad (28)$$

Since

$$\left\| g - \frac{g(a) + g(b)}{2} \right\|_{[a,b], \infty} \leq \frac{1}{2} \bigvee_a^b(g),$$

then by (28) we get the desired result. \square

Corollary 8. *If $f: [a, b] \rightarrow \mathbb{R}$ is Lebesgue integrable and $g: [a, b] \rightarrow \mathbb{R}$ is Lipschitzian in the middle point with the constant $L_{\frac{a+b}{2}}$ and the exponent $q > 0$, then*

a) for $f \in L_{\infty}[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2^q} L_{\frac{a+b}{2}} \frac{(b-a)^q}{(qp+1)^{1/p}} \|f\|_{\infty}^{\Delta},$$

b) for $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2^q} L_{\frac{a+b}{2}} \frac{(b-a)^{q-2/p\beta}}{(qp\alpha+1)^{1/p\alpha}} \|f\|_{p\beta}^{\Delta},$$

c) for $f \in L_p[a, b]$, $g \in L_{\infty}[a, b]$

$$|C(f, g)| \leq \frac{1}{2^q} L_{\frac{a+b}{2}} (b-a)^{q-2/p} \|f\|_p^{\Delta}.$$

Proof. From (17)–(19) we have

a) for $f \in L_{\infty}[a, b]$, $g \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b], p} \|f\|_{\infty}^{\Delta}. \quad (29)$$

Since

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b], p} \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p}}{2^q (qp+1)^{1/p}},$$

then from (29) we deduce the desired result.

b) For $f \in L_{p\beta}[a, b]$, $g \in L_{p\alpha}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{1/p+1/p\beta}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],p\alpha} \|f\|_{p\beta}^\Delta. \tag{30}$$

Since

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],\alpha} \leq L_{\frac{a+b}{2}} \frac{(b-a)^{q+1/p\alpha}}{2^q (qp\alpha + 1)^{1/p\alpha}},$$

then from (30) we deduce the desired result.

c) For $f \in L_p[a, b]$, $g \in L_\infty[a, b]$ we have

$$|C(f, g)| \leq \frac{1}{(b-a)^{2/p}} \left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],\infty} \|f\|_p^\Delta. \tag{31}$$

Since

$$\left\| g - g\left(\frac{a+b}{2}\right) \right\|_{[a,b],\infty} \leq L_{\frac{a+b}{2}} \frac{(b-a)^q}{2^q},$$

then from (31) we deduce the desired result. □

Remark 3. If the function g is Lipschitzian with the constant $L > 0$, then

a) for $f \in L_\infty[a, b]$

$$|C(f, g)| \leq \frac{1}{2} L \frac{b-a}{(p+1)^{1/p}} \|f\|_\infty^\Delta,$$

b) for $f \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$|C(f, g)| \leq \frac{1}{2} L \frac{(b-a)^{1-2/p\beta}}{(p\alpha + 1)^{1/p\alpha}} \|f\|_{p\beta}^\Delta,$$

c) for $f \in L_p[a, b]$

$$|C(f, g)| \leq \frac{1}{2} L (b-a)^{1-2/p} \|f\|_p^\Delta.$$

4 Applications for Mid-point Inequalities

Consider absolutely continuous function $h: [a, b] \rightarrow \mathbb{R}$. We have the following well known representation

$$h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt = \frac{1}{b-a} \int_a^b K(t) h'(t) dt,$$

where the kernel $K: [a, b] \rightarrow \mathbb{R}$ is defined by

$$K(t) := \begin{cases} t-a & \text{if } t \in [a, \frac{a+b}{2}], \\ t-b & \text{if } t \in (\frac{a+b}{2}, b]. \end{cases}$$

Since $\int_a^b K(t) dt = 0$, then

$$\frac{1}{b-a} \int_a^b K(t)h'(t) dt = C(K, h').$$

Utilising Corollary 1 we have

a) for $h' \in L_\infty[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|K\|_{[a,b],p} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],\infty}, \quad (32)$$

b) for $h' \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|K\|_{[a,b],p\alpha} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],p\beta}, \quad (33)$$

c) for $h' \in L_p[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|K\|_{[a,b],\infty} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],p}. \quad (34)$$

Observe that for $q > 0$ we have

$$\begin{aligned} \|K\|_{[a,b],q} &= \left[\int_a^b |K(t)|^q dt \right]^{1/q} \\ &= \left[\int_a^{\frac{a+b}{2}} (t-a)^q dt + \int_{\frac{a+b}{2}}^b (b-t)^q dt \right]^{1/q} \\ &= \left[\frac{(t-a)^{q+1}}{q+1} \Big|_a^{\frac{a+b}{2}} - \frac{(b-t)^{q+1}}{q+1} \Big|_{\frac{a+b}{2}}^b \right]^{1/q} \\ &= \left[\frac{\left(\frac{b-a}{2}\right)^{q+1}}{q+1} + \frac{\left(\frac{b-a}{2}\right)^{q+1}}{q+1} \right]^{1/q} = \frac{(b-a)^{1+1/q}}{2(q+1)^{1/q}}. \end{aligned}$$

Then

$$\|K\|_{[a,b],p} = \frac{(b-a)^{1+1/p}}{2(p+1)^{1/p}}, \quad \|K\|_{[a,b],p\alpha} = \frac{(b-a)^{1+1/p\alpha}}{2(p\alpha+1)^{1/p\alpha}}.$$

We also have

$$\|K\|_{[a,b],\infty} = \frac{1}{2}(b-a).$$

Making use of (32)–(34) we get

a) for $h' \in L_\infty[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{b-a}{2(p+1)^{1/p}} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],\infty},$$

b) for $h' \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{(b-a)^{1-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],p\beta},$$

c) for $h' \in L_p[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(b-a)^{1-1/p} \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],p}.$$

For $p = 1$ we get the simpler inequalities

a) for $h' \in L_\infty[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{4}(b-a) \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],\infty},$$

b) for $h' \in L_1[a, b]$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(b-a) \left\| h' - \frac{h(b)-h(a)}{b-a} \right\|_{[a,b],1}.$$

Utilising Corollary 2 we have

a)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \|K\|_{[a,b],\infty},$$

b) for $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{1}{(b-a)^{1/p\beta}} \|K\|_{[a,b],p\beta},$$

c)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{1}{(b-a)^{1/p}} \|K\|_{[a,b],p},$$

provided that $\gamma \leq h'(t) \leq \Gamma$ for a.e. $t \in [a, b]$.

Utilising the above calculations we then have

a)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq (\Gamma - \gamma)(b-a), \tag{35}$$

b) for $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{(b-a)^{1+1/p\alpha-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}}, \tag{36}$$

c)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{b-a}{2(p+1)^{1/p}}, \quad (37)$$

provided that $\gamma \leq h'(t) \leq \Gamma$ for a.e. $t \in [a, b]$.

In particular, for $p = 1$ in (37) we have

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8}(\Gamma - \gamma)(b-a),$$

which is the best inequality one can get from (35)–(37).

If we use Corollary 3 and assume that h' is of bounded variation on $[a, b]$, then

a)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \sqrt[p]{h'(b-a)},$$

b) for $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2} \sqrt[p]{h'(b-a)} \frac{(b-a)^{1+1/p\alpha-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}},$$

c)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2} \sqrt[p]{h'(b-a)} \frac{b-a}{2(p+1)^{1/p}}. \quad (38)$$

From (38) for $p = 1$ we get

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8}(b-a) \sqrt[p]{h'}.$$

If we use inequalities (14)–(16) and assume that h' is Lipschitzian with the constant $U > 0$, namely

$$|h'(t) - h'(s)| \leq U|t - s| \text{ for } t, s \in (a, b),$$

then

a)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^2}{(p+1)^{1/p}},$$

b)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^{2-1/p\beta+1/p\alpha}}{(p\alpha+1)^{2/p\alpha}},$$

c)

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^2}{(p+1)^{1/p}}.$$

In particular, we get for $p = 1$

$$\left| h\left(\frac{a+b}{2}\right) - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8} (b-a)^2 U.$$

5 Applications for Trapezoid Inequalities

Consider absolutely continuous function $h: [a, b] \rightarrow \mathbb{R}$. We have the following well known representation

$$\frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt = \frac{1}{b-a} \int_a^b V(t) h'(t) dt$$

where the kernel $V: [a, b] \rightarrow \mathbb{R}$ is defined by

$$V(t) := t - \frac{a+b}{2}.$$

Since $\int_a^b V(t) dt = 0$, then

$$\frac{1}{b-a} \int_a^b V(t) h'(t) dt = C(V, h').$$

Utilising Corollary 1 we have

a) for $h' \in L_\infty[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|V\|_{[a,b],p} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],\infty}, \quad (39)$$

b) for $h' \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|V\|_{[a,b],p\alpha} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],p\beta}, \quad (40)$$

c) for $h' \in L_p[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{(b-a)^{1/p}} \|V\|_{[a,b],\infty} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],p}. \quad (41)$$

Observe that, for $q > 0$ we have

$$\begin{aligned} \|V\|_{[a,b],q} &= \left[\int_a^b |V(t)|^q dt \right]^{1/q} = \left[\int_a^{\frac{a+b}{2}} \left(\frac{a+b}{2} - t \right)^q dt + \int_{\frac{a+b}{2}}^b \left(t - \frac{a+b}{2} \right)^q dt \right]^{1/q} \\ &= \left[2 \int_{\frac{a+b}{2}}^b \left(t - \frac{a+b}{2} \right)^q dt \right]^{1/q} = \left[\frac{2 \left(\frac{b-a}{2} \right)^{q+1}}{q+1} \right]^{1/q} = \frac{(b-a)^{1+1/q}}{2(q+1)^{1/q}}. \end{aligned}$$

Then

$$\|V\|_{[a,b],p} = \frac{(b-a)^{1+1/p}}{2(p+1)^{1/p}}, \quad \|V\|_{[a,b],p\alpha} = \frac{(b-a)^{1+1/p\alpha}}{2(p\alpha+1)^{1/p\alpha}}.$$

We also have

$$\|V\|_{[a,b],\infty} = \frac{1}{2}(b-a).$$

Making use of (39)–(41) we get

a) for $h' \in L_\infty[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{b-a}{2(p+1)^{1/p}} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],\infty},$$

b) for $h' \in L_{p\beta}[a, b]$, $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{(b-a)^{1-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],p\beta},$$

c) for $h' \in L_p[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(b-a)^{1-1/p} \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],p}.$$

For $p = 1$ we get the simpler inequalities

a) for $h' \in L_\infty[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{4}(b-a) \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],\infty},$$

b) for $h' \in L_1[a, b]$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(b-a) \left\| h' - \frac{h(b) - h(a)}{b-a} \right\|_{[a,b],p}.$$

Since the p -norms of the kernel V are the same as of K , then we can state the following results as well.

If $\gamma \leq h'(t) \leq \Gamma$ for a.e. $t \in [a, b]$, then we have

a)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq (\Gamma - \gamma)(b-a), \quad (42)$$

b) for $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{(b-a)^{1+1/p\alpha-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}}, \quad (43)$$

c)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2}(\Gamma - \gamma) \frac{b-a}{2(p+1)^{1/p}}. \quad (44)$$

In particular, for $p = 1$ in (44) we have

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8}(\Gamma - \gamma)(b-a),$$

which is the best inequality one can get from (42)–(44).

If h' is of bounded variation on $[a, b]$, then

a)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \bigvee_a^b(h')(b-a),$$

b) for $\alpha > 1$, $\frac{1}{\alpha} + \frac{1}{\beta} = 1$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2} \bigvee_a^b(h') \frac{(b-a)^{1+1/p\alpha-1/p\beta}}{2(p\alpha+1)^{1/p\alpha}},$$

c)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{2} \bigvee_a^b(h') \frac{b-a}{2(p+1)^{1/p}}.$$

From (38) for $p = 1$ we get

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8}(b-a) \bigvee_a^b(h').$$

Assume that h' is Lipschitzian with the constant $U > 0$ then

a)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^2}{(p+1)^{1/p}},$$

b)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^{2-1/p\beta+1/p\alpha}}{(p\alpha+1)^{2/p\alpha}},$$

c)

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq U \frac{1}{4} \frac{(b-a)^2}{(p+1)^{1/p}}.$$

In particular, we get for $p = 1$

$$\left| \frac{h(b) + h(a)}{2} - \frac{1}{b-a} \int_a^b h(t) dt \right| \leq \frac{1}{8}(b-a)^2 U.$$

Some similar inequalities may be stated in terms of the Δ -seminorms. However the details are omitted.

6 Some Exponential Inequalities

We can state the following result.

Theorem 6. *Let $f, g: [a, b] \rightarrow \mathbb{R}$ be Lebesgue integrable functions on $[a, b]$. If $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ is convex and monotonic nondecreasing on \mathbb{R} then we have the inequality*

$$\Phi[C(f, g)] \leq \frac{1}{b-a} \inf_{\mu \in \mathbb{R}} \int_a^b \Phi \left[\left(\frac{f(x) + g(x)}{2} - \mu \right)^2 \right] dx. \quad (45)$$

Proof. From Theorem 1 we have

$$\begin{aligned} & \Phi[C(f, g)] \\ & \leq \frac{1}{b-a} \int_a^b \Phi \left[\left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) \left(g(x) - 2\mu + \frac{1}{b-a} \int_a^b f(t) dt \right) \right] dx \end{aligned}$$

for any $\mu \in \mathbb{R}$.

Utilising the elementary inequality

$$\alpha\beta \leq \left(\frac{\alpha + \beta}{2} \right)^2$$

that holds for any $\alpha, \beta \in \mathbb{R}$, we have

$$\left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) \left(g(x) - 2\mu + \frac{1}{b-a} \int_a^b f(t) dt \right) \leq \left(\frac{f(x) + g(x)}{2} - \mu \right)^2$$

for any $x \in [a, b]$.

Since $\Phi: \mathbb{R} \rightarrow \mathbb{R}$ is monotonic nondecreasing on \mathbb{R} then

$$\begin{aligned} & \Phi \left[\left(f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right) \left(g(x) - 2\mu + \frac{1}{b-a} \int_a^b f(t) dt \right) \right] \\ & \leq \Phi \left[\left(\frac{f(x) + g(x)}{2} - \mu \right)^2 \right] \end{aligned} \quad (46)$$

for any $x \in [a, b]$.

Integrating (46) over x in $[a, b]$ and taking the infimum over $\mu \in \mathbb{R}$, we deduce the desired result (45). \square

Remark 4. Writing the inequality (45) for $\Phi: \mathbb{R} \rightarrow \mathbb{R}$, $\Phi(x) = \exp x$ we have

$$\exp[C(f, g)] \leq \frac{1}{b-a} \inf_{\mu \in \mathbb{R}} \int_a^b \exp \left[\left(\frac{f(x) + g(x)}{2} - \mu \right)^2 \right] dx. \quad (47)$$

This inequality can provide some exponential inequalities as follows.

Assume that $f: [a, b] \rightarrow \mathbb{R}$ is Lipschitzian with constant $L > 0$ and $g: [a, b] \rightarrow \mathbb{R}$ is Lipschitzian with constant $K > 0$. Then by taking

$$\mu = \frac{f\left(\frac{a+b}{2}\right) + g\left(\frac{a+b}{2}\right)}{2}$$

we have

$$\left(\frac{f(x) + g(x)}{2} - \frac{f\left(\frac{a+b}{2}\right) + g\left(\frac{a+b}{2}\right)}{2}\right)^2 \leq \left(\frac{L + K}{2}\right)^2 \left(x - \frac{a + b}{2}\right)^2$$

and by (47) we have

$$\exp[C(f, g)] \leq \frac{1}{b - a} \int_a^b \exp \left[\left(\frac{L + K}{2}\right)^2 \left(x - \frac{a + b}{2}\right)^2 \right] dx .$$

Acknowledgement

The author would like to thank the anonymous referees for their valuable comments that have been implemented in the final version of the paper.

References

- [1] P. Cerone, S.S. Dragomir: Some bounds in terms of Δ -seminorms for Ostrowski-Grüss type inequalities. *Soochow J. Math.* 27 (4) (2001) 423–434.
- [2] P. Cerone, S.S. Dragomir: New bounds for the Čebyšev functional. *App. Math. Lett.* 18 (2005) 603–611.
- [3] P. Cerone, S.S. Dragomir: A refinement of the Grüss inequality and applications. *Tamkang J. Math.* 38 (1) (2007) 37–49. Preprint *RGMIA Res. Rep. Coll.* 5 (2) (2002), Art. 14. [online <http://rgmia.vu.edu.au/v8n2.html>]
- [4] P. Cerone, S.S. Dragomir, J. Roumeliotis: Grüss inequality in terms of Δ -seminorms and applications. *Integral Transforms Spec. Funct.* 14 (3) (2003) 205–216.
- [5] P.L. Chebyshev: Sur les expressions approximatives des intégrals définis par les autres prises entre les même limites. *Proc. Math. Soc. Charkov* 2 (1882) 93–98.
- [6] X.-L. Cheng, J. Sun: Note on the perturbed trapezoid inequality. *J. Ineq. Pure & Appl. Math.* 3 (2) (2002). Art. 29. [online <http://jipam.vu.edu.au/article.php?sid=181>]
- [7] G. Grüss: Über das Maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x) dx - \frac{1}{(b-a)^2} \int_a^b f(x) dx \int_a^b g(x) dx$. *Math. Z.* 39 (1935) 215–226.
- [8] X. Li, R.N. Mohapatra, R.S. Rodriguez: Grüss-type inequalities. *J. Math. Anal. Appl.* 267 (2) (2002) 434–443.
- [9] A. Lupaş: The best constant in an integral inequality. *Mathematica (Cluj, Romania)* 15 (38) (2) (1973) 219–222.
- [10] A.McD. Mercer: An improvement of the Grüss inequality. *J. Inequal. Pure Appl. Math.* 6 (4) (2005). Article 93, 4 pp. (electronic).
- [11] D.S. Mitrinović, J.E. Pečarić, A.M. Fink: *Classical and New Inequalities in Analysis*. Kluwer Academic Publishers, Dordrecht/Boston/London (1993).
- [12] A.M. Ostrowski: On an integral inequality. *Aequat. Math.* 4 (1970) 358–373.
- [13] B.G. Pachpatte: On Grüss like integral inequalities via Pompeiu’s mean value theorem. *J. Inequal. Pure Appl. Math.* 6 (3) (2005). Article 82, 5 pp.

Author's address:

MATHEMATICS, COLLEGE OF ENGINEERING & SCIENCE, VICTORIA UNIVERSITY,
PO BOX 14428, MELBOURNE CITY, MC 8001, AUSTRALIA
SCHOOL OF COMPUTATIONAL & APPLIED MATHEMATICS, UNIVERSITY OF THE
WITWATERSRAND, PRIVATE BAG 3, JOHANNESBURG 2050, SOUTH AFRICA

E-mail: `sever.dragomir@vu.edu.au`

Received: 28 January, 2014

Accepted for publication: 12 April, 2014

Communicated by: Olga Rossi