

Lev Bukovský  
Balcar's theorem on supports

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 59 (2018), No. 4, 443–449

Persistent URL: <http://dml.cz/dmlcz/147549>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## Balcar’s theorem on supports

LEV BUKOVSKÝ

*To the memory of Bohuslav Balcar*

*Abstract.* In *A theorem on supports in the theory of semisets* [Comment. Math. Univ. Carolinae **14** (1973), no. 1, 1–6] B. Balcar showed that if  $\sigma \subseteq D \in M$  is a support,  $M$  being an inner model of ZFC, and  $\mathcal{P}(D \setminus \sigma) \cap M = r\text{“}\sigma$  with  $r \in M$ , then  $r$  determines a preorder “ $\preceq$ ” of  $D$  such that  $\sigma$  becomes a filter on  $(D, \preceq)$  generic over  $M$ . We show that if the relation  $r$  is replaced by a function  $\mathcal{P}(D \setminus \sigma) \cap M = f_{-1}(\sigma)$ , then there exists an equivalence relation “ $\sim$ ” on  $D$  and a partial order on  $D/\sim$  such that  $D/\sim$  is a complete Boolean algebra,  $\sigma/\sim$  is a generic filter and  $[f(u)]_{\sim} = -\sum(u/\sim)$  for any  $u \subseteq D$ ,  $u \in M$ .

*Keywords:* inner model; support; generic filter

*Classification:* 03E40

### 1. Introduction

B. Balcar in [1] has found very nice and important proof of a theorem on supports by P. Vopěnka in [4]. Actually, B. Balcar proved a theorem of the theory of semisets. The translations of the theorem to the set theory is rather immediate. Following ideas of this proof we show a related result.

We shall follow the terminology and notations of T. Jech in [3].

If  $M$  is an inner model then a set  $\sigma \subseteq D \in M$ ,  $\sigma \notin M$  is a *support over  $M$*  if for any binary relations  $r_1, r_2 \in M$  there exists a binary relation  $r \in M$  such that

$$r\text{“}\sigma = r_1\text{“}\sigma \setminus r_2\text{“}\sigma.$$

Note that  $r\text{“}\sigma = \{y \in \text{ran}(r) : \exists x \in \sigma [x, y] \in r\}$ . If  $f$  is a function, then  $f_{-1}(\sigma) = \{x \in \text{dom}(f) : f(x) \in \sigma\}$ . If “ $\sim$ ” is an equivalence relation on a set  $A$ , then for any  $x \in A$  we denote the equivalence class of  $x$  by  $[x]_{\sim}$ . If  $B \subseteq A$ , then  $B/\sim = \{[x]_{\sim} : x \in B\}$ .

By Balcar’s proof in [1], see also [2, pages 365–370], we obtain

**Theorem 1.** *Let  $M$  be an inner model. Let  $\sigma$  be a subset of a set  $D \in M$ ,  $\sigma \notin M$ . Then the following are equivalent*

- a)  $\sigma$  is a support over  $M$ ;
- b) there exists a binary relation  $r \in M$  such that  $\mathcal{P}(D \setminus \sigma) \cap M = r\text{“}\sigma$ ;

- c) there exists a preorder “ $\preceq$ ” on  $D$  such that  $\sigma$  is a filter on  $(D, \preceq)$  generic over  $M$ ;
- d) there exist a Boolean algebra  $B \in M$  complete in  $M$ , a filter  $G \subseteq B$  generic over  $M$ , a binary relation  $r \in M$  and a function  $f \in M$  such that  $G = r\sigma$  and  $\sigma = f_{-1}(G)$ .

P. Vopěnka in [4] has proved the implication a)  $\Rightarrow$  d). B. Balcar in [1] has proved the implication b)  $\Rightarrow$  c) of Theorem 1. The other implications are known from the theory of semisets and from the theory of Boolean valued models.

The condition c) of Theorem 1 is equivalent to the following:

- o there exist an equivalence relation  $\sim \in M$  on  $D$
- o and a partial order  $\leq \in M$  on  $D/\sim$  such that
- o  $\sigma/\sim$  is a filter on  $(D/\sim, \leq)$  generic over  $M$ .

If  $B \in M$  is a Boolean algebra complete in  $M$  and  $G \subseteq B$  is a filter generic over  $M$ , we define a function  $f: \mathcal{P}(B) \cap M \rightarrow B$  as

$$(1) \quad f(u) = - \sum u \quad \text{for } u \subseteq B, u \in M.$$

Then the condition b) of Theorem 1 holds true with  $f_{-1}$  instead a binary relation as

$$(2) \quad P(B \setminus G) \cap M = f_{-1}(G).$$

We shall study how we can change the assertion c) if we replace the relation  $r$  in b) by the inverse of a function  $f_{-1}$  as above. We show

**Theorem 2.** Assume that  $M$  is an inner model. Let  $\sigma$  be a subset of a set  $D \in M$ ,  $\sigma \notin M$ , and let  $f: \mathcal{P}(D) \cap M \rightarrow D$  be a function in the model  $M$ . Then the following are equivalent

- a)  $P(D \setminus \sigma) \cap M = f_{-1}(\sigma)$ ;
- b) there exist an equivalence relation  $\sim \in M$  on  $D$  and a partial order  $\leq \in M$  on  $D/\sim$  such that  $(D/\sim, \leq)$  is a Boolean algebra complete in  $M$ ,  $\sigma/\sim$  is a filter on  $(D/\sim, \leq)$  generic over  $M$ , and for any  $u \subseteq D$ ,  $u \in M$ , we have

$$[f(u)]_{\sim} = - \sum \{[x]_{\sim} : x \in u\}.$$

## 2. Getting a partial order

In the next we assume that

$$D \in M, \sigma \subseteq D, \quad f: P(D) \cap M \rightarrow D, \quad f \in M, \quad P(D \setminus \sigma) \cap M = f_{-1}(\sigma).$$

We set

$$(3) \quad s(x) = \{y \in D : \exists u \in P(D) \cap M \quad u \cap \{x, y\} \neq \emptyset \wedge f(u) \in \{x, y\}\}.$$

The intended interpretation is that  $s(x)$  is the set of elements of the partially preordered set  $D$  incompatible with  $x$ .

Immediately from the definition we obtain

$$y \in s(x) \rightarrow x \in s(y) \quad \text{for any } x, y \in D,$$

and

$$(4) \quad u \subseteq s(f(u)) \quad \text{for any } u \subseteq D, u \in M.$$

If  $x \in s(x)$ , then by definition there is  $u \in P(D) \cap M$  such that  $x \in u$  and  $f(u) = x$ . Hence  $x \notin \sigma$ . Moreover then  $u \cap \{x, y\} \neq \emptyset$  and  $f(u) \in \{x, y\}$  for every  $y \in D$ . Thus

$$(5) \quad \text{if } x \in s(x), \text{ then } x \notin \sigma,$$

and

$$(6) \quad x \in s(x) \text{ if and only if } s(x) = D.$$

If we take  $u = \{x\}$  in the definition (3), we obtain

$$x \in s(f(\{x\})) \quad \text{and} \quad f(\{x\}) \in s(x) \quad \text{for any } x \in D.$$

We define a preorder " $\preceq$ " of the set  $D$  setting

$$x \preceq y \text{ if and only if } s(y) \subseteq s(x).$$

By (6), any  $x \in D$  such that  $x \in s(x)$  plays the role of the least element in this preorder.

The preorder " $\preceq$ " induces an equivalence relation " $\approx$ " defined by

$$x \approx y \text{ if and only if } s(x) = s(y).$$

The preorder " $\preceq$ " becomes a partial order on the quotient set  $D/\approx$ . We identify elements of  $D$  with their equivalence classes and subsets of  $D$  with the set of corresponding equivalence classes. So, speaking about  $x \in D$  we mean the equivalence class  $[x]_{\approx}$  in  $D/\approx$ . Similarly for a subset of  $D$ .

Let

$$A_0 = \{x \in D : x \in s(x)\}, \quad D_0 = D \setminus A_0.$$

It is easy to see that  $A_0$  is hereditary downward, i.e.,

$$(7) \quad (y \preceq x \wedge x \in A_0) \rightarrow y \in A_0.$$

Since  $f(D) \in D = s(f(D))$  we have

$$(8) \quad D \setminus D_0 \neq \emptyset.$$

By (6),  $A_0$  is the  $\preceq$ -least equivalence class of  $D/\approx$  and  $D_0/\approx$  is the set of nonzero elements of  $D/\approx$ .

Note the following:

$$\text{if } z \preceq y \text{ and } y \in s(x), \text{ then } z \in s(x).$$

Indeed, if  $z \preceq y$  and  $y \in s(x)$ , then  $x \in s(y) \subseteq s(z)$ . Hence  $z \in s(x)$ .

Let  $X \subseteq D$ . We say that elements  $x, y \in D$  are *incompatible in  $X$* , if for every  $z \preceq x, z \preceq y$  we have  $z \notin X$ . If  $x, y \in D_0$  are incompatible in  $D_0$  we shall write  $x \perp y$ .

We show that for any  $x \in D_0$ , every element of the set  $s(x) \cap D_0$  is incompatible with  $x$ .

$$\text{If } y \in s(x), z \preceq x, z \preceq y, x, y \in D_0, \text{ then } z \notin D_0.$$

So, assume that  $y \in s(x)$  and  $s(x) \subseteq s(z), s(y) \subseteq s(z)$ . Then  $y \in s(z)$  and therefore also  $z \in s(y)$ . Thus  $z \in s(z)$ . Hence

$$(9) \quad \text{if } y \in s(x), \text{ then } x \perp y \text{ for any } x, y \in D_0.$$

By (4) we obtain that

$$(10) \quad \text{if } s(x) \cup s(f(u)) \subseteq s(y) \text{ for some } x \in u, \text{ then } y \in s(y).$$

In particular,

$$\text{if } s(x) \cup s(f(\{x\})) \subseteq s(y) \text{ for some } x, \text{ then } y \in s(y).$$

**Lemma 3.** a)  $s(x) \subseteq D \setminus \sigma$  for each  $x \in \sigma$ .

b) If  $u \subseteq D \setminus \sigma, u \in M$ , then there exists an  $x \in \sigma$  such that  $u \subseteq s(x)$ .

PROOF: a) Let  $x \in \sigma$ . If  $y \in s(x)$ , then by (3) there exists a set  $u \in P(D) \cap M$  such that

$$u \cap \{x, y\} \neq \emptyset \wedge f(u) \in \{x, y\}.$$

We have four possibilities. If  $x \in u$  and  $f(u) = x$  then  $u \subseteq D \setminus \sigma$ , a contradiction. If  $x \in u$  and  $f(u) = y$  then  $u \not\subseteq D \setminus \sigma$ , hence we obtain that  $y = f(u) \notin \sigma$ . If  $y \in u$  and  $f(u) = x$  then  $u \subseteq D \setminus \sigma$ , therefore  $y \notin \sigma$ . If  $y \in u$  and  $f(u) = y$ , then  $y \in \sigma$  implies that  $u \subseteq D \setminus \sigma$ , a contradiction. Thus  $y \notin \sigma$ .

b) Let  $u \subseteq D \setminus \sigma, u \in M$ . Then  $x = f(u) \in \sigma$ . By (4) we obtain  $u \subseteq s(x)$ .  $\square$

**Lemma 4.** The set  $\sigma$  is a filter on  $(D_0, \preceq)$  generic over  $M$ .

PROOF: By (5) we have  $\sigma \subseteq D_0$ .

Let  $x \in \sigma, x \preceq y$ . Assume that  $y \notin \sigma$ . Then  $s(x) \cup \{y\} \subseteq D_0 \setminus \sigma$ , hence by Lemma 3 b), there exists a  $z \in \sigma$  such that  $s(x) \cup \{y\} \subseteq s(z)$ . Since  $y \in s(z)$  also  $z \in s(y) \subseteq s(x) \subseteq D_0 \setminus \sigma$ , a contradiction.

Let  $x, y \in \sigma$ . Then by Lemma 3 a)  $s(x) \cup s(y) \subseteq D \setminus \sigma$ . Thus, by Lemma 3 b), there exists a  $z \in \sigma$  such that  $s(x) \cup s(y) \subseteq s(z)$ . Then  $z \preceq x$  and  $z \preceq y$ .

Assume that  $u \subseteq D_0 \setminus \sigma, u \in M$ . By Lemma 3 b), there exists an  $x \in \sigma$  such that  $u \subseteq s(x)$ . By (9) every element of  $u$  is incompatible with  $x$ , therefore  $u$  is not dense in  $D_0$ .  $\square$

We denote by  $q_1$  the quotient mapping  $q_1: D_0 \rightarrow D_0/\approx$  defined as  $q_1(x) = [x]_{\approx}$ . Since

$$(x \approx y \wedge x \in \sigma) \rightarrow y \in \sigma,$$

$q_1(\sigma)$  is a filter on  $\langle D_0/\approx, \preceq \rangle$  generic over  $M$ .

The function  $f: \mathcal{P}(B) \cap M \rightarrow B$  defined by (1) may be easily changed still keeping (2) true. E.g. take  $u, v \subseteq B$  such that  $f(u), f(v) \in G$ ,  $f(u) \neq f(v)$ . If you exchange the values  $f(u)$  and  $f(v)$ , (2) is true and (1) fails. However the restriction of  $f$  to  $\mathcal{P}(B \setminus (f(u) \cdot f(v)))$  will satisfy (1).

So we must consider some “inconvenient” elements in  $D$  which we must omit. It turns out that none of those elements is in  $\sigma$ . We show that

$$(11) \quad \text{if } x \perp y \text{ for every } y \in u \text{ and } x \perp f(u), \text{ then } x \notin \sigma.$$

Assume that  $x \in \sigma$ . Then by Lemma 4 we obtain  $u \subseteq D \setminus \sigma$ . Therefore  $f(u) \in \sigma$ , which is a contradiction with  $x \perp f(u)$ .

In particular,

$$\text{if } x \perp y \text{ and } x \perp f(\{y\}), \text{ then } x \notin \sigma.$$

### 3. Getting a complete Boolean algebra

There exist an equivalence relation “ $\asymp$ ” on  $D_0/\approx$  and the quotient mapping  $q_2: D_0/\approx \rightarrow (D_0/\approx)/\asymp$ , see [3, page 205], preserving inequalities “ $\preceq$ ” and “ $\preceq/\asymp$ ” and compatibility of elements in both sides, such that  $((D_0/\approx)/\asymp, \preceq/\asymp)$  is a separative partially ordered set. We denote by “ $\sim$ ” the equivalence relation on  $D_0$  defined as

$$x \sim y \text{ if and only if } [x]_{\approx} \asymp [y]_{\approx},$$

and the partial order “ $\leq$ ” on  $D_0/\sim$ , compare [3, page 205], defined as

$$[x]_{\sim} \leq [y]_{\sim} \equiv \forall z \preceq x \quad [z]_{\approx} \text{ is compatible with } [y]_{\approx} \text{ in } D_0/\approx.$$

Then  $(D_0/\sim, \leq)$  is a separative partially ordered set. We denote

$$q = q_1 * q_2: D_0 \rightarrow D_0/\sim.$$

Then  $q(\sigma)$  is a filter on  $D_0/\sim$  generic over  $M$ .

Hence there exists a Boolean algebra  $B_0 \in M$  complete in  $M$  and a mapping  $e: D_0 \rightarrow B_0$  such that  $e$  “ $D_0$  is dense in  $B_0$ , and

$$(12) \quad \forall x, y \in D_0 \quad x \preceq y \rightarrow e(x) \leq e(y),$$

$$(13) \quad \forall x, y \in D_0 \quad x \perp y \equiv e(x), e(y) \text{ are incompatible in } B_0.$$

We can assume that  $D_0/\sim \subseteq B_0$  and the partial order “ $\leq$ ” coincides with the order of the Boolean algebra  $B_0$ . Then

$$e(x) = [x]_{\sim} \quad \text{for any } x \in D_0.$$

We set

$$A = \{x \in D : \exists u \subseteq D, u \in M \quad (x \perp f(u) \wedge \forall y \in u \quad x \perp y)\},$$

$$C = D \setminus A.$$

Evidently  $A_0 \subseteq A$  and therefore  $C = D_0 \setminus A$ . By (11) we have

$$\sigma \subseteq C.$$

By (7) and by the definition, the set  $A$  is hereditary downward.

Let  $a = e(f(A)) \in B_0$ . We denote  $B = B_0|a = \{x \in B_0 : x \leq a\}$ . We set

$$h(x) = \begin{cases} e(x) \cdot a & \text{if } x \in C, \\ 0 & \text{if } x \in D \setminus C. \end{cases}$$

We show that  $h^{\text{“}C}$  is dense in  $B \setminus \{0\}$ . So let  $b \in B, b \neq 0$ . Then  $b \leq a$ . Since  $e^{\text{“}D_0}$  is dense in  $B_0 \setminus \{0\}$ , there exists  $z \in D_0$  such that  $0 \neq e(z) \leq b$ . Assume that  $z \in A$ . Then  $z \in s(f(A))$ , hence  $z \perp f(A)$ . Hence  $e(z) = e(z) \cdot a = 0$ , a contradiction. Thus  $z \in C$ .

Evidently

$$\forall x, y \in D \quad x \preceq y \rightarrow h(x) \leq h(y).$$

Let  $u \subseteq C, u \in M$ . By (10) we obtain that for any  $x \in u$ , the elements  $x$  and  $f(u)$  are incompatible in  $(D_0, \preceq)$ . Hence by (13) we obtain  $h(f(u)) \cdot \sum h^{\text{“}u} = 0$ .

By the definition of  $A$ , we have  $\sum e^{\text{“}u} + e(f(u)) \geq a$  in  $B_0$ . If  $f(u) \in C$ , then  $\sum h^{\text{“}u} + h(f(u)) = 1$  in  $B$  and

$$h(f(u)) = - \sum h^{\text{“}u} \quad \text{in } B.$$

If  $f(u) \notin C$  then  $\sum e^{\text{“}u} \geq a$ , i.e.,  $\sum h^{\text{“}u} = 1$  in  $B$  and  $-\sum h^{\text{“}u} = 0 = h(f(u))$ .

In particular we obtain that  $h(f(\{x\})) = -h(x)$  for any  $x \in C$ .

If  $x \in B, x \neq 0$  then  $x = \sum h^{\text{“}u}$ , where  $u = \{y \in C : h(y) \leq x\}$ . Since  $\sum h^{\text{“}u} = h(f(\{f(u)\}))$ , we obtain that  $B \setminus \{0\} \subseteq h^{\text{“}C}$ . By (8),  $D \setminus C \neq \emptyset$ , hence  $0 \in h^{\text{“}D}$ . Thus  $h$  is a surjection.

Since  $\sigma \subseteq C$ , we obtain that

$$\sigma / \sim = q(\sigma) \quad \text{is a filter on } B \text{ generic over } M.$$

If we redefine the equivalence relation “ $\sim$ ” as

$$x \sim y \text{ if and only if } h(x) = h(y)$$

for any  $x, y \in D$ , we obtain the assertion of the theorem.

**Acknowledgment.** I should like to thank the anonymous referee for several constructive comments which helped me to improve the presentation of the paper.

## REFERENCES

- [1] Balcar B., *A theorem on supports in the theory of semisets*, Comment. Math. Univ. Carolinae **14** (1973), no. 1, 1–6.
- [2] Balcar B., Štěpánek P., *Set Theory*, Academia, Publishing House of the Czechoslovak Academy of Sciences, Praha, 2001 (Czech).
- [3] Jech T., *Set Theory*, The Third Millenium Edition, Springer Monographs in Mathematics, Springer, 2003.
- [4] Vopěnka P., Hájek P., *The Theory of Semisets*, Academia, Publishing House of the Czechoslovak Academy of Sciences, Praha, 1972.

L. Bukovský:

INSTITUTE OF MATHEMATICS, P. J. ŠAFÁRIK UNIVERSITY, JESENNÁ 5, 04001 KOŠICE,  
SLOVAKIA

*E-mail:* lev.bukovsky@upjs.sk

(Received March 5, 2018, revised July 26, 2018)