

András Hajnal; István Juhász; Lajos Soukup
On saturated almost disjoint families

Commentationes Mathematicae Universitatis Carolinae, Vol. 28 (1987), No. 4, 629--633

Persistent URL: <http://dml.cz/dmlcz/106576>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON SATURATED ALMOST DISJOINT FAMILIES

A. HAJNAL, I. JUHÁSZ and L. SOUKUP

Abstract: An almost disjoint family $\mathcal{A} \subset [X]^\omega$ is called saturated if every subset of X not covered by finitely many elements of \mathcal{A} contains some member of \mathcal{A} . We show that in the model obtained by iteratively adding ω_1 dominating reals to V the following statement is true: On every infinite set there is a saturated almost disjoint family. The question whether this statement is true in ZFC, or even in L , remains open.

Key words: Almost disjoint family, saturated family.

Classification: 03E05, 03E35

Given a set X and a collection \mathcal{A} of subsets of X we denote by $I_{\mathcal{A}}$ the ideal on X generated by $\mathcal{A} \cup [X]^1$, i.e. the members of $I_{\mathcal{A}}$ are the sets that can be almost covered by finitely many elements of \mathcal{A} . As is usual, we write

$$I_{\mathcal{A}}^+ = P(X) \setminus I_{\mathcal{A}}.$$

Definition. An almost disjoint family $\mathcal{A} \subset [X]^\omega$ is called saturated if \mathcal{A} refines $I_{\mathcal{A}}^+$, i.e. if for every set $H \in I_{\mathcal{A}}^+$ there is some $A \in \mathcal{A}$ with $A \subset H$.

The main result of this note may now be formulated as follows:

Theorem. If P is the partial order that adds iteratively ω_1 dominating reals to V , then the following statement $(*)$ holds in V^P :

$(*)$ For every infinite set X there is a saturated almost disjoint family $\mathcal{A} \subset [X]^\omega$.

The proof of this result is based on several lemmas to be given below. We shall use D to denote the standard notion of forcing that adds a dominating real, i.e. a function $r: \omega \rightarrow \omega$ such that $r(n) > f(n)$ for all but finitely many $n \in \omega$ whenever $f \in {}^\omega \omega \cap V$, cf. [3].

Lemma 1. Let $\mathcal{A} \subset [X]^\omega$ be almost disjoint and $H \in I_{\mathcal{A}}^+$, $\mathcal{A}, H \in V$. Then in V^D , there is a set $S \in [H]^\omega$ such that $|S \cap A| < \omega$ for each $A \in \mathcal{A}$, i.e. $A \cap S = \emptyset$.

is almost disjoint.

Proof. If there are only finitely many $A \in \mathcal{A}$ with $|A \cap H| = \omega$, say A_0, \dots, A_n , then clearly every set $S \in [H \setminus \bigcup_{i=0}^n A_i]^\omega$ works, even in V .

Otherwise let $\{A_n : n \in \omega\}$ be distinct members of \mathcal{A} such that $|A_n \cap H| = \omega$ for all $n \in \omega$. Since the A_n 's are almost disjoint, the sets

$$B_n = A_n \cap H \setminus \bigcup \{A_i : i < n\}$$

are disjoint and infinite. Let us write

$$B_n = \{a_{n,i} : i \in \omega\}$$

for each $n \in \omega$.

All this was done in V , but now we claim that the set

$$S = \{a_{n,r(n)} : n \in \omega\} \in [H]^\omega$$

defined in V^D is as required. Indeed, for each $m \in \omega$ we clearly have

$$|S \cap A_m| \leq m < \omega$$

since $A_m \cap B_n = \emptyset$ whenever $n > m$. If, on the other hand, $A \in \mathcal{A} \setminus \{A_n : n \in \omega\}$ then let us consider the function $f_A \in {}^\omega \omega \cap V$ defined as follows:

$$f_A(n) = \max \{i \in \omega : a_{n,i} \in A\},$$

that is well-defined because $|A \cap B_n| < \omega$. But r dominates f_A , hence we clearly have $|A \cap S| < \omega$. \dashv

Lemma 2. (Cf. [6] or [7], Lemma 5.) If W is an extension of V that contains a new real then in W there is an almost disjoint family $\mathcal{B} \subset [\omega]^\omega$ which refines $[\omega]^\omega \cap V$.

Actually, we only need this result in the case where $W = V^D$. In order to make this note self-contained we give a proof for this special case. First recall that D consists of pairs $\langle p, f \rangle$ where p is a strictly increasing map of a natural number into ω and $f \in {}^\omega \omega$. $\langle p, f \rangle \leq \langle q, h \rangle$ iff $p \supset q$, $f(n) \geq h(n)$ for each natural number, n , and for each $k \in \text{dom}(p) \setminus \text{dom}(q)$ we have $p(k) > h(k)$. The generic dominating function will be denoted by r . Next we fix a partition $\{A_n : n < \omega\}$ of ω into ω -many infinite pieces in V .

We choose in V a bijection g between $[\omega]^{<\omega}$ and ω . Then for each $X \in [\omega]^\omega \cap V$ let us consider the set X^* defined as follows:

$$X^* = \{\min(X \cap r'' A_g(X \cap n)) : n < \omega\}.$$

We claim that

$\mathfrak{B} = \{X^* : X \in [\omega]^\omega \cap V\}$ is as required.

A standard density argument shows that whenever $X, A \in [\omega]^\omega \cap V$, we have $X \cap A \neq \emptyset$. Thus X^* is an infinite subset of X . To show that \mathfrak{B} is almost disjoint it is sufficient to observe that $X \cap Y \neq \emptyset$ implies $|X^* \cap Y^*| \leq n$ for each $X, Y \in [\omega]^\omega \cap V$. This completes the proof of the special case.

Let us denote by D_2 the notion of forcing that adds, iteratively, two dominating reals to V .

(Formally, $D_2 = \mathbb{D} * \hat{\mathbb{D}}$, where $\hat{\mathbb{D}}$ names in V the poset in $V^{\mathbb{D}}$ that adds a dominating real.) Lemmas 1 and 2 then easily imply the next result.

Lemma 3. Let $\mathcal{A} \subset [X]^\omega$ be almost disjoint, then in V^{D_2} there exists a family $\mathfrak{B} \subset [X]^\omega$ such that

- (i) $\mathcal{A} \cup \mathfrak{B}$ is almost disjoint,
- (ii) \mathfrak{B} refines $V \cap I_{\mathcal{A}}^+$.

Proof. First, by Lemma 1, we choose in $V^{\mathbb{D}}$ for each $H \in V \cap I_{\mathcal{A}}^+$ a set $S_H \in [H]^\omega$ for which $\mathcal{A} \cup \{S_H\}$ is almost disjoint and put $\mathcal{S} = \{S_H : H \in V \cap I_{\mathcal{A}}^+\}$. Let \mathcal{C} be a maximal almost disjoint subcollection of \mathcal{S} . Then, for each $S \in \mathcal{C}$ we may apply Lemma 2 (with $V^{\mathbb{D}}$ instead of V , V^{D_2} instead of W and S instead of ω) to obtain in V^{D_2} an almost disjoint collection $\mathfrak{B}(S) \subset [S]^\omega$ refining $V^{\mathbb{D}} \cap [S]^\omega$. We claim that

$$\mathfrak{B} = \bigcup \{ \mathfrak{B}(S) : S \in \mathcal{C} \}$$

is as required. That (i) holds is obvious from the choice of \mathcal{S} and \mathcal{C} .

To show (ii), consider any $H \in V \cap I_{\mathcal{A}}^+$. By the maximality of \mathcal{C} there is some $S \in \mathcal{C}$ with $|S \cap S_H| = \omega$, but then we have a set $B \in \mathfrak{B}(S)$ with

$$B \subset S \cap S_H \subset H,$$

which was to be shown.

We are now ready to give the proof of our main result.

Proof of the theorem. We may clearly consider $P = P_{\omega_1}^P$ as given by the finite support iteration

$$\langle P_\alpha : \alpha < \omega_1, Q_\alpha : \alpha < \omega_1 \rangle, \text{ where } V^{P_\alpha} \models Q_\alpha = D_2$$

for each $\alpha < \omega_1$.

To prove that $(*)$ holds in V^P it will clearly suffice to show it for

for $X \in V$. Now, given such an X , we define almost disjoint families

$\mathcal{A}_\alpha \subset [X]^\omega$ with $\mathcal{A}_\alpha \in V_\alpha^P$ by induction on $\alpha \in \omega_1$ as follows.

We set $\mathcal{A}_0 = \emptyset$ and for every limit α we put $\mathcal{A}_\alpha = \bigcup \{ \mathcal{A}_\beta : \beta \in \alpha \}$. Standard tricks (cf. e.g. [5] p.281) concerning the choices made in the successor steps will insure that $\mathcal{A}_\alpha \in V_\alpha^P$.

Now, if $\alpha = \beta + 1$ and \mathcal{A}_β has already been defined then we can apply Lemma 3 to get a collection $\mathcal{B}_\beta \in V_\alpha^P = V_\beta^P \ast D_2$ such that $\mathcal{A}_\beta \cup \mathcal{B}_\beta \subset [X]^\omega$ is almost disjoint and \mathcal{B}_β refines $V_\beta^P \cap I_{\mathcal{A}_\beta}^+$. We then put $\mathcal{A}_\alpha = \mathcal{A}_\beta \cup \mathcal{B}_\beta$.

Since Lemma 3 involves choices (e.g. of the family $\mathcal{C} \subset \mathcal{F}$), the tricks we referred to above consist in making these choices "uniform" by fixing a large enough cardinal κ and a well-ordering \prec of $V(\kappa)$ before we start our induction so that all the relevant sets we have to choose from, or rather names for them, already occur in $V(\kappa)$, and then every choice we have to make will be the \prec -least one.

If someone is not convinced by this argument, there is another way to get around this difficulty that makes use of the fact that each V_α^P is CCC.

This makes sure that when $\langle \mathcal{A}_\beta : \beta \in \alpha \rangle$ has been defined for a limit $\alpha \in \omega_1$ with $\mathcal{A}_\beta \in V_{\mathcal{T}(\beta)}^P$ and $\mathcal{T}(\beta) < \omega_1$ for each $\beta \in \alpha$ then there is a $\mathcal{T}(\alpha) \in \omega_1$ such that $\langle \mathcal{A}_\beta : \beta \in \alpha \rangle \in V_{\mathcal{T}(\alpha)}^P$, and in this case we may define

$$\mathcal{A}_\alpha = \bigcup \{ \mathcal{A}_\beta : \beta \in \alpha \} \text{ in } V_{\mathcal{T}(\alpha)}^P.$$

Having completed the induction, we set $\mathcal{A} = \bigcup \{ \mathcal{A}_\alpha : \alpha \in \omega_1 \}$ and claim that \mathcal{A} is as required, i.e. it refines $I_{\mathcal{A}}^+$.

Thus let $H \in I_{\mathcal{A}}^+$ and note first that there is a $K \in [H]_{\mathcal{A}}^\omega$ with $K \in I_{\mathcal{A}}^+$ as well. Indeed, if

$$\mathcal{H} = \{ A \in \mathcal{A} : |A \cap H| = \omega \}$$

is finite then K may be chosen as any element of $[H \setminus \bigcup \mathcal{H}]^\omega$. Otherwise, let $\{ A_n : n \in \omega \}$ be distinct members of \mathcal{H} , clearly then

$$K = \bigcup \{ A_n \cap H : n \in \omega \}$$

is as required.

Next, since P is CCC, there is some $\alpha \in \omega_1$ with $K \in V_\alpha^P$. Obviously, we have then $K \in V_\alpha^P \cap I_{\mathcal{A}_\alpha}^+$ as well. But then, by our construction, there is some $A \subset \mathcal{A}_{\alpha+1} \subset \mathcal{A}$ with $A \subset K \subset H$, and our proof is complete.

In [2] the following problem was raised: For what cardinals κ is there an almost disjoint family $\mathcal{A} \subset [\kappa]^\omega$ that refines $[\kappa]^{\omega_1}$?

Since, trivially, $[\kappa]^{\omega_1} \subset I_{\mathcal{A}}^+$, we immediately get that every saturated family has this property, and in our V^P a saturated family exists for each κ . In [4] it was shown that an almost disjoint $\mathcal{A} \subset [\kappa]^\omega$ refining $[\kappa]^{\omega_1}$ exists for $\kappa = 2^\omega$ in ZFC and for every $\kappa < \omega_{\omega_1}$ in L. In [1] it was shown that an almost disjoint $\mathcal{A} \subset [\kappa]^\omega$ refining $\{\chi \subset \kappa : \text{tip}(X) \neq \omega^2\}$ exists for $\kappa = (2^\omega)^{+\omega}$, $\omega \in \omega$, in ZFC. Several similar problems are also discussed in [1]. On the other hand it is still unknown whether a saturated $\mathcal{A} \subset [\kappa]^\omega$ exists in ZFC.

To conclude, we note that our notion of forcing P is CCC with $|P| = 2^\omega$, hence V^P has the same cardinal arithmetic as V , moreover P is "mild" and thus will not effect large cardinals. Thus the problem of producing a model in which there is no saturated almost disjoint family on some set X looks very hard.

References

- [1] B. BALCAR, J. DOČKÁLKOVÁ, P. SIMON: Almost disjoint families of countable sets, Proc. Coll. Math. Soc. J. Bolyai 37. Finite and infinite sets, Eger, 1981.
- [2] A. HAJNAL: Some results and problems in set theory, Acta Math. Acad. Sci. Hung. 11(1960), 277-298.
- [3] T. JECH: Set theory, Academic Press, 1978.
- [4] P. KOMJÁTH: Dense systems of almost-disjoint sets, Proc. Coll. Math. Soc. J. Bolyai 37. Finite and infinite sets, Eger, 1981.
- [5] K. KUNEN: Set theory, North Holland, 1980.
- [6] P. NYIKOS, J. PELANT, P. SIMON: private communication.
- [7] S.H. HECHLER: Generalizations of almost disjointness, c -sets, and the Baire number of \mathfrak{N} - \mathfrak{N} , Gen. Top. and its Appl. 8(1978), 93-110.

Mathematical Institute of Hungarian Academy of Sciences, Reáltanoda n.13-15, Budapest V, Hungary

(Oblatum 12.8. 1987)