

Jan Čerych

A characterization of essential sets of function algebras

Archivum Mathematicum, Vol. 40 (2004), No. 3, 229--232

Persistent URL: <http://dml.cz/dmlcz/107905>

Terms of use:

© Masaryk University, 2004

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A CHARACTERIZATION OF ESSENTIAL SETS OF FUNCTION ALGEBRAS

JAN ČERYCH

ABSTRACT. In the present note, we characterize the essential set E of a function algebra A defined on a compact Hausdorff space X in terms of local properties of functions in A at the points of E .

Let X be a compact Hausdorff topological space. Denote by $C(X)$ the commutative Banach algebra, consisting of all continuous complex-valued functions on X (with respect to usual point-wise algebraic operations) endowed with the sup-norm.

By a *function algebra* on X we mean any closed subalgebra of $C(X)$ which contains constant functions on X and which separates points of X .

Definition. A function algebra A on X is said to be a *maximal* one if it is a proper subset (i.e., a proper subalgebra) of $C(X)$ and has the following property: whenever B is a function algebra on X , $B \supset A$, then either $B = A$ or $B = C(X)$.

A being a function algebra on X , a closed subset $E \subset X$ is said to be an *essential set* of A if the following conditions are fulfilled:

- (1) A consists of *all* continuous prolongations of functions in the algebra of restrictions A/E (i.e., the algebra of all restrictions of functions in A from the set X to its subset E).
- (2) Whenever a closed subset F of X has the same property as E in (1), then $E \subset F$ (or, E is a unique minimal closed subset of X satisfying the condition (1)).

The notion “essential set” is due to Bear, who proved in [1] that any maximal algebra on X has an essential set.

Hoffman and Singer in [2] found an essential set of any, not necessarily maximal, function algebra on X .

2000 *Mathematics Subject Classification*: 46J10.

Key words and phrases: compact Hausdorff space X , the sup-norm algebra $C(X)$ of all complex-valued continuous functions on X , its closed subalgebras (called function algebras), measure orthogonal to a function algebra.

Received July 3, 2002.

Denote by $M(X)$ the space of all complex Borel regular measures on X , i.e., by the Riesz Representation Theorem, the dual space of $C(X)$.

The *annihilator* A^\perp of a function algebra A is defined to be the set of all measures $m \in M(X)$ such that $\int f dm = 0$ for any $f \in A$, or the set of all measures *orthogonal* to A . The dual space A' of A is then canonically isomorphic to the quotient space $M(X)/A^\perp$.

Now endow $M(X)$ with the weak-star topology: it is well known that $M(X)$ becomes a locally convex topological linear space with the dual space $C(X)$.

Definition. Let A be a function algebra on X . A (closed nonvoid) set $F \subset X$ is said to be a *peak set* (of A) if there exists a function $f \in A$ with the following properties:

- (1) $f(x) = 1$ for any $x \in F$;
- (2) $|f(y)| < 1$ for any $y \in X \setminus F$.

In this case we say that f *peaks* on F .

In [3], we have proved the following

Theorem 1. *Let A be a function algebra on X . Denote by E the closure of the union of all closed supports of measures in A^\perp . Then E is the essential set of A .*

Our aim here is to characterize the essential set E of a function algebra A in terms of local properties of functions in A at the points off E . More precisely, we shall prove the following

Theorem 2. *Let A be a function algebra on X . Denote by E its essential set. Let $x \in X$. Then $x \in X \setminus E$ if and only if there exists a closed neighbourhood V of x in X such that the following two conditions are fulfilled:*

- (3) $A/V = C(V)$, where A/V means the algebra of all restrictions of functions from A to the set V ;
- (4) V is an intersection of peak sets of A .

Proof. Let at first $x \in X \setminus E$.

Take as V such an closed neighbourhood which does not meet E .

Condition (3) follows immediately from the definition of the essential set.

For any $y \in X \setminus (E \cup V)$ let f_y^0 be a function defined on the set $H_y = E \cup \{y\} \cup V$ such that it is equal to 1 on V and to 0 on $E \cup \{y\}$. We can, by the classical Tietze Theorem, construct a function $\tilde{f}_y \in C(X)$ which is equal to f_y^0 on the set H_y . Finally, put $f_y = \min(1, \tilde{f}_y)$. Then $f_y \in C(X)$ and f is equal to 0 on E ; it follows from the definition of the essential set that $f_y \in A$.

Denote the set on which f_y peaks by F_y . Then $F_y \supset V$ and F_y does not meet $E \cup \{y\}$. It follows that

$$V = \bigcap_{y \in X \setminus (E \cup V)} F_y,$$

the condition (4).

Let, on the contrary, be V such closed neighborhood of x that the conditions (3), (4) are fulfilled. Let m is a measure on X such that $\text{spt } m$, its closed support, has nonvoid intersection with $\text{int } V$, the interior of V . We shall prove that m is not in A^\perp ; it will follow from Theorem 1 that $x \notin E$.

Let $f \in C(V)$ be such that

$$(5) \quad \text{spt } f \subset \text{int } V, \quad \int_V f \, dm \neq 0.$$

It follows from (3) that there exists a function $g \in A$ such that $g|_V = f$. It is $f = g = 0$ on the boundary of V and then the the sets

$$(6) \quad U_n \equiv V \cup \{y \in X; |g(y)| < \frac{1}{n}, n = 1, 2, \dots\}$$

containing V are open.

The set $X \setminus U_n$ is a compact one; the system S of all peak sets of A containing V is a system of compact sets whose intersection is V by (4). It follows that there is a finite subsystem F_1, F_2, \dots, F_k of S such that

$$(7) \quad V_n \equiv \bigcap_{j=1}^k F_j \subset U_n.$$

But the (nonvoid) intersection of peak sets is a peak set: if f_j peaks on F_j , then $\prod f_j$ peaks on $\cap F_j$. We have proved: there exists a sequence $V_n, n = 1, 2, \dots$ of peak sets of A such that

$$(8) \quad V \subset V_n \subset U_n, \quad n = 1, 2, \dots$$

It is easy to see that the intersection $W \equiv \bigcap_{n=1}^\infty V_n$ is a peak set of A : if $h_n \in A$ peaks on V_n , then the function

$$h \equiv \sum_{n=1}^\infty 2^{-n} h_n$$

peaks on W . It follows from (7) and (8) that

$$(9) \quad V \subset W \subset V \cup \{y \in X; g(y) = 0\}.$$

We have

$$\begin{aligned} h^n(y) &= 1 \text{ for } y \in W, \\ h^n(y) &\rightarrow 0 \text{ for } n \rightarrow \infty, y \in X \setminus V. \end{aligned}$$

It follows from (9) that

$$\begin{aligned} (g \cdot h^n)(y) &= g(y) \leq f(y) \text{ for } y \in V, \\ (g \cdot h^n)(y) &\rightarrow 0 \text{ for } n \rightarrow \infty, y \in X \setminus W. \end{aligned}$$

Since $|g \cdot h^n| = |g| \cdot |h|^n = |g|$, we have by the Lebesgue Dominated Convergence Theorem and by (9) and (5)

$$\int g \cdot h^n \, dm \rightarrow \int_W g \, dm = \int_V g \, dm = \int_V f \, dm \neq 0.$$

But $g \cdot h^n \in A$ for $n = 1, 2, \dots$ and then the measure m is not in A^\perp . \square

At first look at Theorem 1 and 2 it would appear that the condition (4) in Theorem 2 is superfluous and could be omitted. The next example shows that it is not the case.

Example. Let A be the *classical disk algebra*, i.e. the algebra of all functions continuous on the closed unit disk K in the complex plane which are holomorphic on the interior of K .

Let B be the restriction of A to the set

$$F \equiv \{z \in K; |z| = 1 \text{ or } z = 0\}.$$

Zero is an isolated point of F ; it follows that $B/\{0\} = C(\{0\})$.

Let μ be such a measure on F that for any $f \in C(F)$

$$\int f \, d\mu = \frac{1}{2\pi i} \int_{|z|=1} \frac{f(z) \, dz}{z} - f(0).$$

Then $\mu \in B^\perp$ by Cauchy Formula. There is $|\mu|(0) = 1$, so 0 is in the essential set of B , by Theorem 1.

REFERENCES

- [1] Bear, H. S., *Complex function algebras*, Trans. Amer. Math. Soc. **90** (1959), 383–393.
- [2] Hoffman, K., Singer, I. M., *Maximal algebras of continuous functions*, Acta Math. **103** (1960), 217–241.
- [3] Čerych, J., *On essential sets of function algebras in terms of their orthogonal measures*, Comment. Math. Univ. Carolin. **36**, 3 (1995), 471–474.

CHARLES UNIVERSITY IN PRAGUE, FACULTY OF MATHEMATICS AND PHYSICS
 SOKOLOVSKÁ 83, 186 00 PRAHA 8, CZECH REPUBLIC