

Gharagoz G. Hamedani

On periodic solutions of nonlinear second order ordinary differential equations

Časopis pro pěstování matematiky, Vol. 103 (1978), No. 4, 380--383

Persistent URL: <http://dml.cz/dmlcz/117995>

Terms of use:

© Institute of Mathematics AS CR, 1978

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

ON PERIODIC SOLUTIONS OF NONLINEAR
SECOND ORDER ORDINARY DIFFERENTIAL EQUATIONS

G. G. HAMEDANI, Tehran

(Received December 29, 1976)

In our previous paper ([2], Theorem 1) we established the existence of w -periodic solutions of the differential equation $x'' + Kx = F(t, x, x')$ for the case $K > 0$. In this note we prove an existence (and uniqueness; Corollary 2) theorem for this differential equation for $K \neq 0$. This theorem is stronger than Theorem 1 of [2] in the sense that there is no restriction on w (except that $[0, w] \subseteq [0, \pi/\sqrt{K}]$ for $K > 0$, and $[0, w] \subseteq [0, +\infty)$ for $K < 0$). Furthermore, its extension (which can be obtained with out difficulties) to a system of nonlinear second order differential equations provides a stronger theorem than Theorems 1 and 2 of [1].

Consider the scalar boundary value problem

$$(1) \quad x'' + f(t, x, x') = 0,$$

$$(2) \quad x(0) - x(w) = x'(0) - x'(w) = 0,$$

where f is a continuous real-valued function with domain $[0, w] \times R^2$.

Theorem 1. *Let there exist constants $K \neq 0$ and $C > 0$ such that*

$$(3) \quad M = \text{Max} \{ |Kx - f(t, x, x')| : t \in [0, w], |x| \leq C, \\ |x'| \leq (\sqrt{|K|}) C \} \leq |K| C.$$

Then in $[0, w] \subseteq [0, \pi/\sqrt{K}]$ if $K > 0$, and in $[0, w] \subseteq [0, +\infty)$ if $K < 0$, the problem (1), (2) has at least one solution $x(t)$ satisfying $|x(t)| \leq C$, $|x'(t)| \leq (\sqrt{|K|}) C$ for $0 \leq t \leq w$.

Proof. If $K > 0$, then problem (1), (2) is equivalent to the integral equation

$$(4) \quad x(t) = \int_0^w G(t, s) F(s, x(s), x'(s)) ds,$$

where $F(t, x, x') = Kx - f(t, x, x')$ and $G(t, s)$ is Green's function

$$(5) \quad G(t, s) = \begin{cases} \frac{1}{2\sqrt{K}} \cdot \frac{\cos(\sqrt{K})(\frac{1}{2}w + s - t)}{\sin(\sqrt{K})w/2} & \text{for } 0 \leq s \leq t \leq w \\ \frac{1}{2\sqrt{K}} \cdot \frac{\cos(\sqrt{K})(\frac{1}{2}w + t - s)}{\sin(\sqrt{K})w/2} & \text{for } 0 \leq t \leq s \leq w. \end{cases}$$

If $K < 0$, then (1), (2) is equivalent to (4) where

$$(6) \quad G(t, s) = \begin{cases} \frac{1}{2\sqrt{|K|}} \cdot \frac{\exp[-(\sqrt{|K|})(t-s)] \exp[(\sqrt{|K|})w] + \exp[(\sqrt{|K|})(t-s)]}{1 - \exp[(\sqrt{|K|})w]} & \text{for } s \leq t \\ \frac{1}{2\sqrt{|K|}} \cdot \frac{\exp[-(\sqrt{|K|})(s-t)] \exp[(\sqrt{|K|})w] + \exp[(\sqrt{|K|})(s-t)]}{1 - \exp[(\sqrt{|K|})w]} & \text{for } t \leq s. \end{cases}$$

Let $S = \{x \in C'[0, w] : |x(t)| \leq C, |x'(t)| \leq (\sqrt{|K|})C\}$ and define an operator U on S by

$$Ux(t) = \int_0^w G(t, s) F(s, x(s), x'(s)) ds.$$

From (3), it follows that

$$|Ux(t)| \leq M \int_0^w |G(t, s)| ds \leq \frac{M}{|K|} \leq C,$$

$$\left| \frac{d}{dt} Ux(t) \right| \leq M \int_0^w |G_t(t, s)| ds \leq \frac{M}{\sqrt{|K|}} \leq (\sqrt{|K|})C,$$

and hence U maps S continuously into itself. Therefore by Schauder's theorem (4) (and hence (1), (2)) has a solution with the desired properties.

Corollary 1. *If in addition to the hypotheses of Theorem 1, the function $f(t, x, x')$ is w -periodic in t and locally Lipschitzian with respect to (x, x') , then (1), (2) has a w -periodic solution.*

Corollary 2. *If in addition to the hypotheses of Theorem 1, the function $f(t, x, x')$ is w -periodic in t and if*

$$|F(t, x_1, x'_1) - F(t, x_2, x'_2)| \leq C_1 \left\{ |x_1 - x_2| + \frac{1}{\sqrt{|K|}} |x'_1 - x'_2| \right\}, \quad 0 \leq t \leq w$$

for $(x, x') \in \Omega = \{(x, x') : |x| \leq C, |x'| \leq (\sqrt{|K|}) C\}$, where $C_1 > 0$ is a constant such that

$$\frac{2C_1}{|K|} < 1,$$

then (1), (2) has a unique w -periodic solution.

Proof. If, for $x \in S$, we let

$$\|x\| = \text{Max} \left\{ |x(t)| + \frac{1}{\sqrt{|K|}} x'(t) : 0 \leq t \leq w \right\},$$

we can easily show that U is a contraction with respect to $\|\cdot\|$ on S .

Applications. Three applications of Theorem 1 for the case $K > 0$ can be found in ([2], pp. 73–75). We give below three applications for the case $K < 0$.

(A₁) Consider the equation

$$(7) \quad x'' + f(x) x'^n + ax = \mu p(t), \quad a < 0, \quad n \geq 2$$

where n is an integer, all coefficients are continuous, $f(x)$ is locally Lipschitzian in x , $0 \leq f(x) \leq b$ for all x , and $|\mu|$ sufficiently small. If $K < a/2$ and if $p(t)$ is periodic of period w , then (7) has a w -periodic solution.

Proof. The hypotheses of Corollary 1 are satisfied by choosing $C = |\mu|^{1/n}$ with $|\mu|$ sufficiently small.

(A₂) Consider the equation

$$(8) \quad x'' + a(t)x + b(t)f(x^2) = \mu p(t),$$

and let

- (i) $a(t), b(t), p(t)$ be continuous and $a(t)$ non-positive,
- (ii) $f(x)$ be locally Lipschitzian, non-negative, non-decreasing for $x \geq 0$ and for some $C > 0$

$$B \frac{f(C^2)}{C} + |\mu| \frac{D}{C} \leq -E$$

where

$$B = \text{Max}_{t \in [0, w]} |b(t)|, \quad D = \text{Max}_{t \in [0, w]} |p(t)|, \quad E = \text{Max}_{t \in [0, w]} a(t).$$

If $K \leq A = \text{Min}_{t \in [0, w]} a(t)$ and if $a(t), b(t), p(t)$ are periodic of period w then (8) has a w -periodic solution.

Proof. The hypotheses of Corollary 1 are satisfied by choosing C as in (ii).

(A₃) Consider the equation

$$(9) \quad x'' + x'(1 - x^2) - x = \mu p(t),$$

where $p(t)$ is continuous in t . If $K < -\frac{1}{2}$ and if $p(t)$ is periodic of period w then (9) has a w -periodic solution.

Proof. If $0 < C \leq 1$, $0 < \varepsilon < 1/\sqrt{|K|}$, and $|\mu| \leq (1 - \varepsilon(\sqrt{|K|})) C/B$, where $B = \text{Max}_{t \in [0, w]} |p(t)|$, then by Corollary 1 (9) has a w -periodic solution.

References

- [1] G. G. Hamedani: Periodic Boundary Value Problems for Nonlinear Second Order Vector Differential Equations. To appear in *Revue Roum. De Math. Pures Et App.*
- [2] B. Mehri and G. G. Hamedani: On the Existence of Periodic Solutions of Nonlinear Second Order Differential Equations, *SIAM J. Appl. Math.* Vol. 29, No. 1, July 1975.

Author's address: Arya-Mehr University of Technology, Tehran, Iran.