

Małgorzata Doman

Weak uniform rotundity of Musielak--Orlicz spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 32 (1991), No. 3, 441--446

Persistent URL: <http://dml.cz/dmlcz/118425>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1991

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Weak uniform rotundity of Musielak–Orlicz spaces

MALGORZATA DOMAN

Abstract. We give necessary and sufficient conditions for weak uniform rotundity of Musielak–Orlicz spaces L_φ with the Luxemburg norm. The result is a generalization of a theorem by Kamińska and Kurc.

Keywords: Musielak–Orlicz space, rotundity

Classification: 46B20, 46B25

Introduction.

Let T be a set, Σ a σ -algebra of subsets of T , μ a non-negative atomless σ -finite complete measure on Σ . A function $\varphi : R_+ \times T \rightarrow R_+$, where $R_+ = [0, +\infty)$, is said to be a Musielak–Orlicz function if $\varphi(0, t) = 0$ for μ -almost every $t \in T$, $\varphi(\cdot, t)$ is a convex function on R_+ for μ -almost every $t \in T$ and $\varphi(u, \cdot)$ is a Σ -measurable function on T for every $u \geq 0$. The complementary function to a function φ is defined by $\varphi^*(v, t) = \sup_{u > 0} (vu - \varphi(u, t))$ for $v \geq 0, t \in T$. We denote by M the set of all Σ -measurable functions $x : T \rightarrow R$. The functions which are different only on a null-set are considered as identical. The Musielak–Orlicz space L_φ is a subset of M such that $I_\varphi(\lambda x) = \int_T \varphi(\lambda|x(t)|, t) d\mu < +\infty$ for some $\lambda > 0$ dependent on x . The functionals $\|x\|_\varphi = \inf\{r > 0 : I_\varphi(\frac{x}{r}) \leq 1\}$ and $\|x\|_\varphi^0 = \sup\{\int_T x(t)y(t) d\mu : y \in L_{\varphi^*}, I_{\varphi^*}(y) \leq 1\}$ are norms in this space, called the Luxemburg and the Orlicz norm, respectively. We say that a function φ satisfies the condition Δ_α , for some $\alpha > 1$, if there are a constant $K_\alpha > 0$ and a function $h_\alpha : T \rightarrow R_+$, such that $\int_T h_\alpha(t) d\mu < +\infty$ and $\varphi(\alpha u, t) \leq K_\alpha \varphi(u, t) + h_\alpha(t)$ for almost every $t \in T$ and for $u \geq u_0$ (u_0 -some positive constant), when $\mu(T) < +\infty$, or for all $u \in R_+$, when $\mu(T) = +\infty$. Recall that a function φ is called strictly convex, if for all $u, v \in R_+, u \neq v, \alpha, \beta \in R_+ \setminus \{0\}, \alpha + \beta = 1$, we have $\varphi(\alpha u + \beta v, t) < \alpha \varphi(u, t) + \beta \varphi(v, t)$ outside of some null-set. For further details concerning Musielak–Orlicz spaces see [7].

We say that a Banach space $(X, \|\cdot\|)$ is weakly uniformly rotund (WUR), if for every $x^* \in X, x^* \neq 0$ and $\varepsilon > 0$ there exists $\delta(x^*, \varepsilon) > 0$, such that if $\|x\| = \|y\| = 1$ and $x^*(x - y) \geq \varepsilon$, then $\|\frac{x+y}{2}\| \leq 1 - \delta(x^*, \varepsilon)$ (cf. [1]). If for all $x, y \in X$ such that $\|x\| = \|y\| = 1$ we have $\|\frac{x+y}{2}\| < 1$, then we say that $(X, \|\cdot\|)$ is rotund.

The aim of this paper is to give necessary and sufficient conditions for WUR of Musielak–Orlicz spaces. The result is a generalization of a theorem by Kamińska and Kurc ([6, Theorem 2.8]).

I am greatly indebted to Professor Anna Kamińska for suggesting the problem discussed here to me. I wish to thank also Professor Henryk Hudzik for his help in preparing this paper

Results.

For the proof of the main theorem, we need some lemmas.

Lemma 1 (cf. [6]). *If an arbitrary Banach space contains an isomorphic copy of l_1 , then X is not WUR.*

Lemma 2. *If φ is a strictly convex Musielak–Orlicz function, then for every $\varepsilon > 0$ and every Σ -measurable functions $p, q : T \rightarrow (0, +\infty), p(t) < q(t)$ for μ -almost every $t \in T$, there exists a Σ -measurable function $r : T \rightarrow (0, 1)$ such that*

$$\varphi\left(\frac{u+v}{2}, t\right) \leq \frac{1-r(t)}{2}\{\varphi(u, t) + \varphi(v, t)\}$$

for μ -almost every $t \in T$ whenever $|u - v| \geq \varepsilon \max\{|u|, |v|\}$ and $\max\{|u|, |v|\} \in [p(t), q(t)]$.

The proof of this lemma is analogous to that of Lemma 1 in [5], so it is omitted here. □

Lemma 3. *Assume that φ is a Musielak–Orlicz function satisfying the Δ_2 -condition. Then for every $\alpha > 1$ and $\varepsilon > 0$, there is a set T_0 of measure 0, a constant $K_{\alpha,\varepsilon} > 0$ and a Σ -measurable function $h_{\alpha,\varepsilon} : T \rightarrow [0, +\infty)$ such that $\int_T h_{\alpha,\varepsilon}(t) d\mu \leq \varepsilon$ and $\varphi(\alpha u, t) \leq K_{\alpha,\varepsilon}\varphi(u, t) + h_{\alpha,\varepsilon}(t)$ for any $t \in T \setminus T_0$ and any $u \in R$.*

The proof for $\alpha = 2$ is given in [4]. The proof for an arbitrary $\alpha > 1$ can proceed in the same way, if we notice that φ satisfies the Δ_2 -condition if and only if it satisfies the Δ_α -condition for every $\alpha > 1$. □

Lemma 4 (cf. [4]). *Let φ be a Musielak–Orlicz function that satisfies the Δ_2 -condition. Then*

- (i) *there is a function $\beta : (0, 1) \rightarrow (0, 1)$ such that $\|x\|_\varphi \leq 1 - \beta(\varepsilon)$ whenever $I_\varphi(x) \leq 1 - \varepsilon$.*
- (ii) *$\|x\|_\varphi = 1$ if and only if $I_\varphi(x) = 1$.*

Lemma 5. *Assume that φ is a Musielak–Orlicz function vanishing only at 0 and that φ and φ^* satisfy the Δ_2 -condition. Let $x^* \in (L_\varphi)^*$ be regular and nontrivial (i.e. there exists $z \in L_{\varphi^*}, z \neq 0$ such that $x^*(x) = \int_T x(t)z(t) d\mu$ for every $x \in L_\varphi$). Let $(B_n)_{n=1}^\infty$ be an increasing sequence of sets with finite and positive measures such that $\bigcup_n B_n = \text{supp } z$. Denote $C_n = \{t \in T : \frac{1}{n} \leq |z(t)| \leq n\}$ and put $D_n = C_n \cap B_n$. Then $(D_n)_{n=1}^\infty$ is increasing, $\bigcup_n D_n = \text{supp } z$ and*

$$\int_{D_n} y(t)z(t) d\mu \rightarrow \int_T y(t)z(t) d\mu$$

uniformly with respect to y in every bounded set in L_φ .

PROOF: In virtue of B. Levi theorem and the Δ_2 -condition for φ^* , we have $\|z - z_n\|_{\varphi^*}^0 \rightarrow 0$ as $n \rightarrow +\infty$, where $z_n = z\chi_{D_n}$. Then

$$\begin{aligned} 0 &\leq \left| \int_T y(t)z(t) d\mu - \int_{D_m} y(t)z(t) d\mu \right| = \\ &= \left| \int_T y(t)z(t) d\mu - \int_T y(t)z_m(t) d\mu \right| \leq \\ &\leq \|y\|_{\varphi} \|z - z_m\|_{\varphi^*}^0 \leq C \|z - z_m\|_{\varphi^*}^0. \end{aligned}$$

Hence the desired result follows. \square

The next two lemmas are analogs of Lemma 2.5 and Lemma 2.6 of [6].

Lemma 6. Let $\mu(T) < +\infty$ and φ be a Musielak–Orlicz function such that for every $t \in T$ $\frac{\varphi(u,t)}{u} \rightarrow +\infty$ as $u \rightarrow +\infty$. Then for every $\varepsilon > 0$, there exist Σ -measurable functions $p, q : T \rightarrow (0, +\infty)$ such that for every $x, y \in L_{\varphi}$ satisfying $I_{\varphi}(x) = I_{\varphi}(y) = 1$ and $\int_T |x(t) - y(t)| d\mu \geq \varepsilon$, we have $\int_A |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{4}$ whenever

$$A = \{t \in T : p(t) \leq \max(|x(t)|, |y(t)|) \leq q(t)\}.$$

PROOF: Define for any $n \in N$ $p_n(t) = \inf\{u > 0 : \frac{\varphi(u,t)}{u} \geq n\}$. Then p_n is a Σ -measurable function and $\varphi(u,t) \geq nu$ for every $u \geq p_n(t)$. Define $A_n = \{t \in T : |x(t)| \leq p_n(t)\}$, $A_n^1 = \{t \in T : |y(t)| \leq p_n(t)\}$. We have

$$\int_{T \setminus A_n} |x(t)| d\mu \leq \frac{1}{n} \int_{T \setminus A_n} \varphi(|x(t)|, t) d\mu \leq \frac{1}{n}.$$

In the same way, we can obtain $\int_{T \setminus A_n^1} |y(t)| d\mu \leq \frac{1}{n}$. Moreover,

$$\begin{aligned} \int_{T \setminus A_n} |y(t)| d\mu &= \int_{(T \setminus A_n) \cap (T \setminus A_n^1)} |y(t)| d\mu + \int_{A_n^1 \setminus A_n} |y(t)| d\mu \leq \\ &\leq \int_{T \setminus A_n^1} |y(t)| d\mu + \int_{T \setminus A_n} |x(t)| d\mu \leq \frac{2}{n}. \end{aligned}$$

Similarly $\int_{T \setminus A_n^1} |x(t)| d\mu \leq \frac{2}{n}$. Hence $\int_{T \setminus (A_n \cap A_n^1)} |x(t) - y(t)| d\mu \leq \int_{T \setminus A_n} |x(t)| d\mu + \int_{T \setminus A_n^1} |y(t)| d\mu + \int_{T \setminus A_n^1} |x(t)| d\mu + \int_{T \setminus A_n} |y(t)| d\mu \leq \frac{6}{n}$. Since $\int_T |x(t) - y(t)| d\mu \geq \varepsilon$ by the assumptions, we have $\int_{A_n \cap A_n^1} |x(t) - y(t)| d\mu \geq \varepsilon - \frac{6}{n} \geq \frac{\varepsilon}{2}$ if n is such that $\frac{6}{n} \leq \frac{\varepsilon}{2}$. Define $A_n^2 = \{t \in T : \frac{\varepsilon}{8\mu(T)} \leq \max(|x(t)|, |y(t)|)\}$. If $t \notin A_n^2$, then $|x(t)| < \frac{\varepsilon}{8\mu(T)}$ and $|y(t)| < \frac{\varepsilon}{8\mu(T)}$. Therefore $\int_{(A_n \cap A_n^1) \setminus A_n^2} |x(t) - y(t)| d\mu \leq \frac{\varepsilon}{8\mu(T)} \mu(T \setminus A_n^2) + \frac{\varepsilon}{8\mu(T)} \mu(T \setminus A_n^2) \leq \frac{\varepsilon}{4}$. Thus $\int_{A_n \cap A_n^1 \cap A_n^2} |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{2} - \frac{\varepsilon}{4} = \frac{\varepsilon}{4}$. Putting $A = A_n \cap A_n^1 \cap A_n^2$, $p(t) = \frac{\varepsilon}{8\mu(T)}$ and $q(t) = p_n(t)$, we get the desired inequality. \square

Lemma 7. *Let φ be a Musielak–Orlicz function satisfying the Δ_2 -condition and let $B \in \Sigma, \varepsilon > 0$ and $\sigma \in (0, 1)$ be such that $I_\varphi((x - y)\chi_B) \geq \varepsilon$ and $I_\varphi(\frac{x+y}{2}) \leq 1 - \frac{\sigma}{2}(I_\varphi(x\chi_B) + I_\varphi(y\chi_B))$, where x, y are arbitrary measurable functions with $I_\varphi(x) = I_\varphi(y) = 1$. Then there exists a constant $q \in (0, 1)$, such that $I_\varphi(\frac{x+y}{2}) \leq 1 - q$.*

PROOF: Let $K = K_{2, \frac{\varepsilon}{2}}$, where $K_{2, \frac{\varepsilon}{2}}$ is the constant from Lemma 3. Then

$$\varepsilon \leq I_\varphi((x - y)\chi_B) \leq \frac{K}{2}(I_\varphi(x\chi_B) + I_\varphi(y\chi_B)) + \frac{\varepsilon}{2}.$$

Hence $I_\varphi(x\chi_B) + I_\varphi(y\chi_B) \geq \frac{\varepsilon}{2} \cdot \frac{2}{K} = \frac{\varepsilon}{K}$. Therefore $I_\varphi(\frac{x+y}{2}) \leq 1 - \frac{\sigma\varepsilon}{2K}$, and it suffices to put $q = \frac{\sigma\varepsilon}{2K}$ □

Theorem 1. *A Musielak–Orlicz space L_φ is WUR if and only if*

- (i) φ is strictly convex,
- (ii) φ satisfies the Δ_2 -condition,
- (iii) φ^* satisfies the Δ_2 -condition.

PROOF: Sufficiency. Assume that the conditions (i), (ii), (iii) are satisfied. Let $x, y \in L_\varphi, \|x\|_\varphi = \|y\|_\varphi = 1, x^* \in (L_\varphi)^*$ and $x^*(x - y) \geq \varepsilon$, where $\varepsilon \in (0, 1)$. In virtue of the representation of x^* , we have $\int_T(x(t) - y(t))z(t) d\mu \geq \varepsilon$ for some $z \in L_{\varphi^*}$. Define z_n as in the proof of Lemma 5. Then in view of this lemma, there is $n_0 \in N$ (n_0 independent of x and y) such that $\int_T(x(t) - y(t))z_{n_0}(t) d\mu \geq \frac{\varepsilon}{2}$. Since $|z_{n_0}(t)| < n_0$, denoting $T_0 = \text{supp } z_{n_0}$, we get $\int_{T_0} |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{2n_0}$. Since, according to Lemma 2.4 of [6], (iii) implies $\varphi(u, t)/u \rightarrow +\infty$ when $u \rightarrow +\infty$ for every $t \in T$, it follows from Lemma 6 that there exist two Σ -measurable functions $p, q : T_0 \rightarrow (0, +\infty)$, such that denoting

$$A = \{t \in T_0 : p(t) \leq \max(|x(t), y(t)|) \leq q(t)\}, \text{ we have}$$

$$\int_A |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{8n_0}.$$

Define $B = \{t \in A : |x(t) - y(t)| \geq \frac{\varepsilon}{8n_0K} \max(|x(t)|, |y(t)|)\}$, where $K = K_{\alpha, \frac{1}{2}}$ is the constant from Lemma 3 corresponding to $\alpha = \max\{\frac{64n_0}{\varepsilon} \|\chi_{T_0}\|_{\varphi^*}, 1\}$. In virtue of Lemma 2 there is a function $r : B \rightarrow (0, 1)$ such that

$$\varphi\left(\frac{|x(t) + y(t)|}{2}, t\right) \leq \frac{1 - r(t)}{2} \{\varphi(|x(t)|, t) + \varphi(|y(t)|, t)\}.$$

Define $B_m = \{t \in B : r(t) \geq \frac{1}{m}\}$. We have $B_m \nearrow$ and $\bigcup_{n=1}^\infty B_m = B$. Thus, defining $C_m = (A \setminus B) \cup B_m$, we obtain the increasing sequence of sets such that $\bigcup_{n=1}^\infty C_n = A$. By Lemma 5 there is $s \in N$ (s independent of x and y) such that

$$\int_{C_s} |x(t) - y(t)| d\mu \geq \int_A |x(t) - y(t)| d\mu - \frac{1}{4} \cdot \frac{\varepsilon}{8n_0}.$$

i.e.

$$(1) \quad \int_{C_s} |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{32n_0}.$$

For $t \in B_s$, we have

$$\varphi\left(\frac{|x(t) + y(t)|}{2}, t\right) \leq \frac{1 - \frac{1}{s}}{2} \{\varphi(|x(t)|, t) + \varphi(|y(t)|, t)\}.$$

Hence, using the convexity of φ and the fact that $I_\varphi(x) = I_\varphi(y) = 1$, we get

$$(2) \quad I_\varphi\left(\frac{x+y}{2}\right) \leq 1 - \frac{1}{2s} \{I_\varphi(x)\chi_{B_s} + I_\varphi(y)\chi_{B_s}\}.$$

If $t \in A \setminus B$, then

$$|x(t) - y(t)| < \frac{\varepsilon}{8n_0K} \max(|x(t)|, |y(t)|).$$

Hence

$$(3) \quad I_\varphi((x-y)\chi_{A \setminus B}) \leq \frac{\varepsilon}{8n_0K} \{I_\varphi(x\chi_{A \setminus B}) + I_\varphi(y\chi_{A \setminus B})\} \leq \frac{\varepsilon}{4n_0K}.$$

Applying the inequality (1) and the Hölder inequality, we get

$$2\|(x-y)\chi_{C_s}\|_\varphi \|\chi_{T_0}\|_{\varphi^*} \geq \int_{C_s} |x(t) - y(t)| d\mu \geq \frac{\varepsilon}{32n_0},$$

i.e.

$$\frac{64n_0}{\varepsilon} \|\chi_{T_0}\|_{\varphi^*} \|(x-y)\chi_{C_s}\|_\varphi \geq 1,$$

hence denoting $\alpha_1 = \frac{64n_0}{\varepsilon} \|\chi_{T_0}\|_{\varphi^*}$, we have $\alpha_1 \leq \alpha$, and

$$1 \leq I_\varphi(\alpha(x-y)\chi_{C_s}) \leq KI_\varphi((x-y)\chi_{C_s}) + \frac{1}{2}.$$

Thus

$$I_\varphi((x-y)\chi_{C_s}) \geq \frac{1}{2K}.$$

Combining this with the inequality (3), we get

$$I_\varphi((x-y)\chi_{B_s}) \geq I_\varphi((x-y)\chi_{C_s}) - I_\varphi((x-y)\chi_{A \setminus B}) \geq \frac{1}{2K} - \frac{\varepsilon}{4n_0K} = \beta.$$

Applying Lemma 7, the inequality (2) and the last inequality, we get

$$I_\varphi\left(\frac{x+y}{2}\right) \leq 1 - q.$$

Now, in view of Lemma 4, we have

$$\left\| \frac{x+y}{2} \right\|_{\varphi} \leq 1 - \beta(q),$$

where $\beta(q) \in (0, 1)$, and depends only on x^* , ε and φ .

Necessity. If φ does not satisfy the condition (i) or the condition (ii), then L_{φ} is not rotund (cf. [5]). Since WUR implies rotundity, L_{φ} is not WUR as well. Assume now that φ satisfies the condition (i) and it does not satisfy the condition (iii). Then $(L_{\varphi})^* = L_{\varphi^*}$, where L_{φ^*} is equipped with the Orlicz norm. Since φ^* does not satisfy the Δ_2 -condition, L_{φ^*} contains an isomorphic copy of l_{∞} . Hence it follows that L_{φ} contains an isomorphic copy of l_1 . Therefore, in view of Lemma 1, L_{φ} is not WUR. The proof is finished. \square

Theorem 1.2 of [3] and Theorem 1.2 of [2] imply the following version of our result.

Theorem 2. *A Musielak–Orlicz space L_{φ} is WUR if and only if it is rotund and reflexive.*

REFERENCES

- [1] Diestel J., *Geometry of Banach spaces—selected topics*, Springer Lecture Notes in Mathematics, vol. 485, 1983.
- [2] Hudzik H., *On some equivalent conditions in Musielak–Orlicz spaces*, Comment. Math. **24** (1984), 57–64.
- [3] Hudzik H., *Strict convexity of Musielak–Orlicz spaces with Luxemburg’s norm*, Bull. Acad. Polon. Sci., Sér. Sci. Math., Astronom. et Phys. **29** (1981), 235–247.
- [4] Hudzik H., *Uniform convexity of Musielak–Orlicz spaces with Luxemburg’s norm*, Comment. Math. **23** (1983), 21–32.
- [5] Kamińska A., *On some convexity properties of Musielak–Orlicz spaces*, Supplemento ai Rendiconti del Circolo Matematico di Palermo, The 12th Winter School on Abstract Analysis, Srní 1984.
- [6] Kamińska A., *Weak uniform rotundity in Orlicz spaces*, Comment. Math. Univ. Carolinae **27** (1986), 651–664.
- [7] Musielak J., *Orlicz spaces and modular spaces*, Springer Lecture Notes in Mathematics, vol. 1034, 1983.

ACADEMY OF ECONOMICS, DEPARTMENT OF MATHEMATICS, AL. NIEPODLEGŁOŚCI 10,
60–967 POZNAŃ, POLAND

(Received November 28, 1990, revised February 26, 1991)