

Charles W. Swartz

A uniform boundedness principle of Pták

Commentationes Mathematicae Universitatis Carolinae, Vol. 34 (1993), No. 1, 149--151

Persistent URL: <http://dml.cz/dmlcz/118564>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1993

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A uniform boundedness principle of Pták

CHARLES SWARTZ

Abstract. The Antosik-Mikusinski Matrix Theorem is used to give an extension of a uniform boundedness principle due to V. Pták to certain metric linear spaces. An application to bilinear operators is given.

Keywords: uniform boundedness, linear operator, bilinear operator

Classification: 46A99

In [P] V. Pták used the classical sliding hump technique to give an extension of the classical uniform boundedness principle for pointwise bounded families of continuous linear operators on Banach spaces and in [NP] used an abstract sliding hump construction to give a further extension. This generalization of the uniform boundedness principle was used to establish automatic continuity results. A similar generalized uniform boundedness principle was established earlier by Lorentz and MacPhail to give a generalization of the Silvermann-Toeplitz Theorem from summability ([LM]). The Antosik-Mikusinski Matrix Theorem ([AS, 2.2]) can also be considered to be an abstraction of the sliding hump technique and has been used to treat a number of topics in classical functional analysis and measure theory including the uniform boundedness principle (see [AS], especially §4 for the uniform boundedness principle). In this note we show that the Antosik-Mikusinski Matrix Theorem can be used to extend V. Pták's version of the uniform boundedness principle to metric \mathcal{K} -spaces. As an application we use the extended form of the uniform boundedness principle to generalize a result of Pták on bilinear operators.

We fix the notation to be used. Throughout the sequel X will denote a metric linear space and Y a normed linear space. A sequence $\{x_k\}$ is \mathcal{K} convergent if every subsequence of $\{x_k\}$ has a further subsequence $\{x_{n_k}\}$ such that the subseries $\sum x_{n_k}$ converges in X . A \mathcal{K} convergent sequence converges to 0, but a sequence can converge to 0 and not be \mathcal{K} convergent ([AS, 3.3]). A space in which every sequence which converges to 0 is \mathcal{K} convergent is called a \mathcal{K} -space. A complete metric linear space is a \mathcal{K} -space ([AS, 3.2]), but there are (normed) \mathcal{K} -spaces which are not complete ([K]). The notion of a \mathcal{K} -space is a weakened form of completeness which has proven to be a good substitute for completeness in several topics in functional analysis ([AS]).

We first establish a special case of Pták's uniform boundedness principle for metric \mathcal{K} -spaces. This result was established for Banach spaces by Lorentz and MacPhail [LM]; see also [M, 4.1].

Theorem 1. *Let X be a metric \mathcal{K} -space. For each $k \in \mathbb{N}$ let $T_k : X \rightarrow Y$ be linear and let M_k be a closed subspace of X such that T_k is continuous on M_k and $M_k \supset M_{k+1}$. If $\{T_k\}$ is pointwise bounded on X , then there exists p such that $\{T_k : k \in \mathbb{N}\}$ is equicontinuous on M_p .*

PROOF: Let $\{U_k\}$ be a neighborhood base at 0 in X with $U_k \supset U_{k+1}$, $\bigcap_{k=1}^{\infty} U_k = \{0\}$. If the conclusion fails, for each k there exist n_k and $x_k \in U_k \cap M_k$ such that $\|T_{n_k}x_k\| > k$, where we may assume $n_{k+1} > n_k$. For notation convenience assume that $n_k = k$. Consider the matrix $M = [(1/i)T_i x_j]$. Since $\{T_i\}$ is pointwise bounded, the columns of M converge to 0. Since $x_j \rightarrow 0$ in X , a \mathcal{K} -space, given any increasing sequence of positive integers $\{p_j\}$ there is a subsequence $\{q_j\}$ of $\{p_j\}$ such that $\sum_{j=1}^{\infty} x_{q_j} = x \in X$. For each fixed i , $\sum_{q_j \geq i} x_{q_j} \in M_i$ since M_i is closed and $M_i \supset M_{k+1}$ so that

$$\sum_{j=1}^{\infty} \frac{1}{i} T_i x_{q_j} = \frac{1}{i} T_i \left(\sum_{q_j < i} x_{q_j} \right) + \frac{1}{i} T_i \left(\sum_{q_j \geq i} x_{q_j} \right) = \frac{1}{i} T_i x$$

since T_i is continuous on M_i . Therefore, since $\{T_i\}$ is pointwise bounded,

$$\lim_i \sum_{j=1}^{\infty} \frac{1}{i} T_i x_{q_j} = \lim_i \frac{1}{i} T_i x = 0.$$

Hence, M is a \mathcal{K} -matrix ([AS, § 2]) and the diagonal of M converges to 0 by the Antosik-Mikusinski Matrix Theorem (AS, 2.2). This contradicts $\|T_i x_i\| > 1$ above. \square

This result was obtained in [LM] for Banach spaces by a sliding hump technique. It was obtained for the case when X is a Banach space and M_k is the (closed) kernel of T_k by Pták in [P]. It is shown in 2.1.3 of [PB] that the assumption that M_k is a closed kernel cannot be replaced with the assumption that M_k is a Baire space. It is also shown in 2.7. of [PB] that the assumption that Y is a normed space cannot be replaced with Y is a metric linear space.

From Theorem 1 we can easily obtain the general form of the uniform boundedness principle given in [NP].

Theorem 2. *Let X be a metric \mathcal{K} -space. For each $a \in A$ let $T_a : X \rightarrow Y$ be linear and M_a be a closed linear subspace of X such that T_a is continuous on M_a . If $\mathcal{T} = \{T_a : a \in A\}$ is pointwise bounded on X , then there exists a finite subset $F \subset A$ such that \mathcal{T} is equicontinuous on $\bigcap_{a \in F} M_a$.*

PROOF: Let $\{U_k\}$ be as in the proof of Theorem 1. Suppose the conclusion fails. Pick $a_1 \in A$. There exist $a_2 \in A$, $x_2 \in M_{a_1} \cap U_1$ such that $\|T_{a_2}x_2\| > 1$. There exist $a_2 \in A$, $x_3 \in M_{a_1} \cap M_{a_2} \cap U_2$ such that $\|T_{a_3}x_3\| > 1$. Continuing this construction produces sequences $\{a_k\} \subset A$, $x_k \in M_{a_1} \cap \dots \cap M_{a_k} \cap U_k$ such that $\|T_{a_k}x_k\| > 1$. Since T_{a_k} is continuous on $M_{a_1} \cap \dots \cap M_{a_k}$, Theorem 1 implies that $\{T_{a_j} : j \in \mathbb{N}\}$ is equicontinuous on some $M_{a_1} \cap \dots \cap M_{a_p}$. Since $x_j \rightarrow 0$, this is impossible by the construction. \square

This result was extended to Fréchet spaces X in 4.9.15 of [PB]. It was also shown in 4.9.16 of [PB] that X is a Banach space cannot be replaced in Theorem 1 by X is normed Baire even if each M_k is a Baire space. Since a metric \mathcal{K} -space is a Baire space but need not be complete, Theorem 2 gives an improvement of 4.9.15 of [PB]. The methods employed above are much simpler than those employed in [PB].

As an application of Theorem 2 we give an extension of Theorem 3.2 of [P] to families of bilinear maps. Let Z be a normed linear space. If $b : X \times Y \rightarrow Z$ is bilinear, for $x \in X$ ($y \in Y$) we let $b(x, \cdot) : Y \rightarrow Z$ ($b(\cdot, y) : X \rightarrow Z$) be the linear map $b(x, \cdot)(y) = b(x, y)$ ($b(\cdot, y)(x) = b(x, y)$). If $\mathcal{B} = \{b_a : a \in A\}$ is a family of bilinear maps from $X \times Y \rightarrow Z$, \mathcal{B} is said to be right equicontinuous if for each $x \in X$ the family $\{b_a(x, \cdot) : a \in A\}$ is an equicontinuous family of linear maps ([AS, §6]).

Theorem 3. *Let X be a metric \mathcal{K} -space and $\mathcal{B} = \{b_a : a \in A\}$ a family of bilinear maps from $X \times Y$ to Z . Assume*

- (1) \mathcal{B} is right equicontinuous,
- (2) for each $a \in A$ $y \in Y$ there is a closed subspace $M(a, y)$ of X such that $b_a(\cdot, y)$ is continuous on $M(a, y)$.

Then there exist $y_1, \dots, y_k \in Y$, $a_1, \dots, a_k \in A$ such that \mathcal{B} is equicontinuous on $M \times Y$ where $M = \bigcap_{j=1}^k M(a_j, y_j)$.

PROOF: Set $\mathcal{T} = \{b_a(\cdot, y) : a \in A, \|y\| \leq 1\}$. Since \mathcal{B} is right equicontinuous, \mathcal{T} is a pointwise bounded family of linear maps from X to Z . By Theorem 2 there exist $a_1, \dots, a_k \in A$, $y_1, \dots, y_k \in Y$ with $\|y_j\| \leq 1$ such that \mathcal{T} is equicontinuous on $M = \bigcap_{j=1}^k M(a_j, y_j)$. Therefore, given $\varepsilon > 0$ there exists a neighborhood U of 0 in X such that $\|b_a(x, y)\| < \varepsilon$ for $x \in U \cap M$, $\|y\| \leq 1$. Hence, \mathcal{B} is equicontinuous on $M \times Y$. \square

The case when \mathcal{B} consists of a single bilinear map is just Theorem 3.2 of [P]. Theorem 3 gives a generalization of the uniform boundedness principle for bilinear maps given in 6.16 of [AS].

REFERENCES

- [AS] Antosik P., Swartz C., *Matrix Methods in Analysis*, Springer-Verlag, Heidelberg, 1985.
 [K] Klis C., *An example of noncomplete normed K -spaces*, Bull. Acad. Polon. Sci. **26** (1978), 415–420.
 [LM] Lorentz G., MacPhail M., *Unbounded operators and a theorem of A. Robinson*, Trans. Royal Soc. Canada **46** (1952), 33–37.
 [M] Maddox I., *Infinite Matrices of Operators*, Springer-Verlag, Heidelberg, 1980.
 [NP] Neumann M., Pták V., *Automatic continuity, local type and casuality*, Studia Math. **82** (1985), 61–90.
 [P] Pták V., *A uniform boundedness theorem and mappings into spaces of operators*, Studia Math. **31** (1968), 425–431.
 [PB] Perez Carreras P., Bonet J., *Barrelled Locally Convex Spaces*, North Holland, Amsterdam, 1987.

DEPARTMENT OF MATHEMATICS, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NM 88003, USA

(Received January 22, 1992)