

Boris Lavrič

The nil radical of an Archimedean partially ordered ring with positive squares

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 35 (1994), No. 2, 231--238

Persistent URL: <http://dml.cz/dmlcz/118661>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## The nil radical of an Archimedean partially ordered ring with positive squares

BORIS LAVRIČ

*Abstract.* Let  $R$  be an Archimedean partially ordered ring in which the square of every element is positive, and  $N(R)$  the set of all nilpotent elements of  $R$ . It is shown that  $N(R)$  is the unique nil radical of  $R$ , and that  $N(R)$  is locally nilpotent and even nilpotent with exponent at most 3 when  $R$  is 2-torsion-free.  $R$  is without non-zero nilpotents if and only if it is 2-torsion-free and has zero annihilator. The results are applied on partially ordered rings in which every element  $a$  is expressed as  $a = a_1 - a_2$  with positive  $a_1, a_2$  satisfying  $a_1 a_2 = a_2 a_1 = 0$ .

*Keywords:* partially ordered ring, Archimedean, nil radical, nilpotent

*Classification:* 06F25, 16N40

### 0. Introduction

In this paper we give some descriptions of the set  $N(R)$  of all nilpotent elements of an Archimedean partially ordered ring  $R$  in which the square of every element is positive. It is shown among other things that  $N(R)$  is the unique nil radical of  $R$ , and that  $N(R)$  is locally nilpotent and even nilpotent with  $N(R)^3 = \{0\}$  when  $R$  is 2-torsion-free. Furthermore, we extend on  $R$  some results of Diem [3] and of Bernau and Huijsmans [1], who have investigated the properties of  $N(R)$  for a lattice-ordered  $R$ . In particular, we prove that the index of nilpotency of each  $a \in N(R)$  does not exceed 4 (3 when  $R$  is 2-torsion-free), and that  $R$  is without non-zero nilpotents if and only if it is 2-torsion-free and has zero annihilator  $\text{ann}(R)$ . Also an application on partially ordered rings in which every element  $a$  is expressed as  $a = a_1 - a_2$  with positive  $a_1, a_2$  satisfying  $a_1 a_2 = a_2 a_1 = 0$  is given at the end of paper.

For the theory of rings and nil radicals we refer the reader to [4], [7] and [9]. We only recall that an ideal  $I$  of a ring  $R$  is called a *nil radical*, if every element of  $I$  is nilpotent and if  $A/I$  does not contain non-zero nilpotent ideals.

For the theory of partially ordered rings we refer the reader to [5]. We briefly review some standard terminology. A ring  $R$  is said to be a *partially ordered ring* if there is a partial ordering  $\leq$  on  $R$  which is compatible with the algebraic structure of  $R$ . The positive cone  $R^+ = \{a \in R : 0 \leq a\}$  of a partially ordered ring  $R$  is closed for the addition and the multiplication, and determines on  $R$  the ordering  $\leq$  by  $a \leq b$  if and only if  $b - a \in R^+$ . A partially ordered ring  $R$  is said to be *Archimedean* if  $a = 0$  whenever  $0 \leq b + na$  for some  $b \in R$  and for

all integers  $n$ . A partially ordered ring  $R$  in which the square of every element is positive will be called a *ps-ring*.

### 1. The nil radical

Throughout this section  $R$  denotes an associative Archimedean *ps-ring*,  $N(R)$  the set of all nilpotent elements of  $R$ , and  $N_k(R) = \{a \in R : a^k = 0\}$  the set of all elements  $a \in N(R)$  with index of nilpotency not greater than  $k$ . An important role will play also the set  $T_2(R) = \{a \in R : 2a = 0\}$  of all elements  $a \in R$  with torsion 2.

The following description of  $N_2(R)$  will be useful in our further work.

**Lemma 1.1.** *Let  $R$  be an Archimedean *ps-ring*. Then*

- (1)  $N_2(R) = \{a \in R : ab + ba = 0 \text{ for all } b \in R\}$ .
- (2)  $abc = bac = bca \in T_2(R)$  for all  $a \in N_2(R)$ ,  $b, c \in R$ .

PROOF: If  $a \in N_2(R)$ ,  $b \in R$ , then

$$0 \leq (b + na)^2 = b^2 + n(ab + ba)$$

holds for all  $n \in \mathbb{Z}$ , hence  $ab + ba = 0$ , since  $R$  is Archimedean. Conversely, if  $a \in R$  satisfies  $ab + ba = 0$  for all  $b \in R$ , then  $2a^2 = 0$ . Since  $R$  has positive squares, this implies  $a \in N_2(R)$ , and (1) follows.

To prove (2) let  $a \in N_2(R)$  and  $b, c \in R$ . Then use three times successively (1) to get

$$a(bc) = -(bc)a = -b(ca) = b(ac) = (ba)c = -(ab)c,$$

which implies (2), so the proof is complete.  $\square$

**Corollary 1.2.** *Let  $m$  be an arbitrary positive integer. Then the left annihilator  $ann_l(R^m)$  and the right annihilator  $ann_r(R^m)$  of  $R^m$  coincide, so*

$$ann_l(R^m) = ann_r(R^m) = ann(R^m).$$

PROOF: If  $a \in N_2(R)$ ,  $b \in R$ , then by Lemma 1.1  $ab = 0$  if and only if  $ba = 0$ . Since  $ann_l(R)$  and  $ann_r(R)$  are contained in  $N_2(R)$ , this implies that  $ann_l(R) = ann_r(R)$ . We now proceed by induction. Suppose that  $ann_l(R^k) = ann_r(R^k)$  for each  $k \leq m$ . Then  $a \in ann_l(R^{m+1})$  is equivalent to  $aR \subset ann_l(R^m) = ann_r(R^m)$ , which holds if and only if  $R^m a \subset ann_l(R) = ann_r(R)$ . Since the latter is equivalent to  $a \in ann_r(R^{m+1})$ , the proof is complete.  $\square$

Diem has proved ([3, Theorem 3.9. (ii)]) that the index of a positive nilpotent element of an Archimedean lattice-ordered ring with positive squares does not exceed 3. Our next result shows that this is true for every Archimedean *ps-ring*.

**Proposition 1.3.** *Let  $R$  be an Archimedean ps-ring. Then*

- (1)  $N(R) = N_4(R) = \{a \in R : 2a^3 = 0\}$ .
- (2)  $N(R) \cap R^+ \subset N_3(R)$ .

PROOF: If  $a \in R$  satisfies  $a^{2m} = 0$  for some natural  $m > 2$ , then  $a^{4m-6} = 0$ , hence

$$0 \leq (a + na^{2m-3})^2 = a^2 + n(2a^{2m-2})$$

holds for all  $n \in \mathbb{Z}$ . Since  $R$  is Archimedean and has positive squares, this implies that  $a^{2m-2} = 0$ . It follows that  $N(R) = N_4(R)$ .

Let now  $a \in N_4(R)$ . Then

$$0 \leq (a + na^2)^2 = a^2 + n(2a^3)$$

holds for all  $n \in \mathbb{Z}$ , hence  $2a^3 = 0$ . Conversely,  $2a^3 = 0$  implies  $2a^4 = 0$ , and consequently  $a^4 = 0$ . The proof of (1) is complete, while (2) evidently follows from (1). □

**Lemma 1.4.** *Let  $R$  be an Archimedean ps-ring and let  $a \in N(R)$ . Then*

- (1)  $(ab)^2 = (ba)^2 = 0$  for all  $b \in R$ .
- (2)  $abc = -cab = bca$  for all  $b, c \in R$ .
- (3)  $abcd = bacd = bcad = bcda \in T_2(R)$  for all  $b, c, d \in R$ .

PROOF: Since by Proposition 1.3 we have  $a^2 \in N_2(R)$ , Lemma 1.1. (2) implies  $2aba^2 = 0$  for all  $b \in R$ . Therefore  $2(aba)^2 = 0$  and consequently  $(aba)^2 = 0$ , since  $R$  has positive squares. It follows that

$$0 \leq (b + naba)^2 = b^2 + n((ab)^2 + (ba)^2)$$

holds for all  $n \in \mathbb{Z}$ , which implies (1).

To prove (2), combine (1) and Lemma 1.1. (1), while to obtain (3), note that by (1)  $ab, ba, ad, da \in N_2(R)$  for all  $b, d \in R$ , and then use Lemma 1.1. (2). □

We are prepared to prove a generalization of a result of Diem [3] and Bernau, Huijsmans [1], Propositions 3.1 and 3.2.

**Theorem 1.5.** *Let  $R$  be an Archimedean ps-ring. Then  $N(R)$  is an order-convex ideal of  $R$ , satisfying*

$$\begin{aligned} N(R) = N_4(R) &= \{a \in R : abc + cab = 0 \text{ for all } b, c \in R\} \\ &= \{a \in R : bca + cab = 0 \text{ for all } b, c \in R\} \\ &= \{a \in R : 2a^3 = 0\} \\ &= \{a \in R : 2a \in \text{ann}(R^3)\}. \end{aligned}$$

PROOF: Combine Proposition 1.3 and Lemma 1.4 to obtain the required equalities for  $N(R)$ . The last one implies that  $N(R)$  is an ideal, and hence it is convex. □

It can be shown similarly that  $N_2(R)$  is an order-convex ideal of  $R$ .

**Corollary 1.6.** *Let  $R$  be an Archimedean  $ps$ -ring and  $Z(R)$  its center. Then*

- (1)  $N(R)^2R = N(R)RN(R) = RN(R)^2 \subset T_2(R)$ ;
- (2)  $N_2(R) \subset Z(R)$  implies  $N(R)R^2 = RN(R)R = R^2N(R) \subset T_2(R)$ ;
- (3)  $R = R^2$  implies  $N(R) \subset Z(R)$ ,  $N(R) = N_2(R)$ ,  $N(R)R = RN(R) \subset T_2(R)$ .

*If  $R$  is also 2-torsion-free, then*

- (4)  $N(R) = N_3(R) = \text{ann}(R^3)$ ;
- (5)  $N_2(R) \subset Z(R)$  implies  $N(R) = \text{ann}(R^2)$ ;
- (6)  $R = R^2$  implies  $N(R) = N_2(R) = \text{ann}(R)$ .

PROOF: (1) Use Lemma 1.4. (2) to see that  $a, b \in N(R)$ ,  $c \in R$  implies  $abc = -cab = bca = -abc$ , and the result follows.

(2) If  $a \in N(R)$ ,  $b, c \in R$ , then  $ab \in N_2(R) \subset Z(R)$ , and therefore  $abc = cab$ . By Lemma 1.4. (2) we have  $2abc = abc + cab = 0$ , so (2) follows easily.

(3) Lemma 1.4. (2) implies that  $N(R) \subset Z(R)$ , thus by (2)  $N(R)R = RN(R) \subset T_2(R)$ . It follows that each  $a \in N(R)$  satisfies  $2a^2 = 0$ , hence  $a \in N_2(R)$ , and the proof is complete.

(4), (5), (6) Use Theorem 1.5 and (1), (2), (3). □

**Remark 1.7.** An Archimedean  $ps$ -ring  $R$  with generating cone  $R^+$  is torsion-free. Indeed, if  $a = a_1 - a_2$  with  $a_1, a_2 \in R^+$  satisfies  $ma = 0$  for some  $m \in \mathbb{N}$ , then it follows that

$$0 \leq m(a_1 + a_2) + na \quad \text{for all } n \in \mathbb{Z},$$

which implies that  $a = 0$ .

It can be proved, that the set  $T(R)$  of all torsion elements of  $R$  is an order-convex ideal of  $R$ , and that the quotient ring  $R/T(R)$  is an Archimedean torsion-free  $ps$ -ring.

**Theorem 1.8.** *Let  $R$  be an Archimedean  $ps$ -ring. Then  $N(R)$  is the unique nil radical of  $R$ . It is locally nilpotent and satisfies  $N(R)^3 \subset T_2(R)$ . If  $R$  is 2-torsion-free, then  $N(R)$  is nilpotent with  $N(R)^3 = \{0\}$ .*

PROOF: A nil radical of bounded index is contained in the lower nil radical ([7, p. 232]), hence by Theorem 1.5  $N(R)$  is the unique nil radical of  $R$ . Thus  $N(R)$  equals the Levitzki nil radical  $L(R)$  of  $R$ , which is locally nilpotent ([9, Proposition 21.2]). The required inclusion follows by Corollary 1.6 and implies the remaining part of the theorem. □

It is proved in [3, Theorem 3.9. (v)] that an Archimedean lattice-ordered ring with positive squares and with zero left or right annihilator has no nonzero positive nilpotents. The following generalization of this result is a simple consequence of the last equation of Theorem 1.5.

**Proposition 1.9.** *If  $R$  is an Archimedean  $ps$ -ring such that  $\text{ann}(R) = \{0\}$ , then  $N(R) = T_2(R)$ .  $\square$*

**Corollary 1.10.** *Let  $R$  be an Archimedean  $ps$ -ring. Then the following statements are equivalent.*

- (i)  $R$  is semiprime.
- (ii)  $R$  is reduced, i.e. without non-zero nilpotents.
- (iii)  $R$  is 2-torsion-free and satisfies  $\text{ann}(R) = \{0\}$ .

$\square$

Note that by Remark 1.7 a lattice-ordered  $R$  is torsion-free, hence the above characterization of reduced rings  $R$  improves [3, Theorem 3.9. (v)].

**Corollary 1.11.** *A unital Archimedean  $ps$ -ring is reduced (or semiprime) if and only if it is 2-torsion-free.  $\square$*

## 2. Examples

By Theorem 1.5 an Archimedean  $ps$ -ring satisfies  $N(R) = N_4(R)$ . We show that in general  $N(R) \neq N_3(R)$ .

**Example 2.1.** Let  $R = \mathbb{Z}a \oplus \mathbb{Z}b \oplus \mathbb{Z}_2c$  be the ring with multiplication defined by

$$a^2 = b, ab = ba = c, b^2 = c^2 = bc = cb = ac = ca = 0,$$

and ordered by the positive cone  $R^+ = \mathbb{Z}^+b$ . It is easy to see that  $R$  is an Archimedean  $ps$ -ring with

$$N_3(R) = \mathbb{Z}(2a) \oplus \mathbb{Z}b \oplus \mathbb{Z}_2c$$

and  $N_4(R) = R$ , thus  $N_3(R) \neq N(R)$ .

By Theorem 1.8 every nil Archimedean  $ps$ -ring  $R$  is locally nilpotent, and even nilpotent with  $R^3 = \{0\}$  when  $R$  is 2-torsion-free. In the following example we present a nil Archimedean  $ps$ -ring  $R$  which is not nilpotent.

**Example 2.2.** If  $m \in \mathbb{Z}^+$ , let  $\beta_0(m), \beta_1(m), \dots$  be the digits in the binary representation

$$m = \sum_{i=0}^{\infty} \beta_i(m)2^i, \beta_i(m) \in \{0, 1\}$$

of  $m$ . Define the function  $\varphi : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$  by

$$\varphi(p, q) = \begin{cases} 0, & \text{if } \beta_i(p) = \beta_i(q) = 1 \text{ for some } i \\ p + q, & \text{otherwise} \end{cases}$$

and consider the ring  $R = (\mathbb{Z}_2)^\infty$  with coordinatewise addition and with the multiplication defined by

$$e_p e_q = e_{\varphi(p, q)}, \quad p, q \in \mathbb{N},$$

where  $e_n = (\underbrace{0, \dots, 1}_n, 0, \dots)$  and  $e_0 = 0 = (0, 0, \dots)$ .

It is easy to see that  $R$  is an associative ring which satisfies  $R = T_2(R) = N_2(R)$ . It follows that the positive cone  $R^+ = \{0\}$  turn  $R$  into a nil Archimedean  $ps$ -ring. Note that for each  $n \in \mathbb{N}$

$$e_{2^n-1} = e_1 e_2 e_4 \cdots e_{2^{n-1}} \in R^n,$$

thus  $R$  is not nilpotent.

If  $R$  is 2-torsion-free, then by Corollary 1.6 the chain  $ann(R) \subset ann(R^2) \subset \dots$  is finite and satisfies  $ann(R^k) = N(R)$  for all  $k \geq 3$ . Example 2.2 show that this is not the case in a general situation. Moreover, there exists an Archimedean  $ps$ -ring  $S$  with strictly increasing chain  $ann(S) \subset ann(S^2) \subset \dots$ .

**Example 2.3.** Let  $R$  be the ring from Example 2.2. For each  $n \in \mathbb{N}$  denote by  $R_n$  the additive subgroup of  $R$  generated by elements  $e_k$  with  $1 \leq k \leq 2^n - 1$ , and observe that  $R_n$  is a subring of  $R$ , since

$$0 \leq i, j \leq 2^n - 1 \text{ implies } \varphi(i, j) \leq 2^n - 1.$$

Let  $S$  be the direct product of all  $R_n$  with componentwise defined operations. Then  $S$  is an Archimedean  $ps$ -ring with the chain of annihilators

$$ann(S) \subset ann(S^2) \subset ann(S^3) \subset \dots$$

strictly increasing and contained in  $N_2(S) = S$ .

It may be asked (see Corollary 1.11) if a unital Archimedean  $ps$ -ring is automatically 2-torsion-free, and therefore reduced. The following example shows that this is not the case.

**Example 2.4.** Denote by  $J$  the principal ideal of  $\mathbb{Z}_4$  generated by 2, and consider its unitization ring  $R = J \oplus \mathbb{Z}$  ordered by the positive cone  $R^+ = \{0\} \oplus \mathbb{Z}^+$ .

It is easy to see that  $R$  is a unital Archimedean  $ps$ -ring satisfying

$$T_2(R) = N(R) = J \oplus \{0\},$$

thus  $R$  is not 2-torsion-free.

### 3. An application

We shall need a result on quotient ring  $R/N(R)$ .

**Lemma 3.1.** *Let  $R$  be an Archimedean  $ps$ -ring. Then  $R/N(R)$  is a reduced Archimedean  $ps$ -ring.*

PROOF: Since by Theorem 1.5  $N(R)$  is an order-convex ideal of  $R$ , the quotient ring  $R/N(R)$  is partially ordered by the positive cone  $\pi(R^+)$ , where  $\pi$  denotes

the canonical projection of  $R$  onto  $R/N(R)$ . A simple verification shows that  $R/N(R)$  is a reduced  $ps$ -ring.

The proof will be complete by proving that  $R/N(R)$  is Archimedean. To this end suppose that elements  $\pi(a), \pi(b) \in R/N(R)$  satisfy  $0 \leq \pi(b) + n\pi(a)$  for all  $n \in \mathbb{Z}$ . Then there exist elements  $a_n \in N(R)$ ,  $n \in \mathbb{Z}$  such that

$$0 \leq b + na + a_n.$$

Multiplying this inequality by  $2a^4$  and observing that  $2a^4a_n = 0$  we get

$$0 \leq 2a^4b + n(2a^5)$$

for all  $n \in \mathbb{Z}$ . Since  $R$  is Archimedean this implies  $2a^5 = 0$ . It follows that  $a \in N(R)$ , hence  $\pi(a) = 0$ , as required.  $\square$

Recall that a partially ordered ring  $R$  is said to be an  $f$ -ring if it is lattice-ordered and if  $a, b \in R$  with  $a \wedge b = 0$  implies  $ac \wedge b = ca \wedge b = 0$  for all  $c \in R^+$ . It is well known that an Archimedean  $f$ -ring is commutative. Let us apply our results on partially ordered rings which are closely related to  $f$ -rings.

We shall say that a partially ordered ring  $R$  is  $f$ -decomposable, if every element  $a \in R$  is expressed as  $a = a_1 - a_2$  with  $a_1, a_2 \in R^+$  and  $a_1a_2 = a_2a_1 = 0$ .

Observe that an Archimedean  $f$ -decomposable ring is a  $ps$ -ring with generating cone  $R^+$ , and therefore torsion-free by Remark 1.7.

**Theorem 3.2.** *Let  $R$  be an Archimedean  $f$ -decomposable partially ordered ring. Then*

- (1)  $R/N(R)$  is an Archimedean  $f$ -ring;
- (2) all triples of elements of  $R$  commute, that is

$$a_1a_2a_3 = a_{\sigma(1)}a_{\sigma(2)}a_{\sigma(3)} \quad \text{for all } a_1, a_2, a_3 \in R, \sigma \in S_3.$$

PROOF: The quotient ring  $R/N(R)$  is reduced and  $f$ -decomposable, hence (1) follows by [6] and Lemma 3.1.

To prove (2) note that  $R$  satisfies

$$N_2(R) = R^+ \cap N_2(R) - R^+ \cap N_2(R),$$

which by Lemma 1.1. (1) implies that  $N_2(R) = ann(R)$ . It follows by Corollary 1.6 that  $N(R) = ann(R^2)$ , hence

$$N(R) \cap R^2 \subset N_2(R) = ann(R).$$

The commutativity of the Archimedean  $f$ -ring  $R/N(R)$  implies that  $ab - ba \in N(R) \cap R^2 \subset ann(R)$ , therefore (2) follows.  $\square$

Applying Proposition 1.9 we get



**Corollary 3.3.** *Let  $R$  be an Archimedean  $f$ -decomposable partially ordered ring.*

- (1) *If  $\text{ann}(R) = \{0\}$ , then  $R$  is an Archimedean semiprime  $f$ -ring.*
- (2) *If  $R = R^2$ , then  $R$  is commutative.* □

Let  $R$  be an Archimedean  $f$ -decomposable ring. It can be seen (using for example [1, Proposition 1.3]) that if  $R$  is lattice-ordered then it is an almost  $f$ -ring, thus Corollary 3.3. (1) generalizes [1, Theorem 1.11 (ii)]. Moreover, in this case  $R$  is commutative by [1, Theorem 2.15], and it might be interesting to know whether  $R$  is commutative also in general case.

#### REFERENCES

- [1] Bernau S.J., Huijsmans C.B., *Almost  $f$ -algebras and  $d$ -algebras*, Proc. Cambridge Philos. Soc. **107** (1990), 287–308.
- [2] Birkhoff G., Pierce R.S., *Lattice-ordered rings*, An. Acad. Brasil Ciênc. **28** (1956), 41–69.
- [3] Diem J.E., *A radical for lattice-ordered rings*, Pacific J. Math. **25** (1968), 71–82.
- [4] Divinsky N., *Rings and Radicals*, Allen, London, 1965.
- [5] Fuchs L., *Partially Ordered Algebraic Systems*, Pergamon Press, Oxford-London-New York-Paris, 1963.
- [6] Hayes A., *A characterization of  $f$ -rings without non-zero nilpotents*, J. London Math. Soc. **39** (1964), 706–707.
- [7] Jacobson N., *Structure of Rings*, Colloquium Publication **37**, Amer. Math. Soc., Providence, 1956.
- [8] Steinberg S.A., *On lattice-ordered rings in which the square of every element is positive*, J. Austral. Math. Soc. Ser. A **22** (1976), 362–370.
- [9] Szász F.A., *Radicals of Rings*, Akadémiai Kiado – John Wiley & Sons, Budapest-Chichester-New York-Brisbane-Toronto, 1981.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF LJUBLJANA, JADRANSKA 19, 61000 LJUBLJANA, SLOVENIA

(Received August 18, 1993)