

Anatoly A. Gryzlov

Cardinal invariants and compactifications

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 2, 403--408

Persistent URL: <http://dml.cz/dmlcz/118679>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Cardinal invariants and compactifications

A. GRYZLOV

Abstract. We prove that every compact space X is a Čech-Stone compactification of a normal subspace of cardinality at most $d(X)^{t(X)}$, and some facts about cardinal invariants of compact spaces.

Keywords: Čech-Stone compactification, cardinal invariants

Classification: 54A25, 54D35

The notions of a compactness and of a compactification are very closed. Really, any compact space is a compactification of any of its dense subsets. But a useful information can be obtained from this fact only in the case when a compact space is a compactification of some certain type.

The classic type is the Čech-Stone compactification. When a compact space is a Čech-Stone compactification of some of its subsets? Theorem 1.1 says that any compact space X is a Čech-Stone compactification of some normal subspace $B \subseteq X$ such that $|B| \leq d(X)^{t(X)}$, where $d(X)$ is a density of X , $t(X)$ is its tightness. The case, when $|X| \leq d(X)^{t(X)}$ is trivial, in this case $B = X$. In the opposite case the theorem says that “extra” points from $X \setminus B$ are constructed by a standard way, as points of Čech-Stone compactification of a “not large” normal subspace. For example, by this theorem, we can say that Fedorchuk’s compact space, that is the hereditarily separable, hereditarily normal compact space of cardinality $2^{\mathfrak{c}}$, is a Čech-Stone compactification of a subspace of a cardinality \mathfrak{c} . We prove some new facts about cardinal invariants.

The definitions and notations used here are standard, one can find them in [2], for example.

We use the notation of a sequential extension of a set A , that is a set B , $A \subseteq B \subseteq [A]$, $|B| \leq |A|^\omega$ such that if a countable set $B' \subseteq B$ has a limit point in X , then there is a limit point of B' in B . One can construct a sequential extension by induction.

Theorem 1.1. *Let X be a compact space, A be a dense subset of X . Then $X = \beta B$, where $A \subseteq B$, $|B| \leq |A|^{t(X)}$, B is a normal and countably compact subspace.*

PROOF: The case $|X| = |A|^{t(X)}$ is trivial. Let $|X| > |A|^{t(X)}$. We construct by induction the family $\{B_\alpha : \alpha < \omega_{\tau+}\}$, where $t(X) = \tau$ such that:

- (1) $B_0 = A$;
- (2) $B_\beta \subseteq B_\alpha$ for $\beta \leq \alpha$;
- (3) a sequential extension of B_α is in $B_{\alpha+1}$;
- (4) $|B_\alpha| \leq |A|^\tau$ for $\alpha < \omega_{\tau+}$;
- (5) $\bigcup\{B_\alpha : \alpha < \omega_{\tau+}\} = B$, where B is a normal, countably compact subspace and $X = \beta B$.

Let $\xi(A)$ be a choice function, defined on the set of all nonempty subsets of X . Let $B_0 = A$. Let $\{B_\beta : \beta < \alpha\}$ with conditions (1)–(4) be constructed. Let B'_α be a sequential extension of the set $\bigcup\{B_\beta : \beta < \alpha\}$ and

$$B_\alpha = B'_\alpha \cup \{\xi([T]_X \cap [T']_X) : T, T' \in \text{exp}_\tau B'_\alpha\}.$$

Then $|B_\alpha| \leq |A|^\tau$ and (1)–(4) hold for $\{B_\beta : \beta \leq \alpha\}$. Note that B is a countably compact space. Let us prove that $X = \beta B$. Let F_1, F_2 be disjoint closed subsets of B . Let $([F_1]_X \cap [F_2]_X) \setminus B \neq \emptyset$ and $x \in ([F_1]_X \cap [F_2]_X) \setminus B$. Since $t(X) = \tau$, there are $F'_1 \subseteq F_1$ and $F'_2 \subseteq F_2$ such that $F'_1, F'_2 \subseteq B_\alpha$ and $|F'_1| \leq \tau$, $|F'_2| \leq \tau$, and $x \in [F'_1]_X \cap [F'_2]_X$. There is $\alpha < \omega_{\tau+}$ such that $F'_1, F'_2 \subseteq B_\alpha$. Since F'_1, F'_2 are disjoint, closed subsets of B , then $[F'_1]_X \cap [F'_2]_X \subseteq ([F_1]_X \cap [F_2]_X) \setminus B$. Then $\xi([F'_1]_X \cap [F'_2]_X) \in X \setminus B$, a contradiction. So $X = \beta B$. It follows from the above proof that B is a normal space. The theorem is proved. \square

Recall that a space X is weakly normal if in every closed, countable, discrete set $A \subseteq X$ there is a countable $A' \subseteq A$ C^* -embedded in X [3].

We say that X is an h -weakly normal space if X is hereditarily weakly normal.

Definition 1.2. A space X is called d -normal if in every closed, countable, discrete set $A \subseteq X$ there is a countable $A' \subseteq A$ with discrete family of neighborhoods. It means that for every point $x \in A'$ there is a neighborhood Ox such that $\{Ox : x \in A'\}$ is a discrete family.

We say that a space X is hd -normal if X is a hereditarily d -normal.

It is clear that a hd -normal (d -normal) space is h -weakly (weakly) normal, and a hereditarily normal space is hd -normal as well as it is compact first countable space or a normal first countable space.

On the other hand, the space $N^* = \beta N \setminus N$, the remainder of the Čech-Stone compactification of a countable discrete space, is an h -weakly normal, but not an hd -normal space. Really, the h -weakly normality of N^* follows from the fact that $[D]_{N^*}$ is homeomorphic to $\beta N = \beta D$ for every countable discrete set $D \subseteq N^*$. But for a discrete set D the space $X = N^* \setminus ([D]_{N^*} \setminus D)$ is not d -normal.

Theorem 1.3. Let X be an hd -normal compact space, or an h -weakly normal space with countable tightness, and $A \subseteq X$ be a dense subset of X . Then $X = \beta B$ where $A \subseteq B$, $|B| \leq |A|^{t(X)}$, B is a normal countably compact space such that every compact space $K \subseteq X \setminus B$ is finite.

PROOF: Again, we suppose that $|X| > |A|^{t(X)}$. Let the set B be as in Theorem 1.1. We prove that every compact space $K \subseteq X \setminus B$ is finite. Let $K \subseteq X \setminus B$

be an infinite compact set. There is a countable discrete (as a subspace) subset $D \subseteq K$. We consider a set $B \cup D$. By h -weakly normality (moreover hd -normality) of X , there is a countable set $D' \subseteq D$ C^* -embedded in $B \cup D$. Then $[D']_X = \beta D'$, so $[D']_X$ is a Čech-Stone compactification of the countable discrete set. But $\beta D' = \beta N$ is not an hd -normal and the tightness of $\beta D'$ is not countable. This contradiction proves the theorem. \square

Theorems 1.1 and 1.3 were announced by the author in [4].

Lemma 1.4. *Let X be an h -weakly normal compact space with countable tightness, $X = \beta B$ for some $B \subseteq X$. The $\Phi \setminus F$ is a discrete (as a subspace) set for closed $F, \Phi \subseteq X$ such that $F \subseteq \Phi$ and $F \cap B = \Phi \cap B$.*

PROOF: Let $x \in \Phi \setminus F$. There is a neighborhood Ox of x such that $[Ox]_X \cap F = \emptyset$. Therefore, $[Ox]_X \cap \Phi = [Ox]_X \cap (\Phi \setminus F) \subseteq X \setminus B$. By Theorem 1.3, the set $[Ox]_X \cap \Phi$ is finite, and therefore $\Phi \setminus F$ is a discrete set. The lemma is proved. \square

Recall that a point $x \in X$ is called a b -point if $x = F \cap \Phi$ where F and Φ are closed sets in X , and x is a limit point for F and Φ [5].

\varkappa -points (limits of sequences of points of X) are b -points as well as points of non-normality (points x of a normal space X such that $X \setminus \{x\}$ is non-normal).

Theorem 1.5. *Let X be an h -weakly normal compact space with countable tightness. Then $|\{x : x \text{ is a } b\text{-point in } X\}| \leq d(X)^\omega$.*

PROOF: By Theorem 1.3, $X = \beta B$ for a normal B such that $|B| \leq d(X)^\omega$ and every compact subset of $X \setminus B$ is finite. We prove that none of the points of $X \setminus B$ is a b -point. Indeed, let $x \in X$ be a b -point, that is, $x = F \cap \Phi$ where F, Φ are closed in X , and x is a limit point for F and Φ . Let $F' = F \cap B$, $\Phi' = \Phi \cap B$. Then $x \in [F']_X \cap [\Phi']_X$. Really, by Lemma 1.4, the sets $F \setminus [F']_X$ and $\Phi \setminus [\Phi']_X$ are discrete and therefore $x \in [F']_X, x \in [\Phi']_X$. But X is a Čech-Stone compactification of the normal space B . This contradiction proves the theorem. \square

Corollary 1.6. *Let X be a weakly normal compact space with countable tightness and let all points of X be b -points. Then $|X| \leq d(X)^\omega$.*

The above results have some connections with the results from [6].

Recall that $x \in X$ is an hb -point if x is a b -point in every closed subspace $X' \subseteq X$, if x is not isolated in X' [5].

By Arhangel'skii's theorem [7] and Theorem 1.5 we have

Theorem 1.7. *Let X be an h -weakly normal compact space with countable tightness, $\chi(X) \leq 2^\omega$ and every non-isolated point in X is an hb -point. Then $|X| \leq 2^\omega$.*

Proposition 1.8. *Let X be a countably compact hd -normal space, $A \subseteq X$ be a dense subset. Then there is $B \subseteq X$ such that $A \subseteq B$, $|B| \leq |A|^\omega$, B is countably compact such that every subset $F \subseteq X \setminus B$ closed in X is finite.*

PROOF: It is clear that B is a sequential extension of A . The only thing we have to explain is the last part. Let $F \subseteq X \setminus B$ be an infinite closed subset of X . There is a countable discrete (as a subspace) set $D \in F$. By an hd -normality of X there is a countable set $D' = \{x_i : i \in \omega\}$, $D' \subseteq D$ with a discrete in $B \cup D$ family of neighborhoods $\{Ox_i : i \in \omega\}$. But $Ox_i \cap B \neq \emptyset$ for every $i \in \omega$, so we have the discrete infinite family in the compact space B . This contradiction proves the proposition. \square

By the same way as Lemma 1.4 we can prove

Lemma 1.9. *Let X be an hd -normal space, $B \subseteq X$ be a dense, countably compact subspace, $F, \Phi \subseteq X$ such that $F \subseteq \Phi$, $F \cap B = \Phi \cap B$. Then $\Phi \setminus F$ is a discrete (as a subspace) set, moreover, $\Phi \setminus F$ is a free sequence in $X \setminus F$.*

Lemma 1.10. *Let X be an hd -normal space, $B \subseteq X$ be a countably compact subspace. Then for every closed set $F \subseteq X$ there is a family $\pi = \{OF\}$ of neighborhoods of F such that $|\pi| \leq |B|$, $(\bigcap\{[OF] : OF \in \pi\} \cap [B]) \setminus F \cap [B]$ is discrete; moreover, if $F \cap B = \emptyset$, then $\bigcap\{[OF] : OF \in \pi\} \cap [B] = \emptyset$.*

PROOF: We consider $F \cap [B]$. It is clear that there is a family $\pi = \{OF\}$ of neighborhoods of F such that $|\pi| = |B|$, $\bigcap\{[OF] : OF \in \pi\} \cap B = F \cap B$. Then $(\bigcap\{[OF] : OF \in \pi\} \cap [B]) \setminus F \cap [B]$ is discrete by Lemma 1.9. If $F \cap B = \emptyset$, then $\bigcap\{[OF] : OF \in \pi\} \cap [B] \subseteq [B] \setminus B$. In the same way as in the proof of Proposition 1.7 we can prove that $\bigcap\{[OF] : OF \in \pi\} \cap [B]$ is finite. We can add a finite number of neighborhoods of F to the family π and get what we need. The lemma is proved. \square

Proposition 1.11. *Let X be an hd -normal, countably compact space such that every point $x \in X$ is a limit point of a closed set of cardinality at most $d(X)^{t(X)}$. Then $|X| \leq d(X)^{t(X)}$.*

PROOF: By Proposition 1.8 there is a dense, countably compact space $B \subseteq X$ such that every closed set $F \subseteq X \setminus B$ is finite and $|B| \leq d(X)^{t(X)}$. Let $x \in X \setminus B$. There is a closed $F_x \subseteq X$ such that $|F_x| \leq d(X)^{t(X)}$ and x is a limit point of F_x . Then by Lemma 1.9, $x \in [F_x \cap B]$. There is $F'_x \subseteq F_x \cap B$ such that $|F'_x| \leq t(X)$ and $x \in [F'_x]$. Then $|X \setminus B| \leq |B|^{t(X)} \leq d(X)^{t(X)}$. The proposition is proved. \square

Proposition 1.12. *Let X be an hd -normal compact space. Then $hl(X) \leq s(X)^\omega$.*

PROOF: We prove that $\chi(F, X) \leq s(X)^\omega$ for every closed $F \subseteq X$. Really, for a closed $F \subseteq X$ there is a family $\pi = \{OF\}$ of neighborhoods of F such that $|\pi| \leq s(X)$ and $d(\bigcap\{[OF] : OF \in \pi\} \setminus F) \leq s(X)$ (this is well known, see for

example [5]). By Proposition 1.8 there is a set $B \subseteq \bigcap \{[OF] : OF \in \pi\}$ such that $|B| \leq s(X)^\omega$, B is countably compact, $[B] \supseteq (\bigcap \{[OF] : OF \in \pi\}) \setminus F$ and every subset of B closed in X is finite. By Lemma 1.9 there is a family $\pi' = \{UF\}$, $|\pi'| \leq s(X)^\omega$ of neighborhoods of F such that $(\bigcap \{[UF] : UF \in \pi\} \cap [B]) \setminus F \cap [B]$ is discrete and therefore has cardinality at most $s(X)$. Finally, $\chi(F, X) \leq s(X)^\omega \cdot s(X) = s(X)^\omega$. The proposition is proved. \square

Recall that a free sequence of cardinality τ is a set $\xi\{x_\alpha : \alpha < \tau\}$ such that for all $\beta < \tau$ $\{x_\alpha : \alpha < \beta\} \cap \{x_\alpha : \alpha \geq \beta\} = \emptyset$ (see [6]).

Define $A(X) = \sup\{\tau : \tau \text{ is cardinality of a free sequence in } X\}$, $\rho A(x, X) = A(X \setminus \{x\})$, $\rho A(X) = \sup\{\rho A(x, X) : x \in X\}$. A. Arhangel'skii proved that $t(X) = A(X)$ for compact spaces [7]; moreover,

$$t(X) = A(X) \leq \rho A(X) \leq s(X).$$

Note that for Alexandroff's double circle $s(X) = 2^\omega$, $\rho A(X) = A(X) = \omega$. The same construction with Fedorchuk's compact space gives the space with $s(X) = 2^c$ and $\rho A(X) = \omega$.

Theorem 1.13. *Let X be an hd -normal compact space. Then $\chi(x, X) \leq \rho A(x, X)^\omega$.*

PROOF: Let there be a point $x \in X$ such that $\rho A(x, X)^\omega < \chi(x, X)$. Define $\rho A(x, X) = \tau$. By induction we construct a set $D = \{y_\alpha : \alpha < \omega_{\tau^+}\}$, a family $\{B_\alpha : \alpha < \omega_{\tau^+}\}$, $|B_\alpha| \leq \tau^\omega$ of neighborhoods of x such that $(\{y_\alpha : \alpha < \delta\} \cap \bigcap \{[Ox] : Ox \in B_\delta\}) \setminus \{x\} = \emptyset$, $\delta < \omega_{\tau^+}$. Let $y_0 \in X$, $B_0 = \{Ox\}$, where $[Ox] \not\ni y_0$. Let $\{y_\alpha : \alpha < \delta\}$ and $\{B_\alpha : \alpha < \delta\}$ be constructed. If $x \notin [\{y_\alpha : \alpha, \delta\}]$, let $B_\delta = \bigcup \{B_\alpha : \alpha < \delta\} \cup \{Ox\}$, where $Ox \cap \{y_\alpha : \alpha < \delta\} = \emptyset$. We choose y_δ in the set $\bigcap \{[Ox] : Ox \in B_\delta\} \setminus \{x\}$. If $x \in [\{y_\alpha : \alpha < \delta\}]$, we use Proposition 1.6 and Lemma 1.9. Since $|\{y_\alpha : \alpha < \delta\}| \leq \tau$, let us consider a family $\pi = \{Ox\}$ of neighborhoods of x , $|\pi| \leq \tau^\omega$ such that $T = (\bigcap \{[Ox] : Ox \in \pi\}) \cap [\{y_\alpha : \alpha < \delta\}] \setminus \{x\}$ is empty or is a free sequence in $X \setminus \{x\}$, and $|T| \leq \rho A(x, X)$. Hence, there is a family π' of neighborhoods of x of cardinality at most τ^ω such that $(\bigcap \{[Ox] : Ox \in \pi\} \cap [\{y_\alpha : \alpha < \delta\}]) \setminus \{x\} = \emptyset$. Let $B_\delta \cup \{B_\alpha : \alpha < \delta\} \cup \pi'$, and choose y_δ from $\bigcap \{[Ox] : Ox \in B_\delta\} \setminus \{x\}$. If we continue until ω_{τ^+} , we get a free sequence of cardinality τ^+ . But this contradicts $\rho A(x, X) = \tau$. The theorem is proved. \square

REFERENCES

- [1] Fedorchuk V.V., *Vpolne zamknutyje otobrazhenija*, Matem. sbornik **99:1** (1976), 3–33.
- [2] Arhangel'skii A.V., *Stroenie i klassifikacija topologicheskikh prostranstv i kardinalnye invarianty*, Uspekhi mat. nauk **33:6** (1978), 29–84.
- [3] Pasynkov B.A., *O teoreme E. Čecha*, Vest. MGU **6** (1979), 54–57.
- [4] Gryzlov A.A., *On cardinal invariants of compact spaces*, Abstr. Baku Intern. Top. Conf., 1987, p. 89.
- [5] Šapirovs'kii B.E., *O vložhenii ekstremalno-normalnyh prostranstv v bikompakty*, Dokl. Akad. Nauk SSSR **223:5** (1975), 1083–1086.

- [6] ———, *O nasledstvenno normalnykh prostranstvakh*, Abstr. VII Vses. Top. Conf. Minsk, 1977, p. 204.
- [7] Arhangel'skii A.V., *O moschnosti bikompaktov s pervoi aksiomoi schetnosti*, Dokl. Akad. Nauk SSSR **187** (1969), 967–970.

DEPARTMENT OF MATHEMATICS, CHAIR OF TOPOLOGY, UDMURTSK STATE UNIVERSITY,
71 KRASNOGEROISKAIA STR., 26031 IJEVSK, RUSSIA

(Received August 30, 1993)