

R. J. Nájares; Luděk Zajíček

A σ -porous set need not be σ -bilaterally porous

Commentationes Mathematicae Universitatis Carolinae, Vol. 35 (1994), No. 4, 697--703

Persistent URL: <http://dml.cz/dmlcz/118711>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1994

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A σ -porous set need not be σ -bilaterally porous

J. NÁJARES R., L. ZAJÍČEK*

Abstract. A closed subset of the real line which is right porous but is not σ -left-porous is constructed.

Keywords: sigma-porous, sigma-bilaterally-porous, right porous

Classification: Primary 26A99; Secondary 28A5

1. Introduction

Let $E \subset \mathbb{R}$ be a set, and let I be an interval. Then we denote by $\lambda(E, I)$ the length of the largest open subinterval of I which does not intersect E . The right porosity of E at $x \in \mathbb{R}$ is defined as

$$p^+(E, x) = \overline{\lim}_{h \rightarrow 0^+} \frac{\lambda(E, (x, x + h))}{h}.$$

The left porosity $p^-(E, x)$ is defined by the symmetrical way.

We say that:

- (i) E is right porous at x if $p^+(E, x) > 0$,
- (ii) E is left porous at x if $p^-(E, x) > 0$,
- (iii) E is bilaterally porous at x if it is porous both on the right and on the left at x .

The set E is said to be right (left, bilaterally) porous if it is right (left, bilaterally) porous at each of its points and σ -right-porous (σ -left-porous, σ -bilaterally-porous) if it is a countable union of right (left, bilaterally) porous sets. It is easy to see that a set is σ -bilaterally porous iff it is bilaterally σ -porous (i.e. it is both σ -right-porous and σ -left-porous). The main aim of the present article is to prove the following result.

Theorem. *There exists a closed set $F \subset \mathbb{R}$ which is right porous but is not σ -left-porous.*

We obtain the example slightly modifying the ideas of [F] and [Za 1].

We essentially use Lemma 5 which is a special case of the generalized Foran lemma [Za 3], which enables us to give a simple proof that our set F is not

*Supported by Research Grant GAUK 363.

σ -left-porous. Another ingredient of our proof is Proposition, which is analogous to Proposition 4.4 from [Za 2]. We believe that it can be also of some independent importance. Note that for symmetrical porosity an analogical proposition does not hold [E-H-S].

2. Proposition and lemmas

Definition 1. If $c > 0$, $M \subset \mathbb{R}$ and $r > 0$ are given, then we define

$$S(c, r, M) = \bigcup \{x \ominus (y - \sigma, y); y \in \mathbb{R}, 0 < \sigma < r, (y - \sigma, y) \cap M = \emptyset\},$$

where $c \ominus (y - \sigma, y) = (y - c\sigma, y)$.

We shall need the following lemmas which are obvious.

Lemma 1. If $p^+(M, x) \geq c > 0$, then $x \in \bigcap \{S(\frac{2}{c}, r, M); r > 0\}$.

Lemma 2. If $c > 1$, $x \in M$ and $x \in \bigcap \{S(c, r, M); r > 0\}$, then $p^+(M, x) \geq \frac{1}{c}$.

Proposition. Let A be a σ -right-porous set (σ -left-porous) and $c < 1$. Then there exists a sequence $\{A_n\}_{n=1}^\infty$ such that $A = \bigcup_{n=1}^\infty A_n$ and $p^+(A_n, x) \geq c$ ($p^-(A_n, x) \geq c$, respectively) for any $n \in \mathbb{N}$ and $x \in A_n$.

PROOF: It is sufficient to give the proof for right porosity only. By definition $A = \bigcup_{n=1}^\infty B_n$ where B_n is a right porous set for any $n \in \mathbb{N}$. Putting

$$B_{n,k} = \{x \in B_n; p^+(B_n, x) \geq \frac{1}{k}\}$$

we have that $A = \bigcup_{n,k=1}^\infty B_{n,k}$ and $p^+(B_{n,k}, x) \geq \frac{1}{k}$ for any $x \in B_{n,k}$.

Thus it is sufficient to prove the following statement:

If $M \subset \mathbb{R}$, $a > 0$ and, for each $x \in M$, the inequality $p^+(M, x) \geq a$ holds, then $M = \bigcup_{i=1}^\infty M_i$, where $p^+(M_i, y) \geq c$ for any $y \in M_i$.

We can suppose $a < c < 1$, the case $a \geq c$ being trivial. Choose $n \in \mathbb{N}$ such that $(\frac{1}{c})^n \geq \frac{2}{a}$ and define $C_k = M \cap \bigcap_{r>0} S(c^{-k}, r, M)$. By Lemma 1 $M = C_n$

and therefore $M = \bigcup_{k=2}^n (C_k \setminus C_{k-1}) \cup C_1$. By Lemma 2, we have $p^+(C_1, x) \geq c$ for any $x \in C_1$.

For $k = 2, \dots, n$ define $T_{k,m} = C_k \setminus S(c^{-k+1}, m^{-1}, M)$. Then

$$\bigcup_{m=1}^\infty T_{k,m} = C_k \setminus \bigcap_{m=1}^\infty S(c^{-k+1}, m^{-1}, M) = C_k \setminus C_{k-1}.$$

Since $T_{k,m} \subset C_k$, for each $z \in T_{k,m}$ and $r > 0$, there exist y and t such that $0 < t < \min(r, m^{-1})$, $(y-t, y) \cap M = \emptyset$ and $z \in c^{-k} \ominus (y-t, y)$. Put $J = c^{-k+1} \ominus (y-t, y)$. Then $z \in c^{-1} \ominus J$ and $J \cap T_{k,m} = \emptyset$, since $J \subset S(c^{-k+1}, m^{-1}, M)$.

Thus, for each $z \in T_{k,m}$, we have $z \in \bigcap_{r>0} S(c^{-1}, c^{-k+1}r, T_{k,m})$ and therefore $p^+(T_{k,m}, z) \geq c$ by Lemma 2, which proves our statement. \square

For the sake of brevity, in the following we shall say that E is V -porous at x if $p^-(E, x) > \frac{100}{101}$. The following lemma is easy to prove.

Lemma 3. *Let $E \subset \mathbb{R}$, $x \in \mathbb{R}$ and a natural number p be given such that $x - 10^{-k}$ or $x - 10^{-(k+1)}$ belongs to E for each natural $k > p$. Then E is not V -porous at x .*

The following lemma is an immediate consequence of Proposition.

Lemma 4. *A set $E \subset \mathbb{R}$ is σ -left-porous iff it is σ - V -porous.*

Definition 2. We say that $\mathcal{F} \subset \text{exp } \mathbb{R}$ is a non- σ - V -porosity family if the following conditions hold:

- (a) \mathcal{F} is a nonempty family of nonempty closed sets,
- (b) for each $F \in \mathcal{F}$ and each open set $G \subset \mathbb{R}$ with $F \cap G \neq \emptyset$, there exists $F^* \in \mathcal{F}$ such that $\emptyset \neq F^* \cap G \subset F \cap G$ and F is V -porous at no point of $F^* \cap G$.

We shall need the following lemma which is a special case of [Za 3, Lemma 4.3].

Lemma 5. *Let \mathcal{F} be a non- σ - V -porosity family. Then no set from \mathcal{F} is σ - V -porous.*

3. Proof of theorem

Our theorem stated in Introduction immediately follows from Lemma 7 and Lemma 8 below. To formulate them, we need some notions.

Definition 3. Let $x \in (0, 1)$. As usual, we write $x = 0, a_1a_2\dots$ if $x = \sum_{i=1}^{\infty} a_i 10^{-i}$ and $a_i \in \{0, 1, \dots, 9\}$. The uniqueness of the expansion is obtained using terminating 0's whenever x has two expansions. Let $a \in \{0, 1, \dots, 9\}$ be a digit. The density and the upper density of a in the expansion of x are defined as

$$d(a, x) = \lim_{n \rightarrow \infty} \frac{\#\{k; 1 \leq k \leq n, a_k(x) = a\}}{n},$$

$$\bar{d}(a, x) = \overline{\lim}_{n \rightarrow \infty} \frac{\#\{k; 1 \leq k \leq n, a_k(x) = a\}}{n}.$$

The following easy fact is well known and easy to prove.

Lemma 6. *The function $x \mapsto \bar{d}(a, x)$ is Borel measurable on $(0, 1)$.*

Definition 4. For a natural n and $x \in (0, 1)$ put

$$c(x, n) = \#\{k; n^2 < k \leq (n + 1)^2, a_k(x) = 9\} \quad \text{and}$$

$$e(x, n) = \#\{k; n^2 < k \leq (n + 1)^2, a_k(x) \neq 9\}$$

Let a natural number N , $\varepsilon > 0$, $1 > \alpha > 0$ and digits $a_1, \dots, a_{N^2} \in \{0, 1, \dots, 9\}$ be given. Then we define the set $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ as the set of all $x \in (0, 1)$ for which

$$(1) \quad a_1(x) = a_1, \dots, a_{N^2}(x) = a_{N^2} \quad \text{and}$$

$$(2) \quad 1 - \frac{\varepsilon}{n^\alpha} \leq \frac{c(x, n)}{2n+1} < 1 \quad \text{whenever } n \geq N$$

Lemma 7. Let $0 < \alpha < 1$, $\varepsilon > 0$ and digits $a_1, \dots, a_{N^2} \in \{0, 1, \dots, 9\}$ such that

$$(3) \quad N > \max\left[\left(1 + \varepsilon\right)^{\frac{1}{\alpha}}, \varepsilon^{\frac{1}{\alpha-1}}\right]$$

be given.

Then $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ is a closed set which is not σ -left-porous.

PROOF: Obviously (2) implies that

$$(4) \quad e(x, n) \neq 0 \quad \text{whenever } n \geq N \quad \text{and } x \in A(\alpha, a_1, \dots, a_{N^2}, \varepsilon).$$

Now suppose that $x_n \in A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ and $x_n \rightarrow x$. On account of (4) we easily obtain that

$$(a_1(x_n), a_2(x_n), \dots) \rightarrow (a_1(x), a_2(x), \dots)$$

in the space $\mathbb{N}^{\mathbb{N}}$ and consequently $x \in A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$. Thus we have that $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ is closed.

The condition (2) is equivalent to

$$c(x, n) \in \left[\left(1 - \frac{\varepsilon}{n^\alpha}\right)(2n+1), 2n+1\right) := I_n \quad \text{for } n \geq N.$$

If $n \geq N$, we have by (3)

$$\begin{aligned} \left(1 - \frac{\varepsilon}{n^\alpha}\right)(2n+1) &> \left(1 - \frac{\varepsilon}{1+\varepsilon}\right)(2(1+\varepsilon)^{\frac{1}{\alpha}}) > 2 \quad \text{and} \\ (2n+1) - \left(1 - \frac{\varepsilon}{n^\alpha}\right)(2n+1) &= \frac{\varepsilon}{n^\alpha}(2n+1) > 2\varepsilon n^{1-\alpha} > 2. \end{aligned}$$

Thus we have $I_n \subset (2, 2n+1)$ and $\text{length}(I_n) > 2$ for $n \geq N$; consequently $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon) \neq \emptyset$ and $c(x, n) > 2$ whenever $x \in A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ and $n \geq N$.

Now let \mathcal{F} denote the family of all sets of the form $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ for which (3) holds. By Lemma 4 and Lemma 5 it is sufficient to prove that \mathcal{F} is a non- σ - V -porosity family. To this end suppose that $F = A(\alpha, a_1, \dots, a_{N^2}, \varepsilon) \in \mathcal{F}$ and an open set $G \subset \mathbb{R}$ such that $F \cap G \neq \emptyset$ are given.

Choose an arbitrary $y \in F \cap G$ and find a natural M so large that

$$(5) \quad M > N, \quad M > \left(\frac{\varepsilon}{2}\right)^{\frac{1}{\alpha-1}} \quad \text{and}$$

$$F^* := A(\alpha, a_1, \dots, a_{N^2}, a_{N^2+1}(y), \dots, a_{M^2}(y), \frac{1}{2}\varepsilon) \subset G.$$

Clearly $F^* \subset F$. On account of (3) and (5) we have

$$M > \max\left(\left(1 + \frac{\varepsilon}{2}\right)^{\frac{1}{\alpha}}, \left(\frac{\varepsilon}{2}\right)^{\frac{1}{\alpha-1}}\right)$$

and therefore $F^* \in \mathcal{F}$. Thus it is sufficient to prove that F is V -porous at no point $z \in F^*$. To prove this, fix an arbitrary $z \in F^*$ and consider an arbitrary natural $k > (M + 1)^2$. By Lemma 3 it is sufficient to prove that at least one of the points $z_k^- = z - 10^{-k}$, $z_{k+1}^- = z - 10^{-(k+1)}$ belongs to F . It is easy to see that

$$(6) \quad c(z, n) - 1 \leq c(z_k^-, n) \quad \text{and} \quad c(z, n) - 1 \leq c(z_{k+1}^-, n), \quad \text{for each } n.$$

Since $z \in F^*$, we have $c(z, M) > 0$ (we know even $c(z, M) > 2$) and therefore

$$(7) \quad a_s(z) = a_s(z_k^-) = a_s(z_{k+1}^-) \quad \text{for } s \leq M^2.$$

Now suppose that $x \in \{z_k^-, z_{k+1}^-\}$. Then (7) says that

$$a_s(x) = a_s(z) \quad \text{for } s \leq M^2.$$

For $n \geq M$ the definition of F^* , (6) and (5) yield

$$\frac{c(x, n)}{2n + 1} \geq \frac{c(z, n) - 1}{2n + 1} \geq 1 - \frac{\varepsilon}{2n^\alpha} - \frac{1}{2n + 1} > 1 - \frac{\varepsilon}{n^\alpha}.$$

Thus it is sufficient to establish that, for $x = z_k^-$ or $x = z_{k+1}^-$,

$$(8) \quad e(x, n) \neq 0, \quad \text{for each } n \geq M.$$

To this end suppose that

$$e(z_k^-, n) = 0 \quad \text{for some } n \geq M.$$

Since $c(z, n) \neq 0$, this condition easily implies that

$$\begin{aligned} k &= n^2 + i \quad \text{where } i \in \{1, \dots, 2n\}, \\ a_{n^2+1}(z) &= 0, \dots, a_{n^2+i}(z) = 0 \quad \text{and} \\ a_{n^2+i+1}(z) &= 9, \dots, a_{(n+1)^2}(z) = 9. \end{aligned}$$

Consequently it is easy to see that (8) holds for $x = z_{k+1}^-$. □

Lemma 8. *If $\frac{1}{2} < \alpha < 1$, then the set $A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$ from Lemma 7 is right porous.*

PROOF: Choose an arbitrary $x \in A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$.

For each natural n , let m_n be the maximum of those natural i , for which there exist natural numbers u, v such that

$$(9) \quad n^2 \leq u < v \leq (n+1)^2, \quad a_s(x) = 9 \quad \text{for each } u < s \leq v$$

and $v - u = i$. It is easy to see that

$$(10) \quad \begin{aligned} 2n+1 - e(x, n) &= c(x, n) \leq m_n(e(x, n) + 1) \quad \text{and consequently} \\ m_n &\geq \frac{2n+1 - e(x, n)}{e(x, n) + 1}. \end{aligned}$$

On account of (2) we have that

$$e(x, n) \leq \frac{\varepsilon(2n+1)}{n^\alpha} \quad \text{for } n \geq N$$

and therefore (10) implies that there exists $c > 0$ and a natural n_0 such that

$$(11) \quad m_n \geq cn^\alpha \quad \text{for all } n \geq n_0.$$

Now, for each n , choose u_n, v_n such that

$$v_n - u_n = m_n \quad \text{and (9) holds for } u = u_n, v = v_n.$$

Put

$$y_n = x + 10^{-v_n} \quad \text{and} \quad z_n = x + 10^{-v_n+1}.$$

It is easy to see that, for each $t \in (y_n, z_n)$, we have

$$a_s(t) = 0, \quad \text{for each } u_n < s \leq v_n - 1$$

and therefore

$$(12) \quad c(t, n) \leq 2n+1 - (m_n - 1).$$

If n is so big that $n > n_0$, $n > N$ and $2n+2 - cn^\alpha < (2n+1)(1 - \frac{\varepsilon}{n^\alpha})$, we have by (12) and (11)

$$c(t, n) \leq 2n+2 - cn^\alpha < (2n+1)(1 - \frac{\varepsilon}{n^\alpha}).$$

Thus we obtain by (2) that $t \notin A(\alpha, a_1, \dots, a_{N^2}, \varepsilon)$. Consequently

$$p^+ \left(A(\alpha, a_1, \dots, a_{N^2}, \varepsilon), x \right) \geq \overline{\lim}_{n \rightarrow \infty} \frac{10^{-(v_n-1)} - 10^{-v_n}}{10^{-v_n+1}} = \frac{9}{10}.$$

REFERENCES

- [F] Foran J., *Continuous functions need not have σ -porous graphs*, Real Anal. Exchange **11** (1985–86), 194–203.
- [Za 1] Zajíček L., *On σ -porous sets and Borel sets*, Topology Appl. **33** (1989), 99–103.
- [Za 2] ———, *Sets of σ -porosity and sets of σ -porosity (q)*, Časopis Pěst. Mat. **101** (1976), 350–359.
- [Za 3] ———, *Porosity and σ -porosity*, Real Anal. Exchange **13** (1987–88), 314–350.
- [E-H-S] Evans M.J., Humke P.D., Saxe K., *A symmetric porosity conjecture of L. Zajíček*, Real Anal. Exchange **17** (1991–92), 258–271.

DEPARTMENT OF MATHEMATICAL ANALYSIS, CHARLES UNIVERSITY, SOKOLOVSKÁ 83,
186 00 PRAHA 8, CZECH REPUBLIC

(Received January 19, 1994)