

Ram U. Verma

On monotone nonlinear variational inequality problems

Commentationes Mathematicae Universitatis Carolinae, Vol. 39 (1998), No. 1, 91--98

Persistent URL: <http://dml.cz/dmlcz/118988>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 1998

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On monotone nonlinear variational inequality problems

RAM U. VERMA

Abstract. The solvability of a class of monotone nonlinear variational inequality problems in a reflexive Banach space setting is presented.

Keywords: nonlinear variational inequality problems, p -monotone and p -Lipschitzian operators, KKM mappings

Classification: 47H15

1. Introduction

General theory of monotone variational inequalities has been applied to various problems in applied mathematics, physics, engineering sciences, and others. A closely associated notion of the complementarity involves several problems in mathematical programming, game theory, economics, and mechanics. For more details on general variational inequalities, we advise to consult [1], [4]–[14].

Let X be a reflexive real Banach space with dual X^* and $[w, x]$ denote a continuous duality pairing between the elements w in X^* and x in X . Let K be a nonempty closed convex subset of X . Here we present the solvability of a class of monotone nonlinear variational inequality (MNVI) problems: Determine an element x in K for a given w in X^* such that

$$(1.1) \quad [Sx - Tx - w, v - x] + f(v) - f(x) \geq 0 \text{ for all } v \in K,$$

where $S, T : K \rightarrow X^*$ are nonlinear operators, and $f : X \rightarrow (-\infty, +\infty]$ is convex lower semicontinuous functional with $f \not\equiv \infty$. Here S and T are, respectively, p -monotone and p -Lipschitz continuous (or p -Lipschitzian).

Next, we recall some definitions needed for the work at hand.

Definition 1.1. An operator $S : K \rightarrow X^*$ is said to be p -monotone if, for all $u, v \in K$, there exist constants $r > 0$ and $p > 1$ such that

$$(1.2) \quad [Su - Sv, u - v] \geq r\|u - v\|^p.$$

The inequality (1.2) implies that S is strictly monotone and coercive for $p > 1$, S is strongly monotone for $p = 2$, and S is uniformly monotone for $p \geq 2$.

Definition 1.2. An operator $T : K \rightarrow X^*$ is called *p-Lipschitz continuous* (or *p-Lipschitzian*) if, for all $u, v \in K$, there exist constants $k > 0$ and $p > 1$ such that

$$(1.3) \quad [Tu - Tv, u - v] \leq k\|u - v\|^p.$$

Let us consider an example of *p-Lipschitzian* operators in the context of generalized pseudocontractions — a mild generalization of the pseudocontractions introduced by Browder and Petryshyn [2] — in a Hilbert space H . Generalized pseudocontractions are more general than Lipschitzian operators and unify certain classes of operators.

Definition 1.3. An operator $T : H \rightarrow H$ is said to be a *generalized pseudocontraction* if, for all $u, v \in H$, there exists a constant $k > 0$ such that

$$(1.4) \quad \|Tu - Tv\|^2 \leq k^2\|u - v\|^2 + \|Tu - Tv - k(u - v)\|^2.$$

This is equivalent to

$$(1.5) \quad \langle Tx - Ty, x - y \rangle \leq k\|x - y\|^2,$$

where $T : H \rightarrow H$ is 2-Lipschitzian.

Example 1.4 ([JY]). Let K be a closed convex subset of a real Hilbert space H , and let $T : K \rightarrow K$ be hemicontinuous and 2-Lipschitzian with a constant $0 < k < 1$. Then T has a unique fixed point in K .

Definition 1.5. A multivalued mapping $F : X \rightarrow P(X)$ is called the *KKM* mapping if, for every finite subset $\{u_1, u_2, \dots, u_n\}$ of X , $\text{conv}\{u_1, u_2, \dots, u_n\}$ is contained in $\bigcup_{i=1}^n F(u_i)$, where $\text{conv}\{A\}$ is the convex hull of set A and $P(X)$ denotes the power set of X .

Before we present our main results, we need to recall some auxiliary results [3].

Lemma 1.6 ([3, Theorem 4]). *Let Y be a convex set in a topological vector space X , and let K be a nonempty subset of Y . For all $x \in K$, let $F(x)$ be a relatively closed subset of Y such that the convex hull of every finite subset $\{x_1, x_2, \dots, x_n\}$ of K is contained in the corresponding union $\bigcup_{i=1}^n F(x_i)$. If there is a nonempty subset K_0 of K such that the intersection $\bigcap_{x \in K_0} F(x)$ is compact and K_0 is contained in a compact convex subset of Y , then $\bigcap_{x \in K} F(x) \neq \emptyset$.*

Lemma 1.7 ([3, Corollary 1]). *Let K be a nonempty set in a topological vector space X . Let $F : K \rightarrow P(K)$ be a KKM mapping from K into the power set of K . If $F(u)$ is closed in X for all $u \in K$ and is compact for at least one $u \in K$, then $\bigcap_{u \in K} F(u) \neq \emptyset$.*

We note that in Lemma 1.6 the hypothesis “ $\bigcap_{x \in K_0} F(x)$ is compact” does not rule out the possibility that it may be empty. However, the conclusion “ $\bigcap_{x \in K} F(x) \neq \emptyset$ ” does imply that $\bigcap_{x \in K_0} F(x)$ is nonempty. The compactness condition in Lemma 1.7 is relaxed in Lemma 1.6.

2. The main results

Theorem 2.1. *Let K be a convex subset of a reflexive real Banach space X with dual X^* and $0 \in K$. Let $S : K \rightarrow X^*$ be hemicontinuous and p -monotone and let $T : K \rightarrow X^*$ be hemicontinuous and p -Lipschitz continuous. Let us further assume that $f : K \rightarrow (-\infty, \infty]$ is a convex functional with $f(0) = 0$, $f(u) > 0$ and $f \not\equiv \infty$. Then, for a given $w \in X^*$, an element u in K is a solution of the MNVI problem*

$$(2.1) \quad [Su - Tu - w, v - u] + f(v) - f(u) \geq 0 \text{ for all } v \in K$$

iff u is a solution of a new MNVI problem

$$(2.2) \quad [Sv - Tv - w, v - u] + f(v) - f(u) \geq c\|v - u\|^p \text{ for all } v \in K,$$

where $c = r - k > 0$ and $p > 1$. Here r is the p -monotonicity constant of S and k is the p -Lipschitz continuity constant of T .

When S and T are monotone and antimonotone, respectively, and $w = 0$, Theorem 2.1 reduces to [8, Lemma 1].

Corollary 2.2. *Let K be a nonempty convex subset of X and let $S : K \rightarrow X^*$ and $T : K \rightarrow X^*$ both be hemicontinuous, and be monotone and antimonotone, respectively. Let f be convex with $f \not\equiv \infty$. Then the following variational inequality problems are equivalent:*

$$(2.3) \quad u \in K : [Su - Tu, v - u] + f(v) - f(u) \geq 0 \text{ for all } v \in K;$$

$$(2.4) \quad u \in K : [Sv - Tv, v - u] + f(v) - f(u) \geq 0 \text{ for all } v \in K.$$

For $T = 0$ and f an indicator functional (that is, $f = 0$ on K and $f = \infty$ off K), Theorem 2.1 reduces to

Corollary 2.3. *Let K be a nonempty closed convex subset of a reflexive real Banach space X with dual X^* and let $S : K \rightarrow X^*$ be hemicontinuous and p -monotone. Then the MNVI problem*

$$(2.5) \quad u \in K : [Su - w, v - u] \geq 0 \text{ for all } v \in K,$$

has a unique solution iff the MNVI problem

$$(2.6) \quad u \in K : [Sv - w, v - u] \geq r\|v - u\|^p \text{ for all } v \in K,$$

has a unique solution for each $w \in X^$.*

PROOF OF THEOREM 2.1: Suppose that (2.1) holds. Since S is p -monotone and T is p -Lipschitz continuous, this implies that

$$[(S - T)v - (S - T)u, v - u] \geq c\|v - u\|^p$$

or

$$\begin{aligned} [(S - T)v, v - u] &\geq c\|v - u\|^p + [(S - T)u, v - u] \\ &\geq c\|v - u\|^p + [w, v - u] + f(u) - f(v). \end{aligned}$$

This implies that

$$[(S - T)v - w, v - u] + f(v) - f(u) \geq c\|v - u\|^p.$$

Conversely, if (2.2) holds, then by choosing an element v with $f(v) < +\infty$, we find that $f(u)$ is finite. Let x be an element of K such that $v_t = (1 - t)u + tx$ satisfies (2.2) for $0 < t < 1$. Then, it follows that $v_t - u = t(x - u)$ and, as a result, we find that

$$[(S - T)v_t - w, v_t - u] + f(v_t) - f(u) \geq c\|v_t - u\|^p$$

or

$$t[(S - T)v_t - w, x - u] + f((1 - t)u + tx) - f(u) \geq c\|v_t - u\|^p.$$

Since f is convex, this implies that

$$\begin{aligned} t[(S - T)v_t - w, x - u] &\geq c\|v_t - u\|^p + f(u) - (1 - t)f(u) - tf(x) \\ &= c\|t(x - u)\|^p + t(f(u) - f(x)). \end{aligned}$$

Thus, given that $t > 0$, we find

$$[(S - T)v_t - w, x - u] + f(x) - f(u) \geq ct^{p-1}\|x - u\|^p.$$

Since the hemicontinuity of S and T implies the hemicontinuity of $S - T$, we find that $(S - T)v_t$ converges weakly to $(S - T)u$ in X^* as $t \rightarrow 0$. Hence, we obtain

$$[(S - T)u - w, x - u] + f(x) - f(u) \geq 0 \text{ for all } x \in K,$$

that is, the variational inequality (2.1) holds. \square

Theorem 2.4. *Let K be a nonempty closed convex subset of a reflexive real Banach space X with $0 \in K$. Let $S : K \rightarrow X^*$ be hemicontinuous and p -monotone with constant $r > 0$, $T : K \rightarrow X^*$ be hemicontinuous and p -Lipschitz continuous with constant $k > 0$, and $f : X \rightarrow (-\infty, +\infty]$ be convex lower semicontinuous with $f \not\equiv \infty$. Then the MNVI problem*

$$(2.7) \quad u \in K : [Su - Tu - w, v - u] + f(v) - f(u) \geq 0 \text{ for all } v \in K$$

has a unique solution for each $w \in X^*$.

For $w = 0$, S strictly monotone, T strictly antimonotone, and K bounded, Theorem 2.4 reduces to [8, Theorem 3].

Corollary 2.5. *Let K be a nonempty bounded closed convex subset of X , and $S, T : K \rightarrow X^*$ both be hemicontinuous and be strictly monotone and antimonotone, respectively. Let $f : X \rightarrow (-\infty, +\infty]$ be convex lower semicontinuous with $f \not\equiv \infty$. Then the variational inequality problem*

$$(2.8) \quad u \in K : [Su - Tu, v - u] + f(v) - f(u) \geq 0 \text{ for all } v \in K$$

has a unique solution.

When $T = 0$ and f is an indicator functional on K (that is, $f = 0$ on K and $f = \infty$ off K), Theorem 2.4 reduces to [5, Theorem 2].

Corollary 2.6. *Let X be a reflexive real Banach space with dual X^* and K be a nonempty closed convex subset X . Let $S : K \rightarrow X^*$ be hemicontinuous and p -monotone. Then the variational inequality problem*

$$(2.9) \quad u \in K : [Su - w, v - u] \geq 0 \text{ for all } v \in K$$

has a unique solution for each $w \in X^*$.

PROOF OF THEOREM 2.4: We first prove the existence of the solution of the MNVI problem (2.7). Let us define the multivalued mappings $F, G : K \rightarrow P(K)$ by

$$F(v) = \{u \in K : [Su - Tu - w, v - u] + f(v) - f(u) \geq 0\} \text{ for all } v \in K$$

and

$$G(v) = \{u \in K : [Sv - Tv - w, v - u] + f(v) - f(u) \geq c\|v - u\|^p\} \text{ for all } v \in K,$$

respectively. We show by a contradiction approach that F is a KKM mapping.

Assume $\{v_1, v_2, \dots, v_n\}$ is in K , $\sum_{i=1}^n t_i = 1$, $t_i > 0$ and $v = \sum_{i=1}^n t_i v_i$ is not in

$$\bigcup_{i=1}^n F(v_i). \text{ Then for } u = v,$$

$$[Su - Tu - w, v_i - u] < f(u) - f(v_i) \text{ for any } i = 1, \dots, n.$$

Thus, we find

$$\begin{aligned}
 0 &= [Su - Tu - w, v - u] = [Su - Tu - w, \sum_{i=1}^n t_i v_i - u] \\
 &= \sum_{i=1}^n t_i [Su - Tu - w, v_i - u] < \sum_{i=1}^n t_i (f(u) - f(v_i)) \\
 &= f(u) - \sum_{i=1}^n t_i f(v_i) \leq f(u) - f(\sum_{i=1}^n t_i v_i) \\
 &= f(u) - f(v) = 0,
 \end{aligned}$$

a contradiction. This implies that $\text{conv}\{v_1, v_2, \dots, v_n\}$ is contained in $\bigcup_{i=1}^n F(v_i)$.

Next, to show $F(v) \subset G(v)$ for all $v \in K$, let u belong to $F(v)$. Then using the p -monotonicity of S and p -Lipschitz continuity of T , we obtain

$$[(S - T)v - (S - T)u, v - u] \geq c\|v - u\|^p.$$

Thus,

$$\begin{aligned}
 [(S - T)v, v - u] &\geq c\|v - u\|^p + [(S - T)u, v - u] \quad \text{or} \\
 [(S - T)v - w, v - u] &\geq c\|v - u\|^p + [(S - T)u - w, v - u] \\
 &\geq c\|v - u\|^p + f(u) - f(v) \quad \text{or} \\
 [(S - T)v - w, v - u] + f(v) - f(u) &\geq c\|v - u\|^p \quad \text{for all } v \in K.
 \end{aligned}$$

This implies that u belongs to $G(v)$ and, consequently, G is a KKM mapping on K . Hence, by Theorem 2.1, we find $\bigcap_{v \in K} F(v) = \bigcap_{v \in K} G(v)$.

Since f is lower semicontinuous and the duality pairing $[\cdot, \cdot]$ is continuous, it follows that $G(v)$ is closed for all $v \in K$. Clearly, K is a weakly compact set in X with weak topology and, as a result, $G(v)$ is weakly compact in K since $G(v)$ is contained in K for each $v \in K$. Now, by Lemma 1.7, we find

$$\bigcap_{v \in K} F(v) = \bigcap_{v \in K} G(v) \neq \emptyset.$$

Hence, there exists an element u_0 in K such that

$$[Su_0 - Tu_0 - w, v - u_0] + f(v) - f(u_0) \geq 0 \quad \text{for all } v \in K.$$

To show the uniqueness of the solution, let x_1, x_2 be two solutions of the MNVI problem (2.7), that is,

$$(2.10) \quad [Sx_1 - Tx_1 - w, v - x_1] + f(v) - f(x_1) \geq 0 \quad \text{for all } v \in K,$$

and

$$(2.11) \quad [Sx_2 - Tx_2 - w, v - x_2] + f(v) - f(x_2) \geq 0 \quad \text{for all } v \in K.$$

Setting $v = x_2$ in (2.10) and $v = x_1$ in (2.11), and adding, we obtain

$$-[Sx_1 - Tx_1 - w, x_1 - x_2] + [Sx_2 - Tx_2 - w, x_1 - x_2] \geq 0,$$

or

$$-[Sx_1 - Sx_2, x_1 - x_2] + [Tx_1 - Tx_2, x_1 - x_2] \geq 0,$$

or

$$[Sx_1 - Sx_2, x_1 - x_2] \leq [Tx_1 - Tx_2, x_1 - x_2].$$

Since S is p -monotone with constant $r > 0$ and T is p -Lipschitz continuous with constant $k > 0$, this implies that

$$r\|x_1 - x_2\|^p \leq [Sx_1 - Sx_2, x_1 - x_2] \leq [Tx_1 - Tx_2, x_1 - x_2] \leq k\|x_1 - x_2\|^p.$$

It follows that

$$(r - k)\|x_1 - x_2\|^p \leq 0.$$

Since $r - k > 0$, we find that $x_1 = x_2$. This completes the proof. \square

Acknowledgment. The author wishes to express his sincere appreciation to the referee for some valuable suggestions leading to the revised version.

REFERENCES

- [1] Browder F.E., *On the unification of the calculus of variations and the theory of monotone nonlinear operators in Banach spaces*, Proc. Nat. Acad. Sci. U.S.A. **56** (1966), 419–425.
- [2] Browder F.E., Petryshyn W.V., *Construction of fixed points of nonlinear mappings in Hilbert spaces*, J. Math. Anal. Appl. **20** (1967), 197–228.
- [3] Fan K., *Some properties of convex sets related to fixed point theorems*, Math. Annal. **266** (1984), 519–537.
- [4] Glowinski R., *Numerical Methods for Nonlinear Variational Problems*, Springer-Verlag, New York, 1984.
- [5] Goeleven D., Motreanu D., *Eigenvalue and dynamic problems for variational and hemivariational inequalities*, Comm. Appl. Nonlinear Anal. **3** (4) (1996), 1–21.
- [6] Noor M.A., *Mixed variational inequalities*, Appl. Math. Lett. **3** (1990), 73–75.
- [7] Noor M.A., *General auxiliary principle for variational inequalities*, PanAmerican Math. J. **4** (1) (1994), 27–44.
- [8] Siddiqi A.H., Ansari Q.H., Kazmi K.R., *On nonlinear variational inequalities*, Indian J. Pure Appl. Math. **25** (9) (1994), 969–973.
- [9] Szulkin A., *Positive solutions of variational inequalities: A degree-theoretic approach*, J. Diff. Equ. **57** (1985), 90–111.
- [10] Verma R.U., *Iterative algorithms for variational inequalities and associated nonlinear equations involving relaxed Lipschitz operators*, Appl. Math. Lett. **9** (4) (1996), 61–63.
- [11] Verma R.U., *Generalized variational inequalities involving multivalued relaxed monotone operators*, Appl. Math. Lett., to appear.

- [12] Verma R.U., *Nonlinear variational and constrained hemi-variational inequalities involving relaxed operators*, Z. Angew. Math. Mech. **77** (1997), 387–391.
- [13] Yao J.-C., *Applications of variational inequalities to nonlinear analysis*, Appl. Math. Lett. **4** (1991), 89–92.
- [14] Zeidler E., *Nonlinear Functional Analysis and its Applications IV*, Springer-Verlag, New York, 1988.

INTERNATIONAL PUBLICATIONS, 12046 COED DRIVE, ORLANDO, FLORIDA 32826, USA

ISTITUTO PER LA RICERCA DI BASE, DIVISION OF MATHEMATICS, I-86075 MONTERODUNI (IS), MOLISE, ITALY

(Received February 3, 1997, revised August 6, 1997)