

Eva Murtinová

A β -normal Tychonoff space which is not normal

Commentationes Mathematicae Universitatis Carolinae, Vol. 43 (2002), No. 1, 159--164

Persistent URL: <http://dml.cz/dmlcz/119308>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A β -normal Tychonoff space which is not normal

EVA MURTIŇOVÁ

Abstract. α -normality and β -normality are properties generalizing normality of topological spaces. They consist in separating dense subsets of closed disjoint sets. We construct an example of a Tychonoff β -normal non-normal space and an example of a Hausdorff α -normal non-regular space.

Keywords: α -normal, β -normal, closed unbounded

Classification: 54A05, 54D15, 03E75

The notions of α and β -normality have been introduced recently by A.V. Arhangel'skii and L. Ludwig in [AL].

Definition. A topological space X is called α -normal (β -normal) whenever for each pair of closed disjoint subsets $A, B \subset X$ there are open sets $U, V \subset X$ such that $\overline{A \cap U} = A$, $\overline{B \cap V} = B$ and $U \cap V = \emptyset$ ($\overline{U} \cap \overline{V} = \emptyset$, respectively).

Arhangel'skii and Ludwig presented several examples of Tychonoff not α -normal and of α -normal not β -normal spaces. They made observations we recall in Propositions 1 and 2.

Proposition 1. *A space X is β -normal if and only if for every A closed and U open, $A \subset U \subset X$, there exists an open $V \subset X$ such that $\overline{A \cap V} = A$ and $\overline{V} \subset U$.*

Proposition 2.

- (a) *Every α -normal T_1 -space is Hausdorff.*
- (b) *Every β -normal T_1 -space is regular.*

This leads naturally to the following questions:

1. Does there exist a Hausdorff α -normal non-regular space? ([AL], Question 1).
2. Does there exist a regular β -normal non-Tychonoff space? We shall return to this topic in Remark 4.
3. Does there exist a Tychonoff β -normal not normal space? ([AL], Question 4).

The aim of the present paper is to answer them in the affirmative. Another ZFC example of a Hausdorff α -normal non-regular space was found independently

This research was supported by grant GAUK 270/2001.

by D. Burke. And there is a consistent example of a Tychonoff β -normal not normal space due to Ludwig and Szeptycki [LS].

Let us observe that a positive answer to the third question yields a positive answer to the first one:

Proposition 3. *Suppose X is a regular β -normal not normal space. Let A, B be closed disjoint subsets of X that cannot be separated by open neighborhoods. Then the quotient space $Y = X/A$ obtained by identifying A with a point is Hausdorff, α -normal and not regular.*

PROOF: Let $q : X \rightarrow Y$ be the canonical quotient map, $\{a\} = q[A]$. It is easy to see that Y is a Hausdorff space and that $q[B]$ and a cannot be separated.

Choose closed disjoint subsets F, H of Y . If $a \notin F \cup H$ then dense parts of F and H can be separated using α -normality of X . Assume that $a \in F$, i.e. $A \subset q^{-1}[F]$ and $A \cap \overline{q^{-1}[H]} = \emptyset$. The space X is β -normal, hence there is O open in X such that $\overline{q^{-1}[H]} \cap O = q^{-1}[H]$ and $q^{-1}[F] \cap \overline{O} = \emptyset$. Put $U = q[O]$, $V = Y \setminus \overline{U}$. As $\overline{O} \cap A = \emptyset$, U is an open subset of Y and $\overline{U} = q[\overline{O}]$. But $H \cap \overline{U} = H$, $V \supset F$. \square

Note that the previous argument is not valid, if α -normality is assumed instead of β -normality.

We give an affirmative answer to the third question in ZFC now.

Example 1. A β -normal Tychonoff space which is not normal.

We shall keep the notation $S = \{\alpha < \omega_2 : \text{cf } \alpha = \omega_1\}$.

Consider the set

$$X = \{(\alpha, \beta) : \beta \leq \alpha \leq \omega_2 \ \& \ (\alpha, \beta) \neq (\omega_2, \omega_2)\}$$

and its partition into

$$\begin{aligned} A &= \{(\alpha, \alpha) : \alpha < \omega_2\}, \\ B &= \{(\omega_2, \beta) : \beta < \omega_2\}, \\ D &= \{(\alpha, \beta) : \beta < \alpha < \omega_2\}. \end{aligned}$$

Topologize X in this way: let each $(\alpha, \beta) \in D$ be isolated, let an open base in $(\alpha, \alpha) \in A$ consist of all sets of the type

$$\{(\gamma, \gamma) : \alpha_0 < \gamma \leq \alpha\} \cup \bigcup \{\{\gamma\} \times C_\gamma : \alpha_0 < \gamma \leq \alpha \ \& \ \gamma \in S\},$$

where $\alpha_0 < \alpha$ and every C_γ is a closed unbounded (club) subset of γ , and let the base in $(\omega_2, \beta) \in B$ consist of all the sets

$$\{(\alpha, \gamma) : \beta_0 < \gamma \leq \beta \ \& \ \alpha_\gamma < \alpha \leq \omega_2\},$$

where $\beta_0 < \beta$ and $\beta \leq \alpha_\gamma < \omega_2$ for each γ .

All above defined basic open neighborhoods are closed, i.e. X is a zero-dimensional (in particular, Tychonoff) space.

Let us disprove normality of X . Consider closed disjoint sets A, B and an open O containing A . Then $\overline{O} \cap B \neq \emptyset$. Indeed, S is a stationary subset of ω_2 and as $\{(\alpha, \alpha) : \alpha \in S\} \subset A \subset O$, for each $\alpha \in S$ there is a C_α club in α such that $\{\alpha\} \times C_\alpha \subset O$. But by Fodor's Pressing Down Lemma, there is an unbounded $R \subset S$ and a $\beta \in \omega_2$ such that $(\forall \alpha \in R) \beta \in C_\alpha$. It follows that $(\omega_2, \beta) \in \overline{O}$.

Using Proposition 1 we shall prove β -normality of X .

Assume that $F \subset O \subset X$, F is closed, O open. We are looking for an open U such that $\overline{F \cap U} = F$ and $\overline{U} \subset O$. Note that the only interesting cases are when $F \subset A$ or $F \subset B$. Indeed, for a general F , denote $F_A = F \cap A$, $F_B = F \cap B$, $F_D = F \cap D$ (F_A and F_B are closed). If there are open sets U_1, U_2 such that

$$\begin{aligned} \overline{F_A \cap U_1} &= F_A \quad \& \quad \overline{U_1} \subset O, \\ \overline{F_B \cap U_2} &= F_B \quad \& \quad \overline{U_2} \subset O, \end{aligned}$$

then $U = U_1 \cup U_2 \cup F_D$ is open, $\overline{F \cap U} \supset (\overline{F_A \cap U_1}) \cup (\overline{F_B \cap U_2}) \cup \overline{F_D} = F$, $\overline{U} = \overline{U_1} \cup \overline{U_2} \cup \overline{F_D} \subset O \cup F = O$.

As $A \simeq B \simeq \omega_2$, $A \cup D$ and $B \cup D$ are (open) normal subspaces of X . Therefore we can assume that $O = X \setminus B$ in case of $F \subset A$ and $O = X \setminus A$ if $F \subset B$. Moreover, for each $\beta_0 < \omega_2$

$$\{(\alpha, \beta) \in X : \beta \leq \beta_0\}$$

is a clopen normal subspace. It follows that if $F \subset A$ ($F \subset B$, respectively) and $|F| \leq \aleph_1$, we are done.

Let $F \subset A$ (or $F \subset B$) be unbounded (i.e. $|F| = \aleph_2$). The set $F^{is} = F \setminus F'$ is open dense in F , so there is an open subset V of A (of B , respectively) such that $F \cap V = F^{is}$. As $F' \cap V = \emptyset$, V is isomorphic to a nonstationary set in ω_2 .

Suppose that $F \subset A$. The set $N = \{\alpha \in S : (\alpha, \alpha) \in V\}$ is nonstationary. Hence there is a nondecreasing regressive function $f : N \rightarrow \omega_2$ such that

$$\forall \beta < \omega_2 \quad |\{\alpha \in N : f(\alpha) = \beta\}| \leq \aleph_1.$$

For each $\alpha \in N$ put $C_\alpha = \{\gamma : f(\alpha) < \gamma < \alpha\}$. Thus

$$U = V \cup \bigcup \{\{\alpha\} \times C_\alpha : \alpha \in N\}$$

is an open subset of X , $\overline{F \cap U} = \overline{F \cap V} = \overline{F^{is}} = F$ and by the choice of f , $\overline{U} = \overline{V} \cup \bigcup \{\{\alpha\} \times C_\alpha : \alpha \in N\}$ does not intersect B , i.e. $\overline{U} \subset O$.

Now, assume that $F \subset B$. The set $C = \omega_2 \setminus \{\beta : (\omega_2, \beta) \in V\}$ is club in ω_2 , $M = S \setminus C' = \{\alpha \in S : \sup(C \cap \alpha) < \alpha\}$ is nonstationary. Fix a regressive nondecreasing function $f : M \rightarrow \omega_2$ such that

$$\forall \beta < \omega_2 \quad |\{\alpha \in M : f(\alpha) = \beta\}| \leq \aleph_1.$$

For $\alpha \in S$ define

$$C_\alpha = \begin{cases} C \cap \alpha, & \text{if } \alpha \in C', \\ \{\gamma : f(\alpha) < \gamma < \alpha\}, & \text{otherwise (i.e. if } \alpha \in M). \end{cases}$$

Thus, $G = A \cup \bigcup \{\{\alpha\} \times C_\alpha : \alpha \in S\}$ is an open neighborhood of A and $\overline{G} \cap V = \emptyset$. Indeed, for any $(\omega_2, \beta) \in V$, $\{\alpha \in S : \beta \in C_\alpha\}$ is bounded: the set is equal to

$$\{\alpha \in S \cap C' : \beta \in C_\alpha\} \cup \{\alpha \in M : f(\alpha) < \beta\},$$

while the first summand is empty and the second one is bounded.

Put $U = X \setminus \overline{G}$. Clearly, U is open and $F = \overline{F \cap V} \subset \overline{F \setminus \overline{G}} = \overline{F \cap U}$. Moreover, $\overline{U} \cap A = (\overline{X \setminus \overline{G}}) \cap A = \emptyset$. Hence $\overline{U} \subset O$. \square

Remark 4. A construction of F.B. Jones that produces a non-Tychonoff space Y starting from a non-normal space X (see [Jo]) was used by Arhangel'skii and Ludwig to find, under CH, an example of a non-Tychonoff α -normal regular space ([AL, Example 3.3]). It is not hard to prove that if X is regular and β -normal, then so is Y . Hence, starting with X from Example 1, Y is a T_3 , non- $T_{3\frac{1}{2}}$ β -normal space.

Let us turn attention to the first question in a particular case.

Proposition 5. *Every first countable Hausdorff α -normal space is regular.*

PROOF: We shall assume that X is a first countable Hausdorff non-regular space and prove that it is not α -normal.

There is an $x \in X$ and a closed $F \subset X$ such that $x \notin F$ but x, F cannot be separated by disjoint open sets. Let $\{O_n : n \in \omega\}$ be an open base in x , $O_n \supset O_{n+1} (\forall n)$.

Pick, inductively, $x_n \in \overline{O_n} \cap F$; as X is Hausdorff, at each step we can assume that $x_n \notin \overline{O_{n+1}}$. Note that, then,

$$x_n \in \overline{O_m} \Leftrightarrow m \leq n.$$

Define $A = \{x_n : n \in \omega\}$. Then $A' = \emptyset$. Indeed, for a $y \neq x$ there is an open set U and $m \in \omega$ such that $y \in U$ and $U \cap \overline{O_m} = \emptyset$. Therefore $U \cap A \subset \{x_0, \dots, x_{m-1}\}$ and $y \notin A'$.

We conclude that A is closed and has no proper dense subset, while $x \notin A$. As $\{x\}$ and A cannot be separated, they refute α -normality of X . \square

The space from the following example is first countable at each point except one. In fact, cardinality of local basis in $\omega_1 \in X$ is equal to the character of Club filter on ω_1 . In particular, $\aleph_2 \leq \mathfrak{b}_{\omega_1} \leq \chi(\omega_1, X) \leq \mathfrak{d}_{\omega_1}$, see, e.g., [BS] (5.23–5.25).

Example 2. An α -normal Hausdorff space which is not regular.

Let X denote the space with underlying set $\omega_1 + 1$ and topology τ such that: ω_1 with ordinal topology is an open subspace and a base in the point ω_1 consists of all sets of the type

$$O_C = \{\omega_1\} \cup \{\alpha + 1 : \alpha \in C\},$$

where $C \subset \omega_1$ is closed unbounded.

The topology τ is stronger than the order topology on $\omega_1 + 1$, in particular, X is Hausdorff.

The space is not regular: consider club $C \subset \{\alpha < \omega_1 : \alpha \text{ is a limit ordinal}\}$. Then $O_C \cap C = \emptyset$, therefore $\omega_1 \notin \overline{C}$ and C is a closed subset of X . But for every club D ,

$$\overline{O_D} \cap C \supset D' \cap C \neq \emptyset,$$

hence ω_1 and C cannot be separated.

It remains to prove that X is α -normal. Pick closed disjoint $A, B \subset X$. If $A, B \subset \omega_1$ then they can be separated in the open normal subspace ω_1 . Let $B = C \cup \{\omega_1\}$ with $C \subset \omega_1$. It is easy to separate A and B , if A is bounded in ω_1 .

Suppose A is unbounded. This implies that C is bounded, in particular $\omega_1 \notin \overline{C}$. Fix open disjoint $U_1, V_1 \subset \omega_1$ such that $A \subset U_1, C \subset V_1$. We shall separate ω_1 from $A^{is} = A \setminus A'$.

Let $A = \{a_\alpha : \alpha < \omega_1\}$ be an increasing enumeration. In this notation $A^{is} = \{a_0\} \cup \{a_{\alpha+1} : \alpha < \omega_1\}$ and it is dense in A . Put

$$\begin{aligned} G_{-1} &= \langle 0, a_0 \rangle, \\ G_\alpha &= \begin{cases} \{a_{\alpha+1}\}, & \text{if } a_{\alpha+1} = a_\alpha + 1, \\ (a_\alpha + 1, a_{\alpha+1}), & \text{if } a_{\alpha+1} > a_\alpha + 1, \end{cases} \\ G &= \bigcup_{-1 \leq \alpha < \omega_1} G_\alpha. \end{aligned}$$

Then G is an open neighborhood of A^{is} .

As $\omega_1 \notin \overline{A} = A$, there is a club D such that $O_D \cap A = \emptyset$, or, equivalently, $\beta \in D \Rightarrow \beta + 1 \notin A$. Moreover, note that $a_\alpha + 1 \in G$ iff $a_{\alpha+1} = a_\alpha + 1$. It is now easy to check that

$$\begin{aligned} G \cap O_{D \cap A} &= \{a_\alpha + 1 : \alpha < \omega_1 \ \& \ a_\alpha \in D \ \& \ a_{\alpha+1} = a_\alpha + 1\} \\ &= \{\beta + 1 : \beta \in D \cap A \ \& \ \beta + 1 \in A\} = \emptyset. \end{aligned}$$

Put $U = U_1 \cap G$, $V = V_1 \cup O_{D \cap A}$. We have shown that $U \cap V = \emptyset$, $B \subset V$ and $\overline{U \cap A} = A$. \square

Referring to Proposition 2, we have constructed an example of an α -normal, not β -normal space.

REFERENCES

- [AL] Arhangel'skii A. V., Ludwig L., *On α -normal and β -normal spaces*, Comment. Math. Univ. Carolinae **42.3** (2001), 507–519.
- [BS] Balcar B., Simon P., *Disjoint refinement*, Handbook of Boolean Algebras (J.D. Monk, R. Bonnet, ed.), Elsevier Science Publishers B.V., 1989, pp. 333–386.
- [Jo] Jones F.B., *Hereditarily separable, non-completely regular spaces*, Topology Conference (Virginia Polytech. Inst. and State Univ., Blacksburg, VA, 1973), Lecture Notes in Math., vol. 375, Springer, Berlin, 1974, pp. 149–152.
- [LS] Ludwig L., Szeptycki P.J., *A consistent example of a β -normal not normal space*, Proceedings of the 2000 Topology and Dynamics Conference (San Antonio, TX), Topology Proc. **25** (2000), 1–4.

DEPARTMENT OF MATH. LOGIC, CHARLES UNIVERSITY, MALOSTRANSKÉ NÁM. 25,
118 00 PRAHA 1, CZECH REPUBLIC

E-mail: emur4296@menza.mff.cuni.cz

(Received May 28, 2001)