

Ryotaro Sato

On the range of a closed operator in an L_1 -space of vector-valued functions

Commentationes Mathematicae Universitatis Carolinae, Vol. 46 (2005), No. 2, 349--367

Persistent URL: <http://dml.cz/dmlcz/119529>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2005

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

On the range of a closed operator in an L_1 -space of vector-valued functions

RYOTARO SATO

Abstract. Let X be a reflexive Banach space and A be a closed operator in an L_1 -space of X -valued functions. Then we characterize the range $R(A)$ of A as follows. Let $0 \neq \lambda_n \in \rho(A)$ for all $1 \leq n < \infty$, where $\rho(A)$ denotes the resolvent set of A , and assume that $\lim_{n \rightarrow \infty} \lambda_n = 0$ and $\sup_{n \geq 1} \|\lambda_n(\lambda_n - A)^{-1}\| < \infty$. Furthermore, assume that there exists $\lambda_\infty \in \rho(A)$ such that $\|\lambda_\infty(\lambda_\infty - A)^{-1}\| \leq 1$. Then $f \in R(A)$ is equivalent to $\sup_{n \geq 1} \|(\lambda_n - A)^{-1}f\|_1 < \infty$. This generalizes Shaw’s result for scalar-valued functions.

Keywords: reflexive Banach space, L_1 -space of vector-valued functions, closed operator, resolvent set, range and domain, linear contraction, C_0 -semigroup, strongly continuous cosine family of operators

Classification: Primary 47A35; Secondary 47A05, 47D06, 47D09

1. Introduction

Let A be a (bounded or unbounded) closed operator in a Banach space Y with range $R(A)$ and domain $D(A)$. By assuming that the resolvent set $\rho(A)$ of A includes a countable set $\{\lambda_n : n \geq 1\}$, with $\lambda_n \neq 0$ for all $n \geq 1$, such that $\lim_{n \rightarrow \infty} \lambda_n = 0$ and $\sup_{n \geq 1} \|\lambda_n(\lambda_n - A)^{-1}\| < \infty$, it was shown in [11] that the obviously necessary condition

$$\sup_{n \geq 1} \|(\lambda_n - A)^{-1}x\| < \infty$$

is sufficient for an element x of Y to be in the range $R(A)$ of A when Y is reflexive. This can be regarded as a generalization of a result of Browder [2]; motivated by a result of Gottschalk and Hedlund (cf. Theorem 14.11 in [5]), he studied the problem of finding a necessary and sufficient condition for an element x of a Banach space Y to be in the range of $T - I$ when T is power-bounded on Y , and proved that the obviously necessary condition

$$(*) \quad \sup_{n \geq 1} \left\| \sum_{k=0}^{n-1} T^k x \right\| < \infty$$

is sufficient when Y is reflexive.

It was shown in [8] that also for T a contraction of $L_1(\mu)$ condition $(*)$ implies $x \in R(T - I)$, and in [6] an analogue for semigroups of contractions in $L_1(\mu)$ was proved. A unified treatment of these two results was given in [11]. The problem whether in $L_1(\mu)$ the norm condition $\|T\| \leq 1$ can be replaced by power-boundedness is still unresolved; a partial answer was given in [1].

In this paper we treat the case of operators in the space $L_1((\Omega, \mathcal{B}, \mu); X)$ of vector-valued norm-integrable functions on a σ -finite measure space $(\Omega, \mathcal{B}, \mu)$, with values in a reflexive Banach space X . The main result (Theorem 1) is the vector-valued version of [11]. The applications extend accordingly the results of [8], [6] and [11].

Let $(X, \|\cdot\|_X)$ be a reflexive Banach space, and $(\Omega, \mathcal{B}, \mu)$ be a σ -finite measure space. For $1 \leq p \leq \infty$, let $L_p(\Omega; X) = L_p((\Omega, \mathcal{B}, \mu); X)$ denote the usual Banach space of all X -valued strongly measurable functions f on Ω with the norm

$$\|f\|_p := \left(\int \|f(\omega)\|_X^p d\mu(\omega) \right)^{1/p} < \infty \quad \text{if } 1 \leq p < \infty,$$

$$\|f\|_\infty := \text{ess sup}\{\|f(\omega)\|_X : \omega \in \Omega\} < \infty \quad \text{if } p = \infty.$$

We consider a closed operator A in $L_1(\Omega; X)$ with range $R(A)$ and domain $D(A)$. We assume that the resolvent set $\rho(A)$ of A includes a countable set $\{\lambda_n : n \geq 1\}$, with $\lambda_n \neq 0$ for all $n \geq 1$, such that $\lim_{n \rightarrow \infty} \lambda_n = 0$ and $\sup_{n \geq 1} \|\lambda_n(\lambda_n - A)^{-1}\| < \infty$. Then we prove that $\sup_{n \geq 1} \|(\lambda_n - A)^{-1}f\|_1 < \infty$ implies $f \in R(A)$, under the additional hypothesis that there exists $\lambda_\infty \in \rho(A)$ such that $\|\lambda_\infty(\lambda_\infty - A)^{-1}\| \leq 1$. It would be interesting to ask whether this implication holds without the additional hypothesis. Concerning the problem the author would like to note that Assani [1] considered a power-bounded linear operator T on L_1 of scalar-valued functions, and under the hypothesis that

$$(**) \quad \lim_{n \rightarrow \infty} h_n = 0 \quad \text{a.e.} \quad \text{implies} \quad \lim_{n \rightarrow \infty} Th_n = 0 \quad \text{a.e.},$$

he proved that $\sup_{n \geq 1} \|\sum_{k=1}^n T^k f\|_1 < \infty$ is equivalent to $f \in R(T - I)$. It seems to the author that it is an open problem to prove this equivalence relation without assuming condition $(**)$. (See also [10], where similar results are proved for vector-valued functions.)

As applications of the result we characterize the range $R(A)$ of A , where A is the generator of a discrete semigroup $\{T^n : n \geq 0\}$, or a C_0 -semigroup $\{T(t) : t \geq 0\}$, or a strongly continuous cosine family $\{C(t) : -\infty < t < \infty\}$ of linear contractions on $L_1(\Omega; X)$. The results obtained below generalize Shaw's results (see [11, Corollaries 4, 6, and 8]) for scalar-valued functions. See also Lin and Sine [8], Krengel and Lin [6] for related topics.

2. The range of a closed operator in $L_1(\Omega; X)$

The following theorem is our main result.

Theorem 1 (cf. Theorem 2 of Shaw [11]). *Let X be a reflexive Banach space, and A be a closed operator in $L_1(\Omega; X)$ with domain $D(A)$ and range $R(A)$. Let $\rho(A)$ denote the resolvent set of A , and assume that $0 \neq \lambda_n \in \rho(A)$ for all $n \geq 1$ and $\lim_{n \rightarrow \infty} \lambda_n = 0$. If $M := \sup_{n \geq 1} \|\lambda_n(\lambda_n - A)^{-1}\| < \infty$, and there exists $\lambda_\infty \in \rho(A)$ such that $\|\lambda_\infty(\lambda_\infty - A)^{-1}\| \leq 1$, then the following conditions are equivalent for $f \in L_1(\Omega; X)$.*

- (I) $\sup_{n \geq 1} \|(\lambda_n - A)^{-1}f\|_1 < \infty$.
- (II) $f \in \bar{R}(A)$.

To prove this theorem we need the following lemma, which may be regarded as a generalization of the Yosida-Hewitt theorem on vector-measures (see [3, p. 30, Theorem I.5.8]).

Lemma 1. *Let X be a reflexive Banach space, and let $\ell \in L_\infty(\Omega; X^*)^*$ ($= L_1(\Omega; X)^{**}$). Then there exist unique ℓ_c and ℓ_p in $L_\infty(\Omega; X^*)^*$ such that*

- (a) $\ell = \ell_c + \ell_p$, and $\|\ell\| = \|\ell_c\| + \|\ell_p\|$;
- (b) there exists $g \in L_1(\Omega; X)$ with

$$(1) \quad \ell_c(f) = \int_{\Omega} \langle g(\omega), f^*(\omega) \rangle d\mu(\omega) \quad \text{for all } f^* \in L_\infty(\Omega; X^*);$$

- (c) if we define a scalar-valued function G_{x^*} on \mathcal{B} for each $x^* \in X^*$ by

$$(2) \quad G_{x^*}(B) := \ell_p(\chi_B(\cdot)x^*) \quad (B \in \mathcal{B}),$$

then G_{x^*} is a purely finitely additive measure on \mathcal{B} , i.e., there is no (countably additive) measure λ on \mathcal{B} satisfying $0 \leq \lambda(B) \leq |G_{x^*}|(B)$ for all $B \in \mathcal{B}$, where $|G_{x^*}|$ denotes the variation of G_{x^*} (cf. [3, p. 2]).

PROOF: For $B \in \mathcal{B}$, define a linear functional $F(B)$ on X^* by

$$(3) \quad F(B)(x^*) := \ell(\chi_B(\cdot)x^*) \quad (x^* \in X^*).$$

Since $|F(B)(x^*)| \leq \|\ell\| \|x^*\|$, it follows that $\|F(B)\| \leq \|\ell\|$. Thus we may regard $F(B)$ as an element of $X^{**} = X$, and hence we can write

$$(4) \quad \langle F(B), x^* \rangle = \ell(\chi_B(\cdot)x^*) \quad (B \in \mathcal{B}, x^* \in X^*).$$

Clearly, $F : \mathcal{B} \rightarrow X$ is finitely additive. To see that F is a finitely additive vector-measure of bounded variation, let $\{B_1, \dots, B_n\}$ be a finite measurable decomposition of Ω , and $x_j^* \in X^* (j = 1, \dots, n)$ be such that $\|x_j^*\| \leq 1$. Then

$$\left| \sum_{j=1}^n \langle F(B_j), x_j^* \rangle \right| = \left| \ell \left(\sum_{j=1}^n \chi_{B_j}(\cdot)x_j^* \right) \right| \leq \|\ell\| \cdot \left\| \sum_{j=1}^n \chi_{B_j}(\cdot)x_j^* \right\|_{L_\infty(\Omega; X^*)} \leq \|\ell\|,$$

so that $\sum_{j=1}^n \|F(B_j)\| \leq \|\ell\|$. Therefore, F is of bounded variation. Let $|F|$ denote the variation of F . Then, since $|F|(\Omega) \leq \|\ell\|$, it follows from Corollary I.5.3 of [3] that F is strongly additive; and hence by the Yosida-Hewitt theorem (cf. [3, p. 30, Theorem I.5.8]), there exist unique strongly additive X -valued measures F_c and F_p on \mathcal{B} (which are of bounded variation) such that

- (i) F_c is countably additive;
- (ii) for each $x^* \in X^*$, x^*F_p is purely finitely additive on \mathcal{B} ;
- (iii) $F = F_c + F_p$;
- (iv) F_c and F_p are mutually singular, i.e., for each $\epsilon > 0$ there exists $E \in \mathcal{B}$ such that $|F_c|(\Omega \setminus E) + |F_p|(E) < \epsilon$;
- (v) $|F| = |F_c| + |F_p|$.

Since X has the Radon-Nikodym property (cf. [3, p. 82, Corollary III.3.4]), there exists $g \in L_1(\Omega; X)$ such that $F_c(B) = \int_B g \, d\mu$ for all $B \in \mathcal{B}$. Using this g , we define a linear functional ℓ_c on $L_\infty(\Omega; X^*)$ by

$$\ell_c(f) := \int_\Omega \langle g(\omega), f^*(\omega) \rangle \, d\mu(\omega) \quad (f^* \in L_\infty(\Omega; X^*)).$$

It is clear that $\ell_c \in L_\infty(\Omega; X^*)^*$ and $\|\ell_c\| = \|g\|_1$. We then put

$$\ell_p := \ell - \ell_c,$$

so that $\ell_p \in L_\infty(\Omega; X^*)^*$ and $\ell = \ell_c + \ell_p$. Let $x^* \in X^*$ and $B \in \mathcal{B}$. Then, by (2) and (3),

$$\begin{aligned} G_{x^*}(B) &= \ell_p(\chi_B(\cdot)x^*) = (\ell - \ell_c)(\chi_B(\cdot)x^*) = \ell(\chi_B(\cdot)x^*) - \ell_c(\chi_B(\cdot)x^*) \\ &= \langle F(B), x^* \rangle - \langle F_c(B), x^* \rangle = \langle F_p(B), x^* \rangle, \end{aligned}$$

which implies that $x^*F_p = G_{x^*}$ on \mathcal{B} for each $x^* \in X^*$. Thus, G_{x^*} is purely finitely additive on \mathcal{B} by (ii).

Next, we prove that $\|\ell\| = \|\ell_c\| + \|\ell_p\|$. To do this, let $\epsilon > 0$. Then, by (iv) there exists $E \in \mathcal{B}$ such that

$$(5) \quad |F_c|(\Omega \setminus E) + |F_p|(E) < \epsilon.$$

Since the set of all countably X^* -valued functions in $L_\infty(\Omega; X^*)$ is a dense subset of $L_\infty(\Omega; X^*)$, there exists $f_1^* \in L_\infty(\Omega; X^*)$ of the form

$$f_1^* = \sum_{n=1}^\infty \chi_{B_n}(\cdot)x_n^*,$$

where $x_n^* \in X^*$, $\|x_n^*\| \leq 1$, and $\{B_n : n \geq 1\}$ is a countable measurable decomposition of Ω , such that

$$(6) \quad |\ell_p(f_1^*)| > \|\ell_p\| - \epsilon.$$

Write $E_n = E \cap (\bigcup_{j=1}^n B_j)$. It follows that $E_n \uparrow E$ as $n \rightarrow \infty$, and

$$\begin{aligned} |F_c|(\Omega \setminus E) &= \int_{\Omega \setminus E} \|g(\omega)\|_X d\mu(\omega) \\ &= \lim_{n \rightarrow \infty} \int_{\Omega \setminus E_n} \|g(\omega)\|_X d\mu(\omega) = \lim_{n \rightarrow \infty} |F_c|(\Omega \setminus E_n). \end{aligned}$$

Hence, we can choose E_N satisfying (5) with E_N in place of E . Then we can choose $f_2^* \in L_\infty(\Omega; X^*)$ of the form

$$f_2^* = \sum_{j=1}^{\infty} \chi_{D_j}(\cdot) y_j^*,$$

where $y_j^* \in X^*$, $\|y_j^*\| \leq 1$, and $\{D_j : j \geq 1\}$ is a countable measurable decomposition of the set E_N , such that

$$(7) \quad |\ell_c(f_2^*)| = \left| \int_{E_N} \langle g(\omega), f_2^*(\omega) \rangle d\mu(\omega) \right| > \int_{E_N} \|g(\omega)\|_X d\mu(\omega) - \epsilon.$$

Lastly, define an X^* -valued function f^* on Ω by

$$f^*(\omega) = \begin{cases} f_1^*(\omega) & \text{if } \omega \in \Omega \setminus E_N, \\ f_2^*(\omega) & \text{if } \omega \in E_N. \end{cases}$$

It is clear that $f^* \in L_\infty(\Omega; X^*)$ and $\|f^*\|_\infty \leq 1$. Furthermore, by (5) with E_N in place of E , (6) and (7),

$$\begin{aligned} |\ell(f^*)| &= |\ell_c(f_2^*) + \ell_p(f_1^*) + \ell_c(\chi_{\Omega \setminus E_N} f_1^*) + \ell_p(f_2^*) - \ell_p(\chi_{E_N} f_1^*)| \\ &> \left(\int_{E_N} \|g(\omega)\|_X d\mu(\omega) - \epsilon \right) + (\|\ell_p\| - \epsilon) - |\ell_c(\chi_{\Omega \setminus E_N} f_1^*)| \\ &\quad - |\ell_p(f_2^*)| - |\ell_p(\chi_{E_N} f_1^*)| \\ &> (\|\ell_c\| - 2\epsilon) + (\|\ell_p\| - \epsilon) - \epsilon - \epsilon - \epsilon = \|\ell_c\| + \|\ell_p\| - 6\epsilon. \end{aligned}$$

Since ϵ was arbitrary, this proves $\|\ell\| \geq \|\ell_c\| + \|\ell_p\|$. Consequently, $\|\ell\| = \|\ell_c\| + \|\ell_p\|$.

The uniqueness of the decomposition $\ell = \ell_c + \ell_p$ follows from the uniqueness of the decomposition $F = F_c + F_p$, and this completes the proof. \square

PROOF OF THEOREM 1: (I) \Rightarrow (II). We may assume here that $\lambda_\infty \neq 0$, because $\lambda_\infty = 0$ implies $R(A) = L_1(\Omega; X)$. Since $\{(\lambda_n - A)^{-1} f : n \geq 1\}$ is a bounded subset of the dual space of $L_\infty(\Omega; X^*)$, it is relatively compact with respect to

the weak*-topology. It follows that there exists $\eta \in L_1(\Omega; X)^{**}$ which is a weak*-cluster point of the sequence $\{(\lambda_n - A)^{-1}f\}_{n=1}^\infty$.

Let $u \in L_\infty(\Omega; X^*)$ and $0 \neq \lambda \in \rho(A)$. Then there exists a subsequence $\{n_j\}_{j=1}^\infty$ of the sequence $\{n\}_{n=1}^\infty$ such that

$$\begin{aligned} \langle (\lambda(\lambda - A)^{-1})^{**}\eta, u \rangle &= \langle \eta, (\lambda(\lambda - A)^{-1})^*u \rangle \\ &= \lim_{j \rightarrow \infty} \langle (\lambda_{n_j} - A)^{-1}f, (\lambda(\lambda - A)^{-1})^*u \rangle \\ &= \lim_{j \rightarrow \infty} \langle \lambda(\lambda - A)^{-1}(\lambda_{n_j} - A)^{-1}f, u \rangle \\ &= \lim_{j \rightarrow \infty} \frac{\lambda}{\lambda - \lambda_{n_j}} \cdot \langle (\lambda_{n_j} - A)^{-1}f - (\lambda - A)^{-1}f, u \rangle \\ &= \langle \eta, u \rangle - \langle (\lambda - A)^{-1}f, u \rangle, \end{aligned}$$

where the last but one equality is due to the resolvent equation. Consequently, we obtain that

$$(8) \quad (\lambda(\lambda - A)^{-1})^{**}\eta = \eta - (\lambda - A)^{-1}f \quad (\lambda \in \rho(A), \lambda \neq 0).$$

Here, we apply Lemma 1 for η as follows. By Lemma 1, there exist unique η_c and η_p in $L_1(\Omega; X)^{**}$ such that

(i) there exists $g \in L_1(\Omega; X)$ with

$$\eta_c(u) = \int_{\Omega} \langle g(\omega), u^*(\omega) \rangle d\mu(\omega) \quad (u \in L_\infty(\Omega; X^*));$$

(ii) if we define a scalar-valued function G_{x^*} on \mathcal{B} for each $x^* \in X^*$ by

$$G_{x^*}(B) := \eta_p(\chi_B(\cdot)x^*) \quad (B \in \mathcal{B}),$$

then G_{x^*} is a purely finitely additive measure on \mathcal{B} ;

(iii) $\eta = \eta_c + \eta_p$, and $\|\eta\| = \|\eta_c\| + \|\eta_p\|$.

By (i) we may identify η_c with g . Then, putting $\lambda = \lambda_\infty$, we have by (8)

$$(\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\eta = g + \eta_p - (\lambda_\infty - A)^{-1}f.$$

On the other hand, we also have

$$\begin{aligned} (\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\eta &= (\lambda_\infty(\lambda_\infty - A)^{-1})^{**}(g + \eta_p) \\ &= \lambda_\infty(\lambda_\infty - A)^{-1}g + (\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\eta_p, \end{aligned}$$

whence

$$(9) \quad (\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\eta_p = \eta_p + g - (\lambda_\infty - A)^{-1}f - \lambda_\infty(\lambda_\infty - A)^{-1}g.$$

Since $\|(\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\| = \|\lambda_\infty(\lambda_\infty - A)^{-1}\| \leq 1$ by hypothesis, it follows that

$$\|\eta_p\| \geq \|(\lambda_\infty(\lambda_\infty - A)^{-1})^{**}\eta_p\| = \|\eta_p + g - (\lambda_\infty - A)^{-1}f - \lambda_\infty(\lambda_\infty - A)^{-1}g\|.$$

Here we notice that $g - (\lambda_1 - A)^{-1}f - \lambda_1(\lambda_1 - A)^{-1}g$ is a function in $L_1(\Omega; X)$ and η_p is an element of $L_1(\Omega; X)^{**}$ satisfying condition (ii). Thus by Lemma 1 we have

$$\|\eta_p\| = \|\eta_p\| + \|g - (\lambda_1 - A)^{-1}f - \lambda_1(\lambda_1 - A)^{-1}g\|_1,$$

which implies

$$g = (\lambda_1 - A)^{-1}f + \lambda_1(\lambda_1 - A)^{-1}g.$$

Consequently, $g \in D(A)$ and $(\lambda_1 - A)g = f + \lambda_1g$, so that $f = A(-g) \in R(A)$.

(II) \Rightarrow (I). If $f = Ag$ for some $g \in L_1(\Omega; X)$, then

$$(\lambda_n - A)^{-1}f = (\lambda_n - A)^{-1}Ag = A(\lambda_n - A)^{-1}g = \lambda_n(\lambda_n - A)^{-1}g - g,$$

and thus $\|(\lambda_n - A)^{-1}f\|_1 \leq \|\lambda_n(\lambda_n - A)^{-1}\| \|g\|_1 + \|g\|_1 \leq (M + 1)\|g\|_1$ for all $n \geq 1$.

This completes the proof of Theorem 1. □

Using the argument of the above proof we can prove the following proposition, which is of independent interest in view of the results of [4] and [12].

Proposition 1. *Let X be a reflexive Banach space, and A be a closed operator in $L_1(\Omega; X)$ with domain $D(A)$ and range $R(A)$. Suppose there exists $\lambda \in \rho(A)$ such that $\|\lambda(\lambda - A)^{-1}\| \leq 1$. Then $A(U \cap D(A))$ is a closed subset of $L_1(\Omega; X)$, where U is the closed unit ball of $L_1(\Omega; X)$, i.e., $U = \{f \in L_1(\Omega; X) : \|f\|_1 \leq 1\}$.*

PROOF: Let $f_n \in U \cap D(A)$, $n = 1, 2, \dots$, and $f \in L_1(\Omega; X)$ be such that $\lim_{n \rightarrow \infty} \|Af_n - f\|_1 = 0$. We must prove that $f \in A(U \cap D(A))$. To do this, let $\eta \in L_1(\Omega; X)^{**}$ be a weak*-cluster point of the sequence $\{f_n\}_{n=1}^\infty (\subset L_1(\Omega; X)^{**})$. Then, for $u \in L_\infty(\Omega; X^*)$ there exists a subsequence $\{n_j\}_{j=1}^\infty$ of the sequence $\{n\}_{n=1}^\infty$ such that

$$\begin{aligned} \langle (\lambda(\lambda - A)^{-1})^{**}\eta, u \rangle &= \langle \eta, (\lambda(\lambda - A)^{-1})^*u \rangle = \lim_{j \rightarrow \infty} \langle f_{n_j}, (\lambda(\lambda - A)^{-1})^*u \rangle \\ &= \lim_{j \rightarrow \infty} \langle \lambda(\lambda - A)^{-1}f_{n_j}, u \rangle = \lim_{j \rightarrow \infty} \langle (I + A(\lambda - A)^{-1})f_{n_j}, u \rangle \\ &= \lim_{j \rightarrow \infty} \langle f_{n_j} + (\lambda - A)^{-1}Af_{n_j}, u \rangle = \langle \eta + (\lambda - A)^{-1}f, u \rangle. \end{aligned}$$

It follows that $(\lambda(\lambda - A)^{-1})^{**}\eta = \eta + (\lambda - A)^{-1}f$. Thus, as in the proof of (I) \Rightarrow (II) of Theorem 1, letting $\eta = \eta_c + \eta_p$ and identifying η_c with a function g in $L_1(\Omega; X)$, we see that

$$\lambda(\lambda - A)^{-1}g + (\lambda(\lambda - A)^{-1})^{**}\eta_p = g + \eta_p + (\lambda - A)^{-1}f,$$

so that

$$(\lambda(\lambda - A)^{-1})^{**}\eta_p = \eta_p + g + (\lambda - A)^{-1}f - \lambda(\lambda - A)^{-1}g.$$

By this and the fact $\|(\lambda(\lambda - A)^{-1})^{**}\| \leq 1$, it follows from Lemma 1 that

$$g + (\lambda - A)^{-1}f - \lambda(\lambda - A)^{-1}g = 0.$$

Hence $(\lambda - A)g + f - \lambda g = 0$, and we see that $f = Ag$ with $g \in D(A)$ and $\|g\|_1 \leq \|\eta\| \leq 1$. This completes the proof. \square

3. Applications

Let T be a bounded linear operator on $L_1(\Omega; X)$. For $\gamma \neq -1, -2, \dots$ we define the Cesàro means of order γ (or γ -Cesàro means) $C_n^\gamma(T)$ of the discrete semigroup $\{T^n : n \geq 0\}$ by

$$(10) \quad C_n^\gamma(T) := \frac{1}{\sigma_n^\gamma} \sum_{k=0}^n \sigma_{n-k}^{\gamma-1} T^k \quad (n \geq 0),$$

where $\sigma_n^\beta := (\beta + 1)(\beta + 2) \dots (\beta + n)/n!$ for $n \geq 1$, and $\sigma_0^\beta := 1$ (cf. [15, Chapter 3]). Among them are the following two particular means: $C_n^0(T) = T^n$ and $C_n^1(T) = (n + 1)^{-1} \sum_{k=0}^n T^k$. As is well-known, only the case $\gamma > -1$ is of interest. The Abel means of $\{T^n : n \geq 0\}$ are the operators

$$(11) \quad A_r(T) := (1 - r) \sum_{n=0}^\infty r^n T^n$$

defined for $0 < r < 1/r(T)$, where $r(T) := \lim_{n \rightarrow \infty} \|T^n\|^{1/n}$ denotes the spectral radius of T . It is known (cf. [15, Chapter 3]) that if $r(T) \leq 1$ and $0 < \gamma < \beta < \infty$, then

$$(12) \quad \sup_{n \geq 0} \|T^n f\|_1 \geq \sup_{n \geq 0} \|C_n^\gamma(T)f\|_1 \geq \sup_{n \geq 0} \|C_n^\beta(T)f\|_1 \geq \sup_{0 < r < 1} \|A_r(T)f\|_1$$

for every $f \in L_1(\Omega; X)$.

The first application of Theorem 1 is the following

Theorem 2 (cf. Theorem 7 of [8]). *Let X be a reflexive Banach space, and T be a linear contraction on $L_1(\Omega; X)$. Assume that $\alpha \geq 1$. Then the following conditions are equivalent for $f \in L_1(\Omega; X)$.*

- (I) $\sup_{n \geq 0} n \|C_n^\alpha(T)f\|_1 < \infty$.
- (II) $\sup_{0 < r < 1} \|\sum_{n=0}^\infty r^n T^n f\|_1 < \infty$.
- (III) $f \in R(T - I)$.

PROOF: (I) \Rightarrow (II). Since

$$nC_n^\alpha(T)f = \frac{n}{\sigma_n^\alpha} \sum_{k=0}^n \sigma_{n-k}^{\alpha-1} T^k f \quad \text{and} \quad \frac{\sigma_n^\alpha}{\sigma_{n-1}^{\alpha-1}} = \frac{\alpha + n}{\alpha} \sim \frac{n}{\alpha} \quad (n \rightarrow \infty),$$

$n \|C_n^\alpha(T)f\|_1 = O(1)$ ($n \rightarrow \infty$) is equivalent to

$$C := \sup_{n \geq 0} \left\| \frac{1}{\sigma_n^{\alpha-1}} \sum_{k=0}^n \sigma_{n-k}^{\alpha-1} T^k f \right\|_1 < \infty.$$

Then, for $0 < r < 1$ we have

$$\begin{aligned} \sum_{n=0}^\infty r^n T^n f &= (1-r)^\alpha (1-r)^{-\alpha} \sum_{n=0}^\infty r^n T^n f \\ &= (1-r)^\alpha \left(\sum_{n=0}^\infty \sigma_n^{\alpha-1} r^n \right) \left(\sum_{n=0}^\infty r^n T^n f \right) \\ &= (1-r)^\alpha \sum_{n=0}^\infty \sigma_n^{\alpha-1} r^n \left(\frac{1}{\sigma_n^{\alpha-1}} \sum_{k=0}^n \sigma_{n-k}^{\alpha-1} T^k f \right), \end{aligned}$$

so that

$$\left\| \sum_{n=0}^\infty r^n T^n f \right\|_1 \leq (1-r)^\alpha \sum_{n=0}^\infty \sigma_n^{\alpha-1} r^n \cdot C = C.$$

(II) \Rightarrow (III). Putting $A = T - I$, we have for $\lambda > 0$

$$(\lambda - A)^{-1} = (\lambda + 1 - T)^{-1} = \frac{1}{\lambda + 1} \sum_{n=0}^\infty \left(\frac{1}{\lambda + 1} \right)^n T^n,$$

whence $\|T\| \leq 1$ implies

$$\|\lambda(\lambda - A)^{-1}\| \leq \frac{\lambda}{\lambda + 1} \sum_{n=0}^\infty \left(\frac{1}{\lambda + 1} \right)^n = 1.$$

Furthermore, we get from (II) that

$$\begin{aligned} \sup_{\lambda > 0} \|(\lambda - A)^{-1}f\|_1 &= \sup_{\lambda > 0} \left\| \frac{1}{\lambda + 1} \sum_{n=0}^{\infty} \left(\frac{1}{\lambda + 1}\right)^n T^n f \right\|_1 \\ &\leq \sup_{\lambda > 0} \left\| \sum_{n=0}^{\infty} \left(\frac{1}{\lambda + 1}\right)^n T^n f \right\|_1 < \infty. \end{aligned}$$

Hence, $f \in R(A) = R(T - I)$ by Theorem 1.

(III) \Rightarrow (I). Suppose $f = (T - I)g$ for some $g \in L_1(\Omega; X)$. Using the fundamental relation $C_n^\beta(T)(T - I) = \frac{\beta}{n+1}(C_{n+1}^{\beta-1}(T) - I)$ for $\beta > 0$ and $n \geq 0$, which can be proved by an elementary calculation (cf. [15, Chapter 3]), we see that

$$nC_n^\alpha(T)f = nC_n^\alpha(T)(T - I)g = \frac{n\alpha}{n + 1}(C_{n+1}^{\alpha-1}(T) - I)g.$$

Then, since $\|C_{n+1}^{\alpha-1}(T)\| \leq 1$ (which comes from the hypotheses that $\|T\| \leq 1$ and that $\alpha \geq 1$), it follows that

$$n\|C_n^\alpha(T)f\|_1 \leq \alpha(\|C_{n+1}^{\alpha-1}(T)\| + 1)\|g\|_1 \leq 2\alpha\|g\|_1 \quad (n \geq 0).$$

This completes the proof of Theorem 2. □

Remarks (on Theorem 2). (a) If $-1 < \alpha < 1$, then (III) \Rightarrow (I) does not hold in general. To see this, first suppose that $\alpha \neq 0$ and $-1 < \alpha < 1$. Then we can use the equation $C_n^\alpha(T)(T - I) = \frac{\alpha}{n+1}(C_{n+1}^{\alpha-1}(T) - I)$. If $f = (T - I)g$, then

$$(13) \quad nC_n^\alpha(T)f = nC_n^\alpha(T)(T - I)g = \frac{n\alpha}{n + 1}(C_{n+1}^{\alpha-1}(T)g - g),$$

so that $\lim_{n \rightarrow \infty} \|C_n^{\alpha-1}(T)g\|_1 = \infty$ implies

$$(14) \quad \|nC_n^\alpha(T)f\|_1 \geq \frac{n|\alpha|}{n + 1}(\|C_{n+1}^{\alpha-1}(T)g\|_1 - \|g\|_1) \longrightarrow \infty \quad \text{as } n \rightarrow \infty.$$

To see the possibility of the case that $\lim_{n \rightarrow \infty} \|C_n^{\alpha-1}(T)g\|_1 = \infty$, let m be the counting measure on the set \mathbb{Z} of all integers, and $L_1(\mathbb{Z}, m)$ be the L_1 -space of real-valued functions on \mathbb{Z} with respect to the measure m . Define a positive linear isometry T on $L_1(\mathbb{Z}, m)$ by $Tf(k) = f(k - 1)$ for $k \in \mathbb{Z}$. Then, the function $g = \chi_{\{0\}}$ satisfies $T^n g = \chi_{\{n\}}$ for $n \geq 0$, and hence

$$\|C_n^{\alpha-1}(T)g\|_1 = \left\| \frac{1}{\sigma_n^{\alpha-1}} \sum_{k=0}^n \sigma_{n-k}^{\alpha-2} T^k g \right\|_1 \geq \frac{1}{|\sigma_n^{\alpha-1}|} \|\sigma_0^{\alpha-2} \chi_{\{n\}\}\|_1 = \frac{1}{|\sigma_n^{\alpha-1}|} \longrightarrow \infty$$

as $n \rightarrow \infty$, since $\sigma_n^{\alpha-1} \sim n^{\alpha-1}/\Gamma(\alpha)$ ($n \rightarrow \infty$) (cf. [15, p.77]). (For related topics we refer the reader to [7].)

Next, suppose that $\alpha = 0$. In this case, for any isometry $T \neq I$ and any $f \neq 0$, with $f \in R(T - I)$, we have $n\|C_n^0(T)f\| = n\|T^n f\| = n\|f\| \rightarrow \infty$ as $n \rightarrow \infty$. This completes the proof.

(b) The implication (I) \Rightarrow (II) holds for every $\alpha > -1$, with $\alpha \neq 0$. To see this, it suffices to consider only the case where $-1 < \alpha < 1$ and $\alpha \neq 0$, by Theorem 2. Now, choose $\beta > 0$ satisfying $\beta + \alpha \geq 1$. Then, since

$$C := \sup_{n \geq 0} \left\| \frac{1}{\sigma_n^{\alpha-1}} \sum_{k=0}^n \sigma_{n-k}^{\alpha-1} T^k f \right\|_1 < \infty,$$

$$\sum_{k=0}^n \sigma_{n-k}^{\beta+\alpha-1} T^k f = \sum_{k=0}^n \sigma_{n-k}^{\beta-1} \sum_{l=0}^k \sigma_{k-l}^{\alpha-1} T^l f, \quad \text{and} \quad \sigma_n^{\beta+\alpha-1} = \sum_{k=0}^n \sigma_{n-k}^{\beta-1} \sigma_k^{\alpha-1},$$

it follows that

$$\left\| \frac{1}{\sigma_n^{\beta+\alpha-1}} \sum_{k=0}^n \sigma_{n-k}^{\beta+\alpha-1} T^k f \right\|_1 \leq \frac{\sum_{k=0}^n \sigma_{n-k}^{\beta-1} \sigma_k^{\alpha-1} C}{\sum_{k=0}^n \sigma_{n-k}^{\beta-1} \sigma_k^{\alpha-1}} = C,$$

whence $\sup_{n \geq 0} n\|C_n^{\beta+\alpha}(T)f\|_1 < \infty$, and thus $f \in R(T - I)$ by Theorem 2.

On the other hand, if $\alpha = 0$, then the implication (I) \Rightarrow (II) fails to hold in general. To see this, let μ be the measure on \mathbb{Z} defined by

$$\mu(\{k\}) = \begin{cases} 1 & \text{if } k \leq 0, \\ (k+1)^{-1} & \text{if } k \geq 1. \end{cases}$$

Let T be the positive linear contraction on $L_1(\mathbb{Z}, \mu)$ defined by $Tf(k) = f(k-1)$ for $k \in \mathbb{Z}$. Then the function $g = \chi_{\{0\}}$ satisfies

$$(15) \quad n\|C_n^0(T)g\|_1 = n\|T^n g\|_1 = n\|\chi_{\{n\}}\|_1 = \frac{n}{n+1} < 1 \quad (n \geq 0).$$

By the definitions,

$$\left\| \sum_{j=0}^n T^j g \right\|_1 = \|\chi_{[0,n]}\|_1 = \sum_{j=0}^n \frac{1}{j+1} \rightarrow \infty$$

as $n \rightarrow \infty$, so $g \notin L_1(T - I)$. Thus (II) with g in place of f does not hold, by Theorem 2.

(c) In Theorem 2 the condition $\sup_{0 < r < 1} \|\sum_{n=0}^{\infty} r^n T^n f\|_1 < \infty$ can be replaced with the weaker condition $\liminf_{r \uparrow 1} \|\sum_{n=0}^{\infty} r^n T^n f\|_1 < \infty$, which follows from Theorem 1.

Next, we consider a C_0 -semigroup $T(\cdot) \equiv \{T(t) : t \geq 0\}$ of linear contractions on $L_1(\Omega; X)$. Thus, $T(s+t) = T(s)T(t)$ for all $s, t \geq 0$, and $\lim_{t \downarrow 0} \|T(t)f - f\|_1 = 0$ for each $f \in L_1(\Omega; X)$. The infinitesimal generator A of $T(\cdot)$ is defined by $Af := \lim_{t \downarrow 0} t^{-1}(T(t)f - f)$, with domain $D(A)$ the set of all those $f \in L_1(\Omega; X)$ for which this limit exists. It is known (cf. e.g. [9]) that A is a densely defined closed operator; and since $\|T(t)\| \leq 1$ for all $t \geq 0$, if $\lambda > 0$, then $\lambda \in \rho(A)$ and $(\lambda - A)^{-1}f = \int_0^{\infty} e^{-\lambda s} T(s)f ds$ for all $f \in L_1(\Omega; X)$. Therefore we have $\sup_{\lambda > 0} \|\lambda(\lambda - A)^{-1}\| \leq 1$. The Cesàro means of order γ (or γ -Cesàro means) $C_t^\gamma(T(\cdot))$ of the semigroup $T(\cdot)$, where $\gamma \geq 0$ and $t > 0$, are the operators defined by $C_t^0(T(\cdot)) := T(t)$ for $\gamma = 0$, and

$$(16) \quad C_t^\gamma(T(\cdot))f := \gamma t^{-\gamma} \int_0^t (t-s)^{\gamma-1} T(s)f ds \quad (\gamma > 0, f \in L_1(\Omega; X)).$$

In particular, if $\gamma = 1$, then we have $C_t^1(T(\cdot))f = t^{-1} \int_0^t T(s)f ds$. The Abel means $A_\lambda(T(\cdot))$ of $T(\cdot)$ are the operators

$$(17) \quad A_\lambda(T(\cdot))f := \lambda \int_0^{\infty} e^{-\lambda s} T(s)f ds \quad (\lambda > 0, f \in L_1(\Omega; X)).$$

Fubini's theorem and an induction argument on n imply easily the following facts:

(i) If $0 < \gamma, \beta < \infty$ then for every $f \in L_1(\Omega; X)$ and $t > 0$,

$$(18) \quad C_t^{\gamma+\beta}(T(\cdot))f = \frac{\int_0^t (t-s)^{\beta-1} [\int_0^s (s-r)^{\gamma-1} T(r)f dr] ds}{\int_0^t (t-s)^{\beta-1} [\int_0^s (s-r)^{\gamma-1} dr] ds}.$$

(ii) If $n \geq 1$ is an integer, then for every $f \in L_1(\Omega; X)$ and $t > 0$,

$$(19) \quad C_t^n(T(\cdot))f = n! t^{-n} \int_0^t \left[\int_0^{s_1} \left(\int_0^{s_2} \left(\dots \left(\int_0^{s_{n-1}} T(s_n)f ds_n \right) \dots \right) ds_3 \right) ds_2 \right] ds_1.$$

Furthermore, as in the discrete case (cf. (12)), we obtain that if $0 < \gamma < \beta < \infty$, then for every $f \in L_1(\Omega; X)$,

$$(20) \quad \begin{aligned} \sup_{t \geq 0} \|T(t)f\|_1 &\geq \sup_{t > 0} \|C_t^\gamma(T(\cdot))f\|_1 \\ &\geq \sup_{t > 0} \|C_t^\beta(T(\cdot))f\|_1 \geq \sup_{\lambda > 0} \|A_\lambda(T(\cdot))f\|_1. \end{aligned}$$

Theorem 3 (cf. Corollary 8 of [11]). *Let X be a reflexive Banach space, and A be the infinitesimal generator of a C_0 -semigroup $T(\cdot)$ of linear contractions on $L_1(\Omega; X)$. Assume that $\alpha \geq 1$. Then the following conditions are equivalent for $f \in L_1(\Omega; X)$.*

- (I) $\sup_{t>0} t \|C_t^\alpha(T(\cdot))f\|_1 < \infty$.
- (II) $\sup_{\lambda>0} \|\int_0^\infty e^{-\lambda t} T(t)f dt\|_1 < \infty$.
- (III) $f \in R(A)$.

PROOF: (I) \Rightarrow (II). We first show that there exists an integer $n \geq \alpha$ such that

$$(21) \quad M_n := \sup_{t>0} t \|C_t^n(T(\cdot))f\|_1 < \infty.$$

To prove this, we may assume that $\alpha > 1$. We then notice by (16) that the condition $\sup_{t>0} t \|C_t^\alpha(T(\cdot))f\|_1 < \infty$ is equivalent to

$$(22) \quad M(\alpha) := \sup_{t>0} \frac{\|\int_0^t (t-s)^{\alpha-1} T(s)f ds\|_1}{\int_0^t (t-s)^{\alpha-2} ds} < \infty.$$

Let $\beta > 0$. By Fubini's theorem

$$\begin{aligned} & \int_0^t (t-s)^{\beta-1} \left(\int_0^s (s-r)^{\alpha-1} T(r)f dr \right) ds \\ &= \int_0^t \left(\int_r^t (t-s)^{\beta-1} (s-r)^{\alpha-1} ds \right) T(r)f dr \\ &= \int_0^t (t-r)^{\beta+\alpha-1} \left(\int_0^1 (1-s)^{\beta-1} s^{\alpha-1} ds \right) T(r)f dr \\ &= B(\beta, \alpha) \int_0^t (t-r)^{\beta+\alpha-1} T(r)f dr \end{aligned}$$

and

$$\int_0^t (t-s)^{\beta-1} \left(\int_0^s (s-r)^{\alpha-2} dr \right) ds = B(\beta, \alpha - 1) \int_0^t (t-r)^{\beta+\alpha-2} dr,$$

where $B(p, q) := \int_0^1 (1-x)^{p-1} x^{q-1} dx$ ($p, q > 0$) is the Beta function. It follows that

$$\begin{aligned} M(\beta + \alpha) &= \sup_{t>0} \frac{\|\int_0^t (t-s)^{\beta+\alpha-1} T(s)f ds\|_1}{\int_0^t (t-s)^{\beta+\alpha-2} ds} \\ &= \frac{B(\beta, \alpha - 1)}{B(\beta, \alpha)} \cdot \sup_{t>0} \frac{\left\| \int_0^t (t-s)^{\beta-1} \left(\int_0^s (s-r)^{\alpha-1} T(r)f dr \right) ds \right\|_1}{\int_0^t (t-s)^{\beta-1} \left(\int_0^s (s-r)^{\alpha-2} dr \right) ds} \\ &\leq \frac{B(\beta, \alpha - 1)}{B(\beta, \alpha)} \cdot M(\alpha) < \infty \quad (\text{by (22)}). \end{aligned}$$

Therefore, (21) holds for any integer n , with $n > \alpha$.

Next, by Fubini's theorem

$$\begin{aligned} \int_0^\infty e^{-\lambda t} T(t) f dt &= \lambda \int_0^\infty e^{-\lambda t} \left[\int_0^t T(s_1) f ds_1 \right] dt = \dots \\ &= \lambda^n \int_0^\infty e^{-\lambda t} \left[\int_0^t \left\{ \int_0^{s_1} \left(\int_0^{s_2} \left(\dots \left(\int_0^{s_{n-1}} T(s_n) f ds_n \right) \dots \right) ds_3 \right) ds_2 \right\} ds_1 \right] dt \\ &= \lambda^n \int_0^\infty e^{-\lambda t} \frac{t^{n-1}}{n!} \cdot [tC_t^n(T(\cdot))f] dt \quad (\text{by (19)}). \end{aligned}$$

Thus we apply (21) to get that for $\lambda > 0$,

$$\begin{aligned} \left\| \int_0^\infty e^{-\lambda t} T(t) f dt \right\|_1 &\leq \lambda^n \int_0^\infty e^{-\lambda t} \frac{t^{n-1}}{n!} \|tC_t^n(T(\cdot))f\|_1 dt \\ &\leq \frac{M}{n!} \lambda^n \int_0^\infty e^{-\lambda t} t^{n-1} dt = \frac{M}{n}. \end{aligned}$$

(II) \Rightarrow (III). Since (II) implies

$$\sup_{\lambda > 0} \|(\lambda - A)^{-1} f\|_1 = \sup_{\lambda > 0} \left\| \int_0^\infty e^{-\lambda t} T(t) f dt \right\|_1 < \infty,$$

(III) follows from Theorem 1.

(III) \Rightarrow (I). Suppose $f = Ag$ for some $g \in L_1(\Omega; X)$. Then, since $\int_0^t T(s) f ds = T(t)g - g$ for $t > 0$, it follows that

$$(23) \quad M_1 = \sup_{t > 0} t \|C_t^1(T(\cdot))f\|_1 = \sup_{t > 0} \|T(t)g - g\|_1 \leq 2\|g\|_1 < \infty.$$

Thus, (I) holds for $\alpha = 1$. If $\alpha > 1$, then by Fubini's theorem

$$\begin{aligned} \int_0^t (t-s)^{\alpha-1} T(s) f ds &= \int_0^t \left(\int_s^t (\alpha-1)(t-r)^{\alpha-2} T(s) f dr \right) ds \\ &= (\alpha-1) \int_0^t (t-r)^{\alpha-2} \left(\int_0^r T(s) f ds \right) dr, \end{aligned}$$

and thus

$$\begin{aligned} \left\| \int_0^t (t-s)^{\alpha-1} T(s) f ds \right\|_1 &\leq (\alpha-1) \int_0^t (t-r)^{\alpha-2} \left\| \int_0^r T(s) f ds \right\|_1 dr \\ &\leq (\alpha-1) \int_0^t (t-r)^{\alpha-2} M_1 dr = M_1 t^{\alpha-1}, \end{aligned}$$

so that

$$\sup_{t > 0} t \|C_t^\alpha(T(\cdot))f\|_1 \leq \alpha M_1.$$

This completes the proof of Theorem 3. □

Remarks (on Theorem 3). **(d)** The implication (III) \Rightarrow (I) does not hold for $0 \leq \alpha < 1$. Indeed, let $L_1(-\infty, \infty)$ be the usual L_1 -space of scalar-valued functions on the real line $\mathbb{R} := (-\infty, \infty)$. Let $T(t)$, $t \in \mathbb{R}$, be the operators on $L_1(-\infty, \infty)$ defined by

$$(24) \quad T(t)f(x) := f(x + t).$$

Then $T(\cdot) := \{T(t) : t \geq 0\}$ is a C_0 -semigroup of positive invertible linear isometries on $L_1(-\infty, \infty)$. The following are well-known:

- (i) $D(A) = \{g \in L_1(-\infty, \infty) : g \text{ is absolutely continuous, and } g' \in L_1(-\infty, \infty)\}$;
- (ii) $Ag = g'$ for $g \in D(A)$.

Hence the function $f = \chi_{[0,1]} - \chi_{[1,3]} + \chi_{[3,4]}$ belongs to $D(A)$, and $f = Ag$, where $g(x) := \int_{-\infty}^x f(s) ds$.

Now, suppose $0 < \alpha < 1$. Then, for every x with $0 < t - 1 < x < t$, we have

$$\begin{aligned} tC_t^\alpha(T(\cdot))f(-x) &= \alpha t^{1-\alpha} \int_0^t (t-s)^{\alpha-1} f(s-x) ds \\ &= \alpha t^{1-\alpha} \int_x^t (t-s)^{\alpha-1} ds = t^{1-\alpha}(t-x)^\alpha, \end{aligned}$$

whence

$$\|tC_t^\alpha(T(\cdot))f\|_1 \geq t^{1-\alpha} \int_{t-1}^t (t-x)^\alpha dx = t^{1-\alpha} \frac{1}{\alpha+1}.$$

This implies $\lim_{t \rightarrow \infty} \|tC_t^\alpha(T(\cdot))f\|_1 = \infty$, because $1 - \alpha > 0$. Next, suppose $\alpha = 0$. Then, clearly, we have $\|tC_t^0(T(\cdot))f\|_1 = t\|T(t)f\|_1 = t\|f\|_1 = 4t \rightarrow \infty$ as $t \rightarrow \infty$.

(e) The computations in the proof of (I) \Rightarrow (II) apply to the case $\alpha > 0$, so that the implication (I) \Rightarrow (II) holds for all $\alpha > 0$. But, if $\alpha = 0$, then the implication (I) \Rightarrow (II) fails to hold in general. This can be seen by modifying the example in Remark (b). Indeed, let w be the function on \mathbb{R} defined by $w(x) = 1$ if $x \geq -1$, and $w(x) = (-x)^{-1}$ if $x < -1$, and let μ be the measure on \mathbb{R} defined by $\mu = w dx$, where dx stands for the Lebesgue measure on \mathbb{R} . Then the operators $T(t)$, $t \geq 0$, of the form $T(t)f(x) = f(x + t)$ define a C_0 -semigroup $T(\cdot)$ of positive linear contractions on $L_1(\mathbb{R}, \mu)$ of scalar-valued integrable functions with respect to μ , and the function $f := \chi_{[0,1]}$ satisfies

$$\sup_{t>0} t\|C_t^0(T(\cdot))f\|_1 = \sup_{t>0} t\|T(t)f\|_1 = \sup_{t>0} t\|\chi_{[-t, -t+1]}\|_1 < 2.$$

But, it is known that

- (i) $D(A) = \{g \in L_1(\mathbb{R}, \mu) : g \text{ is locally absolutely continuous, and } g' \in L_1(\mathbb{R}, \mu)\}$;
- (ii) $Ag = g'$ for $g \in D(A)$.

Thus, if $f = Ah$ for some $h \in D(A)$, then we must have $f = \chi_{[0,1]} = h'$, and

$$h(x + t) - h(x) = \int_0^t f(x + s) ds \quad (t \geq 0, x \in \mathbb{R}).$$

Therefore, $h(x) = h(1)$ for $x \geq 1$, and $h(x) = h(0)$ for $x \leq 0$. But, since $h(1) - h(0) = \int_0^1 h'(s) ds = 1$, this proves that h cannot belong to $L_1(\mathbb{R}, \mu)$, a contradiction. Thus, $f \notin R(A)$, and (II) does not hold by Theorem 3.

Lastly, we give an application to the infinitesimal generator A of a strongly continuous cosine family $C(\cdot) \equiv \{C(t) : t \in \mathbb{R}\}$ of linear contractions on $L_1(\Omega; X)$. By definition, the family $C(\cdot)$ satisfies

- (i) $C(s + t) + C(s - t) = 2C(s)C(t)$ for all $s, t \in \mathbb{R}$;
- (ii) $C(0) = I$;
- (iii) $C(t)f$ is continuous in $t \in \mathbb{R}$ for each $f \in L_1(\Omega; X)$.

The infinitesimal generator A is defined by $Af := \lim_{t \rightarrow 0} (C(2t)f - f)/2t^2$, with domain $D(A)$ the set of all those $f \in L_1(\Omega; X)$ for which this limit exists. Since $\|C(t)\| \leq 1$ for all $t \in \mathbb{R}$, it is known (cf. e.g. [13], [14]) that A is a densely defined closed operator such that if $\lambda > 0$, then $\lambda \in \rho(A)$ and $\lambda(\lambda^2 - A)^{-1}f = \int_0^\infty e^{-\lambda s} C(s)f ds$ for all $f \in L_1(\Omega; X)$. Therefore we have $\sup_{\lambda > 0} \|\lambda(\lambda - A)^{-1}\| \leq 1$.

The associated sine family $S(\cdot) \equiv \{S(t) : t \in \mathbb{R}\}$ of linear operators on $L_1(\Omega; X)$ is defined by

$$(25) \quad S(t)f := \int_0^t C(s)f ds \quad (t \in \mathbb{R}, f \in L_1(\Omega; X)).$$

Elementary properties of $S(\cdot)$ and $C(\cdot)$ can be found in [14]. The Cesàro means of order γ (or γ -Cesàro means) $C_t^\gamma(S(\cdot))$ of the sine family $S(\cdot)$, where $\gamma \geq 0$ and $t > 0$, are the operators defined by $C_t^0(S(\cdot)) := S(t)$, and

$$(26) \quad C_t^\gamma(S(\cdot))f := \gamma t^{-\gamma} \int_0^t (t - s)^{\gamma-1} S(s)f ds \quad (\gamma > 0, f \in L_1(\Omega; X)).$$

It is direct to see that (18), (19) and (20) hold with $S(\cdot)$, $S(r)$, $S(s_n)$ and $S(t)$ in place of $T(\cdot)$, $T(r)$, $T(s_n)$ and $T(t)$, respectively.

Theorem 4 (cf. Corollary 8 of [11]). *Let X be a reflexive Banach space, and A be the infinitesimal generator of a strongly continuous cosine family $C(\cdot)$ of linear contractions on $L_1(\Omega; X)$. Assume that $\alpha \geq 1$. Then the following conditions are equivalent for $f \in L_1(\Omega; X)$.*

- (I) $\sup_{t > 0} t \|C_t^\alpha(S(\cdot))f\|_1 < \infty$.
- (II) $\sup_{\lambda > 0} \|\int_0^\infty e^{-\lambda t} S(t)f dt\|_1 < \infty$.
- (III) $f \in R(A)$.

PROOF: (I) \Rightarrow (II). We first notice, as in the proof of (I) \Rightarrow (II) of Theorem 3, that there exists an integer $n \geq \alpha$ such that

$$(27) \quad M'_n := \sup_{t>0} t \|C_t^n(S(\cdot))f\|_1 < \infty.$$

Then, since

$$\int_0^\infty e^{-\lambda t} S(t)f \, dt = \lambda^n \int_0^\infty e^{-\lambda t} \frac{t^{n-1}}{n!} \cdot [tC_t^n(S(\cdot))f] \, dt$$

(cf. (19) with $S(\cdot)$ and $S(s_n)$ in place of $T(\cdot)$ and $T(s_n)$, respectively), it follows that

$$\left\| \int_0^\infty e^{-\lambda t} S(t)f \, dt \right\|_1 \leq \frac{M'_n}{n!} \lambda^n \int_0^\infty e^{-\lambda t} t^{n-1} \, dt = \frac{M'_n}{n} \quad (\lambda > 0).$$

(II) \Rightarrow (III). Since $(\lambda^2 - A)^{-1}f = \lambda^{-1} \int_0^\infty e^{-\lambda s} C(s)f \, ds = \int_0^\infty e^{-\lambda t} S(t)f \, dt$ for $\lambda > 0$, (II) implies

$$\sup_{\lambda>0} \|(\lambda^2 - A)^{-1}f\|_1 = \sup_{\lambda>0} \left\| \int_0^\infty e^{-\lambda t} S(t)f \, dt \right\|_1 < \infty.$$

Hence (III) follows from Theorem 1.

(III) \Rightarrow (I). Assume that $f = Ag$ for some $g \in L_1(\Omega; X)$. By Lemma 2.15 of [13] we have $\int_0^t S(s)f \, ds = \int_0^t S(s)Ag \, ds = C(t)g - g$ for $t > 0$. Thus $M'_1 = \sup_{t>0} \left\| \int_0^t S(s)f \, ds \right\|_1 \leq 2\|g\|_1$, and hence (I) holds for $\alpha = 1$. If $\alpha > 1$, then we can obtain, as in the proof of (III) \Rightarrow (I) of Theorem 3, that $\sup_{t>0} t \|C_t^\alpha(S(\cdot))f\|_1 \leq \alpha M'_1$.

This completes the proof of Theorem 4. □

Remarks (on Theorem 4). (f) The implication (III) \Rightarrow (I) does not hold for $0 \leq \alpha < 1$. Indeed, if $C(t)$, $t \in \mathbb{R}$, are the operators on $L_1(-\infty, \infty)$ defined by $C(t)f(x) := 2^{-1}(f(x+t) + f(x-t))$, then $C(\cdot) := \{C(t) : t \in \mathbb{R}\}$ becomes a strongly continuous cosine family of positive linear contractions on $L_1(-\infty, \infty)$. It is known (cf. e.g. [13, Theorem 4.12]) that

$$(i) \quad D(A) = \left\{ g \in L_1(-\infty, \infty) : \begin{array}{l} g \text{ and } g' \text{ are absolutely continuous,} \\ \text{and } g', g'' \in L_1(-\infty, \infty) \end{array} \right\};$$

$$(ii) \quad Ag = g'' \text{ for } g \in D(A).$$

Thus the function

$$f = \chi_{[0,1)} - \chi_{[1,3)} + \chi_{[3,4)} - \chi_{[4,5)} + \chi_{[5,7)} - \chi_{[7,8)}$$

belongs to $R(A)$. Since $S(t)f(x) = \int_0^t 2^{-1}(f(x+s) + f(x-s)) ds$, it follows that if $t > 2$ then

$$(28) \quad S(t)f(x) \geq 1/4 \quad \text{for all } x \in [-t + (1/2), -t + (3/2)],$$

and thus if $s \in [t - (1/4), t]$, then, for all $x \in [-t + (3/4), -t + (3/2)]$,

$$(29) \quad S(s)f(x) = \int_0^s 2^{-1}(f(x+r) + f(x-r)) dr \geq 1/4.$$

Now, suppose $0 < \alpha < 1$. Then by (29), for $t > 2$ and $x \in [-t + (3/4), -t + (3/2)]$ we have

$$\begin{aligned} tC_t^\alpha(S(\cdot))f(x) &= \alpha t^{1-\alpha} \int_0^t (t-s)^{\alpha-1} S(s)f(x) ds \\ &\geq \alpha t^{1-\alpha} \int_{t-\frac{1}{4}}^t (t-s)^{\alpha-1} \cdot \frac{1}{4} ds = \frac{t^{1-\alpha}}{4^{\alpha+1}}, \end{aligned}$$

therefore

$$t\|C_t^\alpha(S(\cdot))f\|_1 \geq \int_{-t+(3/4)}^{-t+(3/2)} \frac{t^{1-\alpha}}{4^{\alpha+1}} dx = \frac{3t^{1-\alpha}}{4^{\alpha+2}} \longrightarrow \infty \quad (t \rightarrow \infty).$$

Next, suppose $\alpha = 0$. Then by (28) we get

$$t\|C_t^0(S(\cdot))f\|_1 = t\|S(t)f\|_1 \geq t \int_{-t+(1/2)}^{-t+(3/2)} \frac{1}{4} dx = \frac{t}{4} \longrightarrow \infty \quad (t \rightarrow \infty).$$

(g) The implication (I) \Rightarrow (II) of Theorem 4 holds for all $\alpha > 0$, as observed in Remark (e). Here it may be of some interest to note that if $\alpha = 0$ then the implication (I) \Rightarrow (II) is trivial. Indeed, if (I) holds for $\alpha = 0$, then we have $\sup_{t>0} t\|S(t)f\|_1 < \infty$ and hence $\lim_{t \rightarrow \infty} \|S(t)f\|_1 = 0$, from which we deduce that $f = 0$ as follows. For a moment, assume that $f \neq 0$. Then there exists $s_0 > 0$ such that $g := S(s_0)f \neq 0$. Then by Proposition 2.1 of [14]

$$C(t)g = C(t)S(s_0)f = 2^{-1}(S(t+s_0)f - S(t-s_0)f) \longrightarrow 0 \quad \text{as } t \rightarrow \infty,$$

and thus $g = -C(2t)g + 2C(t)^2g \rightarrow 0$ as $t \rightarrow \infty$. But this is a contradiction. (This proof was communicated to the author by Professor S.-Y. Shaw.)

REFERENCES

- [1] Assani I., *A note on the equation $y = (I - T)x$ in L^1* , Illinois J. Math. **43** (1999), 540–541.
- [2] Browder F.E., *On the iteration of transformations in non-compact minimal dynamical systems*, Proc. Amer. Math. Soc. **9** (1958), 773–780.
- [3] Diestel J., Uhr J.J., Jr., *Vector Measures*, Amer. Math. Soc., Providence, 1977.
- [4] Fonf V., Lin M., Rubinov A., *On the uniform ergodic theorem in Banach spaces that do not contain duals*, Studia Math. **121** (1996), 67–85.
- [5] Gottschalk W.H., Hedlund G.A., *Topological Dynamics*, Amer. Math. Soc. Colloq. Publ. **36**, Amer. Math. Soc., Providence, 1955.
- [6] Krengel U., Lin M., *On the range of the generator of a Markovian semigroup*, Math. Z. **185** (1984), 553–565.
- [7] Li Y.-C., Sato R., Shaw S.-Y., *Boundedness and growth orders of means of discrete and continuous semigroups of operators*, preprint.
- [8] Lin M., Sine R., *Ergodic theory and the functional equation $(I - T)x = y$* , J. Operator Theory **10** (1983), 153–166.
- [9] Pazy A., *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer, New York, 1983.
- [10] Sato R., *Solvability of the functional equation $f = (T - I)h$ for vector-valued functions*, Colloq. Math. **99** (2004), 253–265.
- [11] Shaw S.-Y., *On the range of a closed operator*, J. Operator Theory **22** (1989), 157–163.
- [12] Shaw S.-Y., Li Y.-C., *On solvability of $Ax = y$, approximate solutions, and uniform ergodicity*, Rend. Circ. Mat. Palermo (2) Suppl. **2002**, no. 68, part II, 805–819.
- [13] Sova M., *Cosine operator functions*, Rozprawy Mat. **49** (1966), 1–47.
- [14] Travis C.C., Webb G.F., *Cosine families and abstract nonlinear second order differential equations*, Acta Math. Acad. Sci. Hungar. **32** (1978), 75–96.
- [15] Zygmund A., *Trigonometric Series*, Vol. I, Cambridge University Press, Cambridge, 1959.

DEPARTMENT OF MATHEMATICS, OKAYAMA UNIVERSITY, OKAYAMA, 700-8530 JAPAN

E-mail: satoryot@math.okayama-u.ac.jp

(Received June 7, 2004, revised January 13, 2005)