

Petr Hájek

Automatic listing of important observational statements. III

Kybernetika, Vol. 10 (1974), No. 2, (95)--124

Persistent URL: <http://dml.cz/dmlcz/124863>

Terms of use:

© Institute of Information Theory and Automation AS CR, 1974

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these

Terms of use.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*
<http://project.dml.cz>

Automatic Listing of Important Observational Statements III

PETR HÁJEK

The ALIOS theory (of Automatic Listing of Important Observational Statements) was introduced and developed in the previous papers of this series. In the present paper, which is a free continuation, we study an important particular class of observational functor calculi and develop its logic.

Part III — Nominal calculi with incomplete information

INTRODUCTION

In [2] we introduced the notions of a semantic system, a problem and of a solution of a problem in a model; in [3] we studied functor calculi and corresponding semantic system, in particular, observational monadic functor calculi (OMFC's, see also below). In [4] we called the problematics of the ALIOS theory the problematic of *Automated Research*; this seems to be an acceptable term (first of all since it is short). It can be said that the logic of Automated Research is the *logic of observational functor calculi*, especially (at least at the present stage), of OMFC's. In the present part we are going to study OMFC's with nominal values and incomplete information, briefly, \times -nominal calculi (see below); they form an important particular class of OMFC's including the classical observational monadic predicate calculus. The generalization of the classical calculus consists in generalizing (i) truth values to more general values, (ii) connectives to some junctors and (iii) quantifiers to some operators. Values are generalized only slightly and in two directions: we allow values $0, 1, \dots, h$ corresponding to some nominal quantities and also the values \times — "unknown". (Cf. [3] 8.10–11 and [5].)*

* I promised in [3] a paper denoted there by [12] on nominal quantities; what I planned to write in that paper is included here. I also present here some results of [4].

The main attention is paid to the study of some particular classes of operators. The value \times is treated in Kleene-Körner's style as it was in [5]. (Cf. [6]; note that Kleene-Körner calculus is studied in the recent paper by Cleave [1].) The reader interested only in the logical aspects of the present paper but not in Automated Research should consult § 1 of [2] and § 7 of [3] for terminology.

12. BASIC DEFINITIONS; OPEN FORMULAS

OMFC's are functor calculi whose all models are finite, all associated functions are calculable (say, recursive), all functors are unary, there is only one variable (omitted at all occurrences) and (an added condition) all operators are of a type $\langle 1, \dots, 1 \rangle$ (i.e. the operator binds the variable in each of the joined formulas).

12.1. Definition. Let $h \geq 1$. Numbers $0, 1, \dots, h$ are called *regular values* ($V_{reg} = \{0, \dots, h\}$); the symbol \times is the *singular value*. So $V = V_{reg} \cup \{\times\}$. We put $0 < \times < 1$ (one might identify \times e.g. with $\frac{1}{2}$). Fix a type $\langle \underbrace{1, \dots, 1}_{n \text{ times}} \rangle$ (say, 1^n). A

V -structure $\langle M, f_1, \dots, f_n \rangle = \mathbf{M}$ of this type is *quantitative* if the ranges of all f 's are included in $\{0, \times, 1\}$; \mathbf{M} is a structure with *complete information* if all f 's take only regular values. A structure $\mathbf{M}' = \langle M, g_1, \dots, g_n \rangle$ with complete information is a *completion* of $\mathbf{M} = \langle M, f_1, \dots, f_n \rangle$ if, for each $a, i, f_i(a) \in V_{reg}$ implies $f_i(a) = g_i(a)$.

12.2. Remarks. (1) A *partial V_{reg} -structure* is a tuple $\langle M, h_1, \dots, h_n \rangle$ where each h_i is a mapping whose domain is a subset of M and whose range is a subset of V_{reg} . There is an obvious one-one correspondence between V -structures and partial V_{reg} -structures ($\langle M, f_1, \dots, f_n \rangle$ corresponds to $\langle M, h_1, \dots, h_n \rangle$ iff, for each i, f_i extends h_i and f_i takes the value \times on $Dom(f) - Dom(h)$).

(2) A *partition* in M is a set p of pairwise disjoint nonempty subsets of M . A *partitional structure* is a tuple $\langle M, p_1, \dots, p_n \rangle = \mathbf{M}$ where p_i 's are partitions in M ; we suppose that the cardinality of each p_i is $\leq h + 1$. Let, for each i, e_i be a one-one mapping of p_i into $\{0, \dots, h\}$ (*enumeration*). Then $\langle \mathbf{M}, e \rangle = \langle M, p_1, e_1, \dots, p_n, e_n \rangle$ is an *enumerated partitional structure* called an *enumeration of \mathbf{M}* . There is an obvious one-one correspondence between all enumerated partitional structures and all V -structures ($\langle M, f_1, \dots, f_n \rangle$ corresponds to $\langle M, p_1, e_1, \dots, p_n, e_n \rangle$ iff $f_i(a) = e_i(a)$ for $a \in A \in p_i$ and $f_i(a) = \times$ for $a \in M - \cup p_i$).

(3) There is another obvious one-one correspondence between all V -structures and all qualitative structures $\langle M, q_1, \dots, q_{n,h} \rangle$ of the type $1^{n \cdot h}$ such that the following holds for each i and each $a \in M$: if one of $q_{i,h+1}(a), \dots, q_{i(h+1)}(a)$ is \times then all of them; if no member of the above sequence is \times then exactly one member is 1. (Call such a structure a *disjointed structure* of the type $\langle n; h \rangle$.)

12.3. Discussion. It follows from the intuitive notion of a nominal quantity (as a quantity whose values merely enumerate factors of an equivalence relation) that speaking about a \times -nominal V -structure \mathcal{M} we in fact want to speak about a partial structure \mathcal{M}_0 such that \mathcal{M} corresponds to an enumeration of \mathcal{M}_0 . This will influence the choice of our language. Note that theoretically we could deal only with qualitative structures, namely with disjointed structures of the type $\langle n; h \rangle$ but practically we prefer V -structures e.g. for saving place in the computer's memory, (See also 12.7.)

12.4. Definition. (1) Take functors F_1, \dots, F_n . We introduce a unary junctor (X) (called a *coefficient*) for each $X \subseteq V_{reg}$ putting $Asf_{(X)}(u) = 1$ iff $u \in X$, $Asf_{(X)}(u) = 0$ iff $u \in V_{reg} - X$ and $Asf_{(X)}(u) = \times$ iff $u = \times$.

(2) A function $\alpha : V^j \rightarrow V$ is *qualitative* if there is a function $\alpha_0 : \{0, \times, 1\}^j \rightarrow \{0, \times, 1\}$ such that $\alpha(u_1, \dots, u_j) = \alpha_0(\bar{u}_1, \dots, \bar{u}_j)$, where $\bar{0} = 0$, $\bar{\times} = \times$ and $\bar{1} = 1$ for $u \geq 1$. We say that α *corresponds* to α_0 . (Evidently, for each $\alpha_0 : \{0, \times, 1\}^j \rightarrow \{0, \times, 1\}$ the corresponding operation on V is determined uniquely.)

(3) We introduce junctors $\neg, \vee, \&$ by associated functions corresponding to the respective three-valued associated functions (see [5], [6] or [1]), e.g. $Asf_{\&}(u, v) = 1$ iff $u \geq 1$ and $v \geq 1$, $Asf_{\&}(u, v) = 0$ iff $u = 0$ or $v = 0$ and $Asf_{\&}(u, v) = \times$ otherwise.

(4) (Auxiliary.) For a while, call any OMFC whose language has the functors F_1, \dots, F_n and the junctors just defined an *openly \times -nominal OMFC*. (Nothing is assumed on operators; we study open formulas in such OMFC's.)

(5) Each formula of the form $(X)F$ where F is a functor and $0 \subset X \subset V$ is called a *literal*. An *elementary disjunction* (ED) is a disjunction of literals in which each functor occurs at most once; similarly we define *elementary conjunctions* (EC).

(6) Two open formulas φ, ψ are *strongly equivalent* in \mathcal{M} if $\|\varphi\|_{\mathcal{M}}(a) = \|\psi\|_{\mathcal{M}}(a)$ for each $a \in M$. (Denotation: $\varphi \cong_{\mathcal{M}} \psi$.)

(7) A formula is *qualitative* if $(\forall \mathcal{M}) (\forall a \in M) (\|\varphi\|_{\mathcal{M}}(a) \in \{0, \times, 1\})$.

(8) Let φ, ψ be qualitative open formulas. φ *strongly implies* ψ in \mathcal{M} if $\|\varphi\|_{\mathcal{M}}(a) \leq \|\psi\|_{\mathcal{M}}(a)$ for each $a \in M$. (Denotation: $\varphi \supseteq_{\mathcal{M}} \psi$, cf. [1].)

12.5. Remark. (1) Each non-atomic open formula is qualitative. (2) $\|F\|_{\mathcal{M}}(a) = \times$ iff $\|(X)F\|_{\mathcal{M}}(a) = \times$ for each coefficient (X) .

(3) If φ, ψ are qualitative open then $\varphi \cong_{\mathcal{M}} \psi$ iff $(\varphi \supseteq_{\mathcal{M}} \psi$ and $\psi \supseteq_{\mathcal{M}} \varphi)$.

12.6. Lemma. (1) $\neg(X)\varphi \cong_{\mathcal{M}} (V_{reg} - X)\varphi$,

(2) $(X)\varphi \cong_{\mathcal{M}} \bigvee_{k \in X} (\{k\})\varphi$ for $X \neq \emptyset$,

(3) $(X)\varphi \cong_{\mathcal{M}} \bigvee_{k \notin X} \neg(\{k\})\varphi$ for $X \neq V_{reg}$,

(4) $(X)\varphi \vee (Y)\varphi \cong_{\mathcal{M}} (X \cup Y)\varphi$,

(5) $(X)\varphi \& (Y)\varphi \cong_{\mathcal{M}} (X \cap Y)\varphi$.

Easy proofs are left to the reader.

12.7. Discussion. Call a mapping of M into $\{0, \times, 1\}$ a *partial subset* of M . Let \mathbf{M} be a V -structure and let N be the corresponding disjointed structure of the type $\langle n; h \rangle$. Let \mathfrak{F} be an openly \times -nominal OMFC and let \mathfrak{B} be the three-valued observational monadic predicate calculus with $n \cdot h$ predicates. One easily shows that a partial subset of M is definable in \mathbf{M} by an open qualitative formula of \mathfrak{F} iff it is definable in N by an open formula of \mathfrak{B} . So theoretically one could deal with \mathfrak{B} ; but practically we prefer (openly) \times -nominal calculi not only because of the reasons of 12.3 but also since we have some reasonable notions of complexity of formulas of \mathfrak{F} not preserved by the transition to corresponding formulas of \mathfrak{B} ; in particular, ED's and EC's seem to be very natural open formulas (one can construct "normal forms" for non-atomic open formulas in the obvious way) and in 12.8 (4) we introduce a natural "simpler than" relation for EC's and ED's.

12.8. Definition. (1) An EC $\kappa = \bigwedge_{i \in I} (X_i) F_i$ is *poorer than* $\lambda = \bigwedge_{i \in I} (Y_i) F_i$ if for each $i \in I$ we have $X_i \subseteq Y_i$. $\kappa \subseteq \lambda$ means that κ is a subconjunction of λ .

(2) An EC κ is *incompressible* in \mathbf{M} (or \mathbf{M} -incompressible) if κ is in \mathbf{M} not strongly equivalent to any κ_0 strictly poorer than κ . κ is *incancellable* in \mathbf{M} if κ is in \mathbf{M} not strongly equivalent to any κ_0 which is a proper subconjunction of κ , κ is *prime* in \mathbf{M} if it is both incompressible and incancellable in \mathbf{M} .

(3) If κ is $\bigwedge_{i \in I} (X_i) F_i$ then $neg(\kappa)$ is $\bigvee_{i \in I} (V_{reg} - X_i) F_i$.

(4) Let $\kappa = \bigwedge_{i \in I} (X_i) F_i$ and $\lambda = \bigwedge_{i \in J} (Y_i) F_i$. We put $\kappa \triangleleft \lambda$ (κ is simpler than λ) if $I \subseteq J$ and for each $i \in I$ we have $X_i \subseteq Y_i$.

(5) We put $\kappa \triangleleft \lambda$ if $I \subseteq J$ and for each $i \in I$ we have $Y_i \subseteq X_i$.

(6) All the above definitions are made analogously for ED's.

12.9. Remark. If κ, λ are EC's if \mathbf{M} is a model and κ is poorer than λ then $\kappa \approx_{\mathbf{M}} \lambda$; if $\kappa \subseteq \lambda$ then $\lambda \approx_{\mathbf{M}} \kappa$. Hence if $\kappa \triangleleft \lambda$ then $\lambda \approx_{\mathbf{M}} \kappa$; analogously, if γ, δ are ED's and $\gamma \triangleleft \delta$ then $\gamma \approx_{\mathbf{M}} \delta$. If $\kappa \triangleleft \lambda$ (κ, λ EC's) then in general neither $\kappa \approx_{\mathbf{M}} \lambda$ nor $\lambda \approx_{\mathbf{M}} \kappa$. Nevertheless, we have the following

12.10. Lemma. If $\kappa_0 \cong_{\mathbf{M}} \kappa$ and $\kappa_0 \triangleleft \kappa$ then there is a κ_2 such that $\kappa_0 \subseteq \kappa_2$, κ_2 is poorer than κ and $\kappa \cong_{\mathbf{M}} \kappa_2$.

Proof. Let κ_0 be poorer than $\kappa_1 \subseteq \kappa$, $\kappa = \bigwedge_I (X_i) F_i$, $\kappa_1 = \bigwedge_{I_0} (X_i) F_i$, $\kappa_0 = \bigwedge_{I_0} (X_i^0) F_i$ ($I_0 \subseteq I$, $X_i^0 \subseteq X_i$). Put $X_i^0 = X_i$ for $i \in I - I_0$ and put $\kappa_2 = \bigwedge_I (X_i^0) F_i$. Then $\kappa_0 \subseteq \kappa_2$, κ_2 is poorer than κ ; hence if $\|\kappa_2\|_{\mathbf{M}}(a) = 1$ then $\|\kappa\|_{\mathbf{M}}(a) = 1$. If $\|\kappa\|_{\mathbf{M}}(a) = 1$ then $\|\kappa_1\|_{\mathbf{M}}(a) = 1$ and $\|\bigwedge_{I - I_0} (X_i) F_i\|_{\mathbf{M}}(a) = 1$, hence $\|\kappa_2\|_{\mathbf{M}}(a) = 1$. If $\|\kappa_2\|_{\mathbf{M}}(a) = \times$ then $\|\kappa\|_{\mathbf{M}}(a) \in \{\times, 1\}$; conversely, if $\|\kappa\|_{\mathbf{M}}(a) = \times$ then $\|\kappa_0\|_{\mathbf{M}}(a) = \times$ and $\|\bigwedge_{I - I_0} (X_i) F_i\|_{\mathbf{M}}(a) \in \{\times, 1\}$, so $\|\kappa_2\|_{\mathbf{M}}(a) = \times$.

12.11. Corollary. For each M and each κ , κ is M -prime iff there is no $\kappa_0 \neq \kappa$ such that $\kappa_0 \triangleleft \kappa$ and $\kappa_0 \cong_M \kappa$.

12.12. Lemma. (1) Each subconjunction of an M -incompressible EC is M -incompressible. (2) Each M -incompressible EC κ contains an M -prime subconjunction strongly equivalent to κ .

Proof. (1) follows immediately from 12.10; (2) is then evident.

13. OPERATORS; IMPROVEMENT OPERATORS

13.1. Definition. (1) Let \mathfrak{R} be a set of V -structures. A mapping $\alpha : \mathfrak{R} \rightarrow \{0, \times, 1\}$ is *qualitative* if for each $M = \langle M, f_1, \dots, f_n \rangle$ we have $\alpha(M) = \alpha(\langle M, \bar{f}_1, \dots, \bar{f}_n \rangle)$ where $f_i(a) = \bar{f}_i(a)$ for each $a \in M$ (cf. 12.4).

(2) Let \mathfrak{F} be an openly \times -nominal OMFC and let Q be an operator of \mathfrak{F} . Q is *qualitative* if its associated function is qualitative. (Evidently, such a function is determined by its values on qualitative models.)

13.2. Lemma. Q is qualitative iff, for each M , $\|Q(F_1, \dots, F_n)\|_M = \|Q((X)F_1, \dots, (X)F_n)\|_M$ where $X = V_{reg} - \{0\}$. (Since $\langle M, \|(X)F_1\|_M, \dots, \|(X)F_n\|_M \rangle$ is $\langle M, \bar{f}_1, \dots, \bar{f}_n \rangle$.)

13.3. Definition and Remark. (1) Introduce a qualitative operator \cong of the type 1^2 (*strong equivalence*) whose associated function takes for a qualitative model $M = \langle M, f_1, f_2 \rangle$ value 1 if $f_1 = f_2$, otherwise $Asf \cong(M) = 0$. Then, for qualitative open formulas φ, ψ we have: $\varphi \cong_M \psi$ iff $\|\varphi \cong \psi\|_M = 1$.

(2) We further introduce a qualitative operator \simeq of the type 1^2 (*weak equivalence*) whose associated function takes for a qualitative model $M = \langle M, f_1, f_2 \rangle$ value 1 if $(\forall a \in M)(f_1(a) = 1 \text{ iff } f_2(a) = 1)$ and otherwise $Asf \simeq(M) = 0$.

(3) Evidently, an EC κ is M -compressible iff there is a $\kappa_0 \neq \kappa$, κ_0 poorer than κ such that $\|\kappa_0 \cong \kappa\|_M = 1$. Analogously we define: κ is *weakly M -compressible* if there is a $\kappa_0 \neq \kappa$, κ_0 poorer than κ such that $\|\kappa_0 \simeq \kappa\|_M = 1$. Caution: the negation of “weakly M -compressible” is “*strongly M -incompressible*”.

13.4. Definition. An openly \times -nominal OMFC \mathfrak{F} is a \times -nominal OMFC (or an OMFC with *nominal values and incomplete information*) if all operators of \mathfrak{F} are qualitative.

13.5. Remark. This definition corresponds to our interest in qualitative formulas; Lemma 13.2 shows that every non-atomic formula (not necessarily open) is strongly equivalent to a qualitative formula.

13.6. Definition. Let \mathfrak{F} be a \times -nominal OMFC. An operator Q is *secured* if for each model M of the type of Q we have: $Asf_Q(M) = 1$ iff for each completion M' of M $Asf_Q(M') = 1$, $Asf_Q(M) = 0$ iff for each completion M' of M $Asf_Q(M') = 0$, $Asf_Q(M) = \times$ otherwise.

13.7. Remark. (1) Evidently, it suffices when the above conditions are satisfied for all qualitative models (since Q is qualitative, \mathfrak{F} being \times -nominal). Hence, the semantics of a secured operator is determined by its semantics on qualitative models with complete information.

(2) One often defines an operator using a real function $S(M)$ (e.g. a statistic) putting for $\{0, 1\}$ -models $Asf_Q(M) = 1$ if $S(M) \leq \alpha$ and $= 0$ otherwise. If one uses such a definition to define a secured operator, it is advantageous to have an effective procedure which, given a qualitative model M produces its completion for which the value $S(M_{\max})$ is maximal among all $S(M')$ for M' being a completion of M .

13.8. Discussion. The definition of a secured operator is inspired by the idea of a "heavenly model" (cf. [5]): we treat \times as a sign for missing information; if $\|\varphi\|_M(a) = \times$ then in fact the object a either has the property φ or has not; we only do not know what case occurs. Since the right "heavenly" completion is not available, wanting to be sure that a formula $Q(\varphi_1, \dots, \varphi_n)$ is true in it we must assure that it is true in all the completions. A philosophical investigation of the value \times in connection with "inexact classes" is contained in [6].

Non-secured operators are auxiliary from our point of view; but they can be quite helpful. For example, \cong is not secured. In this paper we shall study certain non-secured operators called improvement operators. We find important particular improvement operators in Sections 15 and 16.

In the rest of the present section we consider an arbitrary OMFC \mathfrak{F} whose set of values contains 0, 1. "Tautology" means $\{1\}$ -tautology etc.

13.9. Definition. Let H be an operator of the type 1", let Of be a (finite) set of n -tuples of open formulas, let \subseteq be an ordering of Of such that the supremum of any two elements exists; finally, let \ll be an operator of the type 1²ⁿ; Φ, Ψ, Ω denote elements of Of .

(1) \ll is a *closure operator* for Of and \subseteq if $\Phi \ll \Phi$ is a tautology for each Φ and if the following rules are sound:

$$\begin{array}{l}
 \text{(I)} \quad \left. \begin{array}{c} \frac{\Phi \ll \Psi, \Psi \ll \Omega}{\Phi \ll \Omega} \\ \\ \text{(II)} \quad \frac{\Phi \ll \Omega}{\Phi \ll \Psi, \Psi \ll \Omega} \end{array} \right\} \text{ for } \Phi \subseteq \Psi \subseteq \Omega,
 \end{array}$$

$$(III) \quad \frac{\Phi \ll \Psi_1, \Phi \ll \Psi_2}{\Phi \ll \Omega} \quad \text{for } \Omega = \text{sup}_{\subseteq}(\Psi_1, \Psi_2).$$

(2) \ll satisfies *modus ponens* w.r.t. H, Of, \subseteq if the following rule is sound:

$$\frac{H(\Phi), \Phi \ll \Psi}{H(\Psi)} \quad \text{for } \Phi \subseteq \Psi.$$

(3) \ll is an *improvement operator* for H, Of, \subseteq if it is a closure operator Of, \subseteq and satisfies *modus ponens* w.r.t. H, Of, \subseteq .

13.10. Lemma and Definition. (1) Let \ll be a closure operator for Of, \subseteq . Then for each $\Phi \in Of$ and each M there is a uniquely determined $\Psi \supseteq \Phi$ which is \subseteq -maximal such that $\|\Phi \ll \Psi\|_M = 1$; put $\Psi = \text{Reg} \ll_M(\Phi)$. $\text{Reg} \ll_M(\Phi)$ is the supremum of all $\Psi \subseteq \Phi$ such that $\|\Phi \ll \Psi\|_M = 1$.

(2) The following holds for each Φ, Ψ :

$$\Phi \subseteq \text{Reg} \ll_M(\Phi); \Phi \subseteq \Psi \subseteq \text{Reg} \ll_M(\Phi) \rightarrow \text{Reg} \ll_M(\Psi) = \text{Reg} \ll_M(\Phi).$$

Proofs are easy from the definition.

13.11. Remark. Suppose that $\langle Of, \subseteq \rangle$ has the following algebraic property: whenever $\Phi \subset \Psi \subset \Omega$ then there is a $\bar{\Psi}$ such that $\Phi \subset \bar{\Psi} \subset \Omega$, $\bar{\Psi}$ is an immediate successor of Φ and $\Psi, \bar{\Psi}$ are incomparable. Then for $\Phi \subset \Omega$, Ω is the supremum of all immediate successors Ψ of Φ such that $\Psi \subseteq \Omega$. In particular, $\text{Reg} \ll_M(\Phi)$ is the supremum of all the immediate successors Ψ of Φ such that $\|\Phi \ll \Psi\|_M = 1$ provided there are any; if not, $\text{Reg} \ll_M(\Phi) = \Phi$.

13.12. Theorem. Let \ll be an improvement operator for H, Of, \subseteq ; put $P = \langle F, \{1\}, IC \rangle$ where $F = \{H(\Phi); \Phi \in Of\}$ and IC is the following rule:

$$\frac{H(\Phi) \& \Phi \ll \Omega}{H(\Psi)} \quad \text{for } \Phi \subseteq \Psi \subseteq \Omega.$$

Then IC is sound and hence P is a problem; for each model M , call $H(\Phi)$ M -prime if $\|H(\Phi)\|_M = 1$ and there is no $\Psi \subset \Phi$ such that $\|H(\Psi)\|_M = 1$ and $\|\Psi \ll \Phi\|_M = 1$. Put $X = \{H(\Phi) \& \Phi \ll \Psi; H(\Phi) \text{ } M\text{-prime and } \Psi = \text{Reg} \ll_M(\Phi)\}$. Then X is a \subseteq -minimal solution of P in M .

Proof. $X \subseteq \text{Tr}(M)$ is trivial; if $\|H(\Omega)\|_M = 1$ then consider the set Pr_Ω of all $\Phi \subseteq \Omega$ such that $\|H(\Phi) \& \Phi \ll \Omega\|_M = 1$. Let Φ be a \subseteq -minimal element of Pr_Ω . Then $H(\Phi)$ is M -prime and $\Phi \subseteq \Omega \subseteq \text{Reg} \ll_M(\Phi)$. Hence if $\text{Reg} \ll_M(\Phi) = \Psi$ then $(H(\Phi) \& \Phi \ll \Psi)$ is in X and $H(\Omega)$ is an immediate consequence of $H(\Phi) \& \Phi \ll \Psi$. If $(H(\Phi) \& \Phi \ll \Psi) \in X$ then this formula is the only one from which $H(\Phi)$ immediately follows; so X is \subseteq -minimal.

13.13. Remarks. (1) Compare theorem 13.12 with [3] 9.3–9.9. One can form a hierarchicity statement as in 9.4: Let

$$\begin{aligned} cf(H(\Phi)) &= \Phi, \quad cf(H(\Phi) \& \Phi \ll \Psi) = \Phi, \\ Sent &= \{H(\Phi); \Phi\} \cup \{H(\Phi) \& \Phi \ll \Psi; \Phi \subseteq \Psi\}. \end{aligned}$$

Let, for $\Sigma_1, \Sigma_2 \in Sent$, $\Sigma_1 \leq_R \Sigma_2$ iff $cf(\Sigma_1) \subseteq cf(\Sigma_2)$. Let H be a hierarchy on $Sent$ such that $R \subseteq R_H$. Then $\{X \cap h; h \in H\}$ is a solution of the hierarchical problem $\langle P, H \rangle$. (The proof is routine; cf. [2] 5.4 and 5.9.)

(2) Let \ll_1 and \ll_2 be two improvement operators for H, Of, \subseteq ; define problems P_1, P_2 and solutions X_1, X_2 as above. \ll_1 is said to be stronger than \ll_2 if $(\Phi \ll_2 \Psi) : (\Phi \ll_1 \Psi)$ is a sound rule. Evidently, if \ll_1 is stronger than \ll_2 then $card(X_1) \leq card(X_2)$ so that X is better.

(3) Improvement operators can be used also to improve solutions of problems having already a non-trivial relation of immediate consequence; one strengthens the relation using properties of improvement operators and diminishes the cardinality of solutions. An example will be considered in Section 16.

(4) Let us mention here that various notions of minimality of a solution do not formalize all intuitive criteria for a solution to be good but only some of them. We want the solution (as the machine output e.g. in printed form) to be both as small as possible and as transparent as possible. Preference of a certain (sort of a) solution can depend also on one's taste.

14. CLOSURE OPERATORS

Closure operators were defined in 13.9; we shall now study some particular cases

14.1. Definition. A qualitative operator Q of the type 1^n is *universally definable* if there is a set $U \subseteq \{0, \times, 1\}^n$ such that, for each qualitative model $M = \langle M, f_1, \dots, f_n \rangle$ of the type 1^n we have the following:

$$\begin{aligned} Asf_Q(M) &= 1 \quad \text{iff} \quad (\forall a \in M) \langle f_1(a), \dots, f_n(a) \rangle \in U, \\ Asf_Q(M) &= 0 \quad \text{otherwise.} \end{aligned}$$

(In words: $Asf_Q(M) = 1$ iff all the cards in M belong to U .)

14.2. Remark. The set U defines a junctor $\iota(Asf_\iota(u_1, \dots, u_n) = 1$ iff $\langle u_1, \dots, u_n \rangle \in U$, $= 0$ otherwise) such that, for each M , $Q(\varphi_1, \dots, \varphi_n) \cong_M \forall \iota(\varphi_1, \dots, \varphi_n)$ (\forall is the universal quantifier). We restrict ourselves to universally definable closure operators.

14.3. Definition. If \ll is a closure operator (for Of, \subseteq) and if \ll is universally defined by U then U is called a *closure set* (for Of, \subseteq).

14.4. Definition. (1) A *pseudoliteral* is a formula of the form $(X)F$ where $X \subset V_{reg}$ ($X = \emptyset$ is not excluded). Pseudoelementary conjunctions and disjunctions (psEC, psED) are defined from pseudoliterals as EC's and ED's from literals.

(2) The orderings \triangleleft , \triangleleft_e are extended in the obvious manner to psEC's and psED's.

14.5. Lemma. If $\kappa = \bigwedge_I (X_i)F_i$ and $\lambda = \bigwedge_J (Y_i)F_i$ are psEC's then, in the sense of \triangleleft_e , the supremum of κ , λ is $\bigwedge_{I \cup J} (Z_i)F_i$ where $Z_i = X_i \& Y_i$ for $i \in I \cap J$, $Z_i = X_i$ for $i \in I - J$ and $Z_i = Y_i$, for $i \in J - I$; for each M , $\sup(\kappa, \lambda) \cong_M \kappa \& \lambda$.

14.6. Theorem. (1) The strong equivalence operator \cong is a universally definable closure operator for psAC's and \triangleleft_e .

(2) If κ is a psEC then $Reg \cong_M(\kappa)$ is M -incompressible.

(3) If κ is an EC satisfiable in M (i.e. $(\exists a \in M) (\|\kappa\|_M(a) = 1)$) then $Reg \cong_M(\kappa)$ is an EC (no coefficient is empty).

(4) The weak equivalence operator \cong_e is a universally definable closure operator for psEC's and \triangleleft_e .

(5) If κ is a psEC then $Reg \cong_e(\kappa)$ is strongly M -incompressible (i.e. not weakly M -compressible).

(6) If κ is an EC satisfiable in M then $Reg \cong_e(\kappa)$ is an EC.

Proof. (1) It follows from the definition using 14.5 that is a closure operator for psEC's and \triangleleft_e (observe that $\kappa \triangleleft_e \lambda$ implies $\|\kappa\|_M(a) \cong \|\lambda\|_M(a)$ for each M and each $a \in M$. Clearly, \cong is universally defined by $U = \{\langle 0, 0 \rangle, \langle \times, \times \rangle, \langle 1, 1 \rangle\}$.

(2) Let $Reg \cong_M(\kappa)$ be $\kappa_1 \& (X)F$ and suppose $\|\kappa_1 \& (X)F\|_M = 1$ for some $X_0 \subseteq X$. Then, since $\|\kappa \cong \kappa_1 \& (X)F\|_M = 1$, we have $\|\kappa \cong \kappa_1 \& (X_0)F\|_M = 1$, hence $\kappa_1 \& (X_0)F \triangleleft_e \kappa_1 \& (X)F$ by the properties of Reg . Consequently, $X \subseteq X_0$.

(3) Suppose $Reg \cong_M(\kappa)$ to be $\kappa_1 \& (\emptyset)F$ and let $\|\kappa\|_M(a) = 1$. Then either $\|F\|_M(a) \in \{0, 1\}$, $\|(\emptyset)F\|_M(a) = 0$ and $\|\kappa_1 \& (\emptyset)F\|_M(a) = 0$ or $\|F\|_M(a) = \times$ and $\|\kappa_1 \& (\emptyset)F\|_M(a) = \times$; in either case we arrive at a contradiction with $\kappa \cong_M Reg \cong_M(\kappa)$.

Proofs of (4)–(6) are similar.

14.7. Corollary. For each psEC κ and each M , (1) there is a uniquely determined psEC κ_0 poorer than κ , M -incompressible and such that $\kappa_0 \cong_M \kappa$;

(2) there is a uniquely determined psEC κ_1 poorer than κ , strongly M -incompressible and such that $\|\kappa_0 \cong \kappa\|_M = 1$. (The last fact could be denoted by $\kappa_0 \cong_M \kappa$ if you like.)

Proof. (1) Take the subconjunction κ_0 of $Reg \cong_M(\kappa)$ containing exactly the same functors as κ ; then κ_0 is poorer than κ (since $\kappa \triangleleft_e Reg \cong_M(\kappa)$), κ_0 is M -incompressible by 12.12 and $\kappa \cong_M Reg \cong_M(\kappa) \supseteq_M \kappa_0 \supseteq_M \kappa$, hence $\kappa \cong_M \kappa_0$. The uniqueness is obvious (e.g. by 14.5).

14.8. Remark and Definition. In the sequel, we shall be interested in closure operators for sets of pairs of open formulas, in particular, in the set CC of pairs $\langle \kappa, \lambda \rangle$ of psEC's and in the set CD of pairs $\langle \kappa, \delta \rangle$ where κ is a psEC and δ is a psED. (We could consider also DC and DD in the obvious meaning and way.) We define \ll for elements of CC as the "product" ordering: $\langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle$ if $\kappa \ll \bar{\kappa}$ and $\lambda \ll \bar{\lambda}$. Similarly, $\langle \kappa, \delta \rangle (\ll \ll) \langle \bar{\kappa}, \bar{\delta} \rangle$ if $\kappa \ll \bar{\kappa}$ and $\delta \ll \bar{\delta}$.

By a *CC-closure set* we mean a closure set U for CC and \ll satisfying the following additional condition: $\langle u, v, \bar{u}, \bar{v} \rangle \in U$ implies $u \geq \bar{u}, v \geq \bar{v}$ for each u, v, \bar{u}, \bar{v} . A *CC-closure operator* is an operator defined universally by a CC-closure set. (The condition is natural since having a closure operator \ll for CC and \ll we are interested only in values $\| \langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle \|_M$ for $\langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle$; but then, for each $a \in M$, we have $\| \kappa \|_M(a) \geq \| \bar{\kappa} \|_M(a)$ and $\| \lambda \|_M(a) \geq \| \bar{\lambda} \|_M(a)$. Similarly for a CD-closure set (operator). The following lemma shows that the theory of CD-closure sets reduces completely to the theory of CC-closure sets (cf. 14.10):

14.9. Lemma Let \ll, \ll^* be two operators of the type 1^n and let $[\langle \kappa, \delta \rangle \ll^* \langle \bar{\kappa}, \bar{\lambda} \rangle] \cong_M \cong_M [\langle \kappa, \text{neg}(\bar{\delta}) \rangle \ll \langle \bar{\kappa}, \text{neg}(\bar{\lambda}) \rangle]$ for each model M and any EC's $\kappa \ll \bar{\kappa}$ and ED's $\delta \ll \bar{\delta}$. Then \ll is a CC-closure operator iff \ll^* is a CD-closure operator.

Evident; note that $\delta \ll \bar{\delta}$ iff $\text{neg}(\delta) \ll \text{neg}(\bar{\delta})$.

14.10. Corollary. For each $\langle u, v, \bar{u}, \bar{v} \rangle \in \{0, \times, 1\}^4$, let $N(\langle u, v, u, v \rangle) = \langle u, \neg v, \bar{u}, \neg \bar{v} \rangle$ (where, of course, $\neg 1 = 0, \neg \times = \times, \neg 0 = 1$). Let $U \subseteq \{0, \times, 1\}^4$. Then U is a CC-closure set iff N^*U is a CD-closure set.

14.11. Remark and Definition. (1) Note that if \sim is an operator of the type 1^2 , it is not true that improvement operators for \sim, CC and \ll reduce to improvement operators for \sim, CD and $(\ll \ll)$! See Section 16.

(2) We shall find reasonable necessary and sufficient conditions for a set $U \subseteq \{0, \times, 1\}^4$ to be a CC-closure set. We shall deal with U as with a binary relation on $\{0, \times, 1\}^2$.

(3) \geq_{CC} is the following ordering of $\{0, \times, 1\}^2$. $\langle u, v \rangle \geq_{CC} \langle \bar{u}, \bar{v} \rangle$ iff $u \leq \bar{u}$ and $v \leq \bar{v}$.

(4) If $u = \langle u_1, u_2 \rangle$ and $v = \langle v_1, v_2 \rangle$ then $u \& v = \langle u_1 \& v_1, u_2 \& v_2 \rangle$.

14.12. Lemma. A set $U \subseteq \{0, \times, 1\}$ is a closure set iff the following holds for each $u, v, w \in \{0, \times, 1\}^2$:

- (I) uUu ;
- (II) uUv implies $u \geq_{CC} v$;
- (III) uUv and uUw implies uUw ;
- (III) uUv and uUw implies uUw ;
- (IV) uUw and $u \geq_{CC} v \geq_{CC} w$ implies uUv, vUw ;
- (V) uUv and uUw implies $uUv \& w$.

Obvious from the definition 13.9 and 14.9 (of a closure operator and a closure set).

14.13. Theorem. A set $U \subseteq \{0, \times, 1\}^4$ is a CC-closure set iff there is an equivalence E on $\{0, \times, 1\}^2$ such that


- (i) $u \equiv_E w$ and $u \geq_{CC} v \geq_{CC} w$ implies $u \equiv_E v \equiv_E w$;
- (ii) $u \equiv_E v$ implies $u \equiv_E u \& v \equiv_E w$;
- (iii) uUv iff $u \geq_{CC} v$ and $u \equiv_E v$.


Proof. (1) Consider U as a graph on $\{0, \times, 1\}^2$ and let $u \equiv_E v$ mean that u and v are in the same component of U . Suppose U to be a CC-closure set. We prove (iii). If uUv then $u \geq_{CC} v$ by (II) and evidently $u \equiv_E v$; to prove the converse suppose we have a chain u_0, \dots, u_n such that, for each i , either u_iUu_{i+1} or $u_{i+1}Uu_i$ (i.e. u_0 and u_n are in the same component). We prove $u_0U(u_0 \& u_i)$ for $i = 1, \dots, n$. This is obvious for $i = 1$ by (I). Let the statement hold for some i and consider $i + 1$. If $u_i \geq_{CC} u_{i+1}$ then u_iUu_{i+1} hence $u_iU(u_0 \& u_i \& u_{i+1})$ by (V), hence $u_iU(u_0 \& u_{i+1})$ and $u_0 \& u_iUu_0 \& u_{i+1}$ by (IV). Then $u_0U(u_0 \& u_{i+1})$ by (III) and $u_{i+1}U(u_0 \& u_{i+1})$ by (IV) (from $u_iU(u_0 \& u_{i+1})$). On the other hand, if $u_{i+1} \geq_{CC} u_i$ then $u_{i+1}Uu_iUu_0 \& u_i$; hence $u_{i+1}Uu_0 \& u_i$; furthermore, $u_{i+1} \geq_{CC} u_0 \& u_{i+1} \geq_{CC} u_0 \& u_i$, which implies $u_{i+1}U(u_0 \& u_{i+1})$ by (IV). Similarly, $u_0U(u_0 \& u_i)$ and $u_0 \geq_{CC} u_0 \& u_{i+1} \geq_{CC} u_0 \& u_i$ implies $u_0U(u_0 \& u_{i+1})$ by (IV). This proves (iii), since if $u = u_0, v = u_n$ and $u \leq_{CC} v$ then uUv ($v = u \& v$).

We prove (i). Let $u \equiv_E w$ and $u \geq_{CC} v \geq_{CC} w$. By (iii), uUw ; hence $uUvUw$ by (IV) and hence $u \equiv_E v \equiv_E w$. We prove (ii). Let $u \equiv_E v, u = u_0, v = u_n, u_0, \dots, u_n$ as above. Then it follows $u_0U(u_0 \& u_i)$ as above; we obtain $uU(u \& v)$ and $vU(v \& u)$. This completes the proof if the implication \rightarrow in the theorem.

(2) Suppose now that (i)–(iii) are satisfied; we prove that U is a closure set. We verify (I)–(V). (I) and (II) are evident from (iii). (III) follows from (iii) using transitivity of E and of \geq_{CC} . (IV) follows by (i) and (V) follows by (I). This completes the proof.

14.14. Examples. The previous theorem enables us to represent a closure set as an appropriate decomposition of $\{0, \times, 1\}^2$. The set $\{0, \times, 1\}^2$ is represented as a square matrix where the first row (column) corresponds to the value 1, the second to the value \times and the third to the value 0. Thick lines define subsets of $\{0, \times, 1\}^2$ that form the decomposition. The conditions that must be satisfied are (i) and (ii) of 14.13: (iii) defines U .





- Comments: (a) $u \equiv_E v$ iff $u = v$;
 (b) $u \equiv_E v$ iff $u_1 \& u_2 = v_1 \& v_2$ ($u = \langle u_1, u_2 \rangle, v = \langle v_1, v_2 \rangle$);
 (c) $u \equiv_E v$ iff ($u = \langle 1, 1 \rangle$ is equivalent to $v = \langle 1, 1 \rangle$);
 (d) $u \equiv_E v$ iff (whenever u or v are in $\{\langle 1, 1 \rangle, \langle \times, 1 \rangle, \langle 1, \times \rangle\}$ then $u = v$).
 (e) This equivalence does not determine a CC-closure set since (i) is violated:
 $\langle 1, \times \rangle \equiv_E \langle \times, 0 \rangle, \langle 1, \times \rangle \not\equiv_{CC} \langle 1, 0 \rangle \not\equiv_{CC} \langle \times, 0 \rangle$ but not $\langle 1, \times \rangle \equiv_E \langle 1, 0 \rangle$.
 (f) Here (ii) is violated: $\langle \times, 0 \rangle \equiv_E \langle 0, \times \rangle, \langle \times, 0 \rangle \& \langle 0, \times \rangle = \langle 0, 0 \rangle$ but not
 $\langle \times, 0 \rangle \equiv_E \langle 0, 0 \rangle$.

14.15. Remark. (1) We shall meet some of the above examples in Sections 15 and 16.

(2) We are interested in EC's; we are dealing with psEC's since they are closed under supremum (w.r.t. \ll). Having a CC-closure operator, two questions arise:

- (i) If $\langle \kappa, \lambda \rangle$ is a pair of EC's, is $Reg \ll_M(\kappa, \lambda) = \langle \bar{\kappa}, \bar{\lambda} \rangle$ a pair of EC's?
- (ii) Is $\bar{\kappa}, \bar{\lambda}$ a pair of incompressible conjunctions? Is $\bar{\kappa} \& \bar{\lambda}$ incompressible? The following theorems give some information.

14.16. Theorem. Let U be a CC-closure set defining a CC-closure operator \ll . The following are equivalent:

- (i) $\langle 1, 1, u, v \rangle \in U$ implies $\langle u, v \rangle = \langle 1, 1 \rangle$;
- (ii) For each M such that $(\exists a \in M) (\|\kappa \& \lambda\|_M(a) = 1$ (κ, λ are EC), $Reg \ll_M(\kappa, \lambda)$ is a pair of EC's.

Proof. Suppose (i). Let $\|\kappa, \lambda\| \ll (\kappa \& (X)F, \lambda)\|_M = 1$ and $\|\kappa \& \lambda\|_M(a) = 1$. We want to show that $X = \emptyset$ is impossible. Suppose $X \neq \emptyset$. Then $\|\kappa \& (\emptyset)F\|_M(a) \in \{0, \times\}$, hence $\langle \|\kappa\|_M(a), \|\lambda\|_M(a), \|\kappa \& (\emptyset)F\|_M(a), \|\lambda\|_M(a) \rangle \notin U$, a contradiction.

Suppose non (i). Then either $\langle 1, 1, 1, \times \rangle \in U$ or $\langle 1, 1, \times, 1 \rangle \in U$; suppose $\langle 1, 1, 1, \times \rangle \in U$. Let F be a functor not in $\kappa \& \lambda$ and let M be a model in which, for each object a , $\|\kappa\|_M(a) = \|\lambda\|_M(a) = 1$ and $\|F\|_M(a) = \times$. Then $\|\kappa, \lambda\| \ll (\kappa \& (\emptyset)F, \lambda)\|_M = 1$, hence $Reg \ll_M(\kappa, \lambda)$ is not a pair of EC's.

14.17. Theorem. (1) If \ll is a CC-closure operator and if $\langle \kappa, \lambda \rangle$ is a pair of psEC's then for each M , $Reg \ll_M(\kappa, \lambda)$ is a pair of M -incompressible psEC's.

(2) Let \ll be universally defined by a CC-closure set U . The following are equivalent:

(i) $\langle 1, 0, 0, 0 \rangle, \langle 0, 1, 0, 0 \rangle \in U$ (and hence all pairs containing at least one 0 are equivalent in the equivalence E of 14.13).

(ii) For each M and each pair $\langle \kappa, \lambda \rangle$ of psC's, if we put $Reg \ll_M \langle \kappa, \lambda \rangle = \bar{\kappa}, \bar{\lambda}$ then $\kappa \& \lambda$ is M -incompressible.

Proof. (1) follows from 14.7.

(2) Suppose (i). It suffices to verify the following: If $\|(\kappa, \lambda) \ll (\kappa \& (X)F, \lambda)\|_M = 1$ and $\|\kappa \& \lambda \& (X)F \cong \kappa \& \lambda \& (X_0)F\|_M = 1$ for and $X_0 \subseteq X$ then $\|(\kappa, \lambda) \ll (\kappa \& (X_0)F, \lambda)\|_M = 1$ (and similarly for $\kappa, \lambda \& (X)F$). Indeed, if for an object a the value of $\kappa \& \lambda \& (X)F$ is 1 then the value of $\kappa \& \lambda \& (X_0)F$ is also 1 and hence $\|(X_0)F\|_M(a) = \|(X)F\|_M(a) = 1$; hence the quadruple $\langle u, v, \bar{u}, \bar{v} \rangle$ of values of $\kappa, \lambda, \kappa \& (X)F, \lambda$ equals to the quadruple $\langle u, v, \hat{u}, \hat{v} \rangle$ of values of $\kappa, \lambda, \kappa \& (X_0)F, \lambda$ and so the latter one is in U . If the value of $\kappa \& \lambda \& (X)F$ is \times then either $\|(X)F\|_M(a) = \times$ and hence $\|(X)F\|_M(a) = \|(X_0)F\|_M(a) = \times$ or $\|(X)F\|_M(a) = 1$ and then $\|(X_0)F\|_M(a) = 1$ which follows from $\kappa \& \lambda \& (X)F \cong_M \kappa \& \lambda \& (X_0)F$. If $\|\kappa \& \lambda \& (X)F\|_M(a) = 0$ then $\|(X_0)F\|_M(a)$ can be different from $\|(X)F\|_M(a)$ and we still have $\kappa \& \lambda \& (X)F \cong_M \kappa \& \lambda \& (X_0)F$. But in the present case we have $(\bar{u} = 0$ or $\bar{v} = 0$ and $(\hat{u} = 0$ or $\hat{v} = 0)$, hence $\langle \bar{u}, \bar{v} \rangle \equiv_E \langle \hat{u}, \hat{v} \rangle$; furthermore, $\langle \bar{u}, \bar{v} \rangle \geq_{cc} \langle \hat{u}, \hat{v} \rangle$, hence $\langle \bar{u}, \bar{v} \rangle U \langle \hat{u}, \hat{v} \rangle$, which together with $\langle u, v \rangle U \langle \bar{u}, \bar{v} \rangle$ yields $\langle u, v, \hat{u}, \hat{v} \rangle \in U$.

Suppose not (i) and let e.g. $\langle 1, 0 \rangle \not\equiv_E \langle 0, 0 \rangle$. Let $M, \kappa, \lambda, X_0 \subset X$ be such that for each a , $\|\kappa\|_M(a) = \|(X)F\|_M(a) = 1$, $\|\lambda\|_M(a) = \|(X_0)F\|_M(a) = 0$. Then $\|(\kappa, \lambda) \ll (\kappa \& (X)F, \lambda)\|_M = 1$, $\|(\kappa, \lambda) \ll (\kappa \& (X_0)F, \lambda)\|_M = 0$, $\kappa \& \lambda(X)F \cong_M \cong_M \kappa \& \lambda \& (X_0)F$. Hence if $\langle \bar{\kappa}, \bar{\lambda} \rangle = Reg \ll_M \langle \kappa, \lambda \rangle$ then $\bar{\kappa} \& \bar{\lambda}$ is M -compressible.

14.18. Theorem. If both 14.16 (i) and (14.17 (i) holds and if $\kappa \& \lambda$ is an EC then for $Reg \ll_M \langle \kappa, \lambda \rangle = \langle \bar{\kappa}, \bar{\lambda} \rangle$ we have: $\bar{\kappa} \& \bar{\lambda}$ is an EC (in the following sense: whenever $(X)F$ occurs in $\bar{\kappa}$ and $(Y)F$ occurs in $\bar{\lambda}$ then $X = Y$; all coefficients are non-empty).

Proof. The fact that all coefficients are non-empty follows by 14.16 If $\|(\kappa, \lambda) \ll (\kappa \& (X)F, \lambda \& (Y)F)\|_M = 1$ observe that $(\kappa \& \lambda \& (X)F) \& (Y)F \cong_M \cong_M (\kappa \& \lambda \& (X)F) \& (X \cap Y)F$ so that, by the proof of 14.17, we have $\|(\kappa, \lambda) \ll (\kappa \& (X)F, \lambda \& (X \cap Y)F)\|_M = 1$. Similarly we obtain $\|(\kappa, \lambda) \ll (\kappa \& (X \cap Y)F, \lambda \& (X \cap Y)F)\|_M = 1$.

14.19. Definition. Let U be an CC-closure set defining a closure operator \ll . Define

$$\|Ant(\varphi_1, \varphi_2, \psi)\|_M = \|(\varphi_1, \varphi_2) \ll (\varphi_1 \& \psi, \varphi_2)\|_M,$$

$$\|Suc(\varphi_1, \varphi_2, \psi)\|_M = \|(\varphi_1, \varphi_2) \ll (\varphi_1 \& \psi)\|_M \text{ for each } M.$$

(Hence *Ant* and *Suc* are operators of the type 1^3 called the *antecedent operator* and the *succedent operator* corresponding to \ll).

14.20. Theorem. If \ll is a CC-closure operator and if κ, λ are psEC's then $Reg \ll_M(\kappa, \lambda) = \langle \bar{\kappa}, \bar{\lambda} \rangle$ where $\bar{\kappa}$ is the conjunction of all pseudoliterals $(X) F$ such that (X) is the smallest coefficient of all pseudoliterals (Z) such that $\|Ant(\kappa, \lambda, (Z) F)\|_M = 1$ and $\bar{\lambda}$ is the conjunction of all pseudoliterals $(X) F$ such that (X) is the smallest coefficient (Z) such that $\|Suc(\kappa, \lambda, (Z) F)\|_M = 1$.

Proof. Put $Reg \ll_M(\kappa, \lambda) = \langle \hat{\kappa}, \hat{\lambda} \rangle$ and let $\bar{\kappa}, \bar{\lambda}$ be defined as in the theorem. Evidently, $\|(\kappa, \lambda) \ll (\bar{\kappa}, \bar{\lambda})\|_M = 1$, hence $\bar{\kappa} \ll \hat{\kappa}$ and $\bar{\lambda} \ll \hat{\lambda}$. On the other hand, let $(X) F$ be a literal from $\hat{\kappa}$; then $\|Ant(\kappa, \lambda, (X) F)\|_M = 1$ by the definition of *Ant*. Suppose that there is an $X_0 \subset X$ such that $\|Ant(\kappa, \lambda, (X_0) F)\|_M = 1$. Then obviously $\|(\kappa, \lambda) \ll (\hat{\kappa}((X_0) F|(X) F), \hat{\lambda})\|_M = 1$ ($\hat{\kappa}(\dots)$ results from $\hat{\kappa}$ by replacing $(X) F$ by $(X_0) F$) and $\hat{\kappa}((X_0) F|(X) F) \supseteq_M \hat{\kappa}$, which contradicts to $\langle \hat{\kappa}, \hat{\lambda} \rangle = Reg \ll_M(\kappa, \lambda)$. Hence (X) is the smallest coefficient (Z) such that $\|Ant(\kappa, \lambda, (Z) F)\|_M = 1$ and $(X) F$ is in $\bar{\kappa}$. We have proved $\bar{\kappa} = \hat{\kappa}$ (the proof of $\bar{\lambda} = \hat{\lambda}$ is similar).

14.21. Remark. The last theorem shows that one can find $Reg \ll_M(\kappa, \lambda)$ quickly; one has to consider separate pseudoliterals and not all pairs $\langle \bar{\kappa}, \bar{\lambda} \rangle$ such that $\langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle$.

14.22. Theorem. Let both 14.16 (i) and 14.17 (i) hold. The following are equivalent:
 (i) $\langle u, v, \bar{u}, \bar{v} \rangle \in U$ implies $u \& v = \bar{u} \& \bar{v}$

$$\left(\text{hence } \begin{matrix} (\kappa, \lambda) \ll (\bar{\kappa}, \bar{\lambda}) \\ \kappa \& \lambda \cong \bar{\kappa} \& \bar{\lambda} \end{matrix} \text{ is sound} \right).$$

(ii) If $\kappa \& \lambda$ is M -incompressible and $\langle \bar{\kappa}, \bar{\lambda} \rangle = Reg \ll_M(\kappa, \lambda)$ then $\kappa \& \lambda \subseteq \bar{\kappa} \& \bar{\lambda}$ (subconjunction).

Proof. Suppose (i) We have $\kappa \& \lambda \cong_M \bar{\kappa} \& \bar{\lambda}$ and $\kappa \& \lambda \subseteq \bar{\kappa} \& \bar{\lambda}$. Let $\kappa_0 \& \lambda_0$ be the subconjunction of $\bar{\kappa} \& \bar{\lambda}$ having the same functors as $\kappa \& \lambda$. Then obviously $\kappa \& \lambda \cong_M \kappa_0 \& \lambda_0$ and $\kappa_0 \& \lambda_0$ is poorer than $\kappa \& \lambda$, so $\kappa_0 \& \lambda_0$ equals to $\kappa \& \lambda$, since $\kappa \& \lambda$ is incompressible.

Suppose not (i). Consider e.g. the case $\langle \times, 1, 0, 1 \rangle \in U$ (other cases similarly). Consider a $\{0, 1, 2, \times\}$ - model of the type 1^3 with two objects given by the following table. Put $\kappa = (1, 2) F \& (1) G, \bar{\kappa} = 1(1) F \& (1) G, \lambda = (1) H$.

F	G	H	$(X) F$	$(X_0) F$	κ	$\bar{\kappa}$	λ
2	\times	1	1	0	\times	0	1
1	1	1	1	1	1	1	1

Clearly, κ is M -incompressible, $\bar{\kappa}$ is strictly poorer than κ , and $\|(\kappa, \lambda) \ll (\bar{\kappa}, \bar{\lambda})\|_M = 1$.

In the present section we are going to study secured (a fortiori, qualitative) operators of the type 1^2 that for $\{0, 1\}$ -models mean: coincidence of the two properties predominates over difference. It should be clear that this meaning can be made precise in many ways, both statistical and non-statistical.

15.1. Definition. (1) If M is a $\{0, 1\}$ -model then a_M, b_M, c_M, d_M denote the cardinality of the set of objects having the card $\langle 1, 1 \rangle, \langle 1, 0 \rangle, \langle 0, 1 \rangle, \langle 0, 0 \rangle$ respectively. (As usual, the *card* of an object $a \in M$ in $M = \langle M, f_1, f_2 \rangle$ is $\langle f_1(a), f_2(a) \rangle$). We put $q_M = \langle a_M, b_M, c_M, d_M \rangle$.

(2) In the sequel saying “quadruple” we mean a quadruple of natural numbers whose sum is positive (so that for such a quadruple $\langle a, b, c, d \rangle$ there is an M such that $a = a_M, \dots, d = d_M$).

15.2. Definition. (1) A quadruple $q_1 = \langle a_1, b_1, c_1, d_1 \rangle$ is *a-better* than $q_2 = \langle a_2, b_2, c_2, d_2 \rangle$ iff $a_1 + b_1 + c_1 + d_1 = a_2 + b_2 + c_2 + d_2$ and $a_1 \geq a_2, b_1 \leq b_2, c_1 \leq c_2, d_1 \geq d_2$.

(2) A secured operator \sim of the type 1^2 is *associational* if the following holds for any $\{0, 1\}$ -models M_1, M_2 of the type 1^2 : If $Asf_{\sim}(M_1) = 1$ and if q_{M_2} is a-better than q_{M_1} then $Asf_{\sim}(M_2) = 1$.

15.3. Examples of associational operators: (1) the “relatively more” operator of [3] 8.9: $Asf_{\sim}(M) = 1$ iff $a_M d_M > b_M c_M$.

(2) The Fisher operator (see [5] p. 430–431): $Asf_{\sim\sigma}(M) = 1$ iff $a_M d_M > b_M c_M$ and $\Delta(a_M, b_M, c_M, d_M) \leq \sigma$ where Δ is the Fisher statistic.

(3) The χ^2 -operator: $Asf_{\chi^2}(M) = 1$ iff $a_M d_M > b_M c_M$ and $[(a_M d_M - b_M c_M)^2] : [(a_M + b_M)(a_M + c_M)(b_M + d_M)(c_M + d_M)] \geq \chi^2$ (see e.g. [7]).

15.4. Definition. (1) A $\{0, 1\}$ -model M_2 is *a-better* than a $\{0, 1\}$ -model M_1 if q_{M_2} is a-better than q_{M_1} .

(2) A $\{0, \times, 1\}$ -model M_2 is *a-better* than a $\{0, \times, 1\}$ -model M_1 if for each completion N_2 of M_2 there is a completion N_1 of M_1 such that N_2 is a-better than N_1 .

(3) M_1 is *a-equivalent* to M_2 if M_1 is a-better than M_2 and vice versa.

15.5. Remark. Evidently, both definitions of “a-better” coincide for $\{0, 1\}$ -models. The “a-better” relation is reflexive and transitive, hence a quasiordering; a-equivalence is the canonical equivalence given by the “a-better” quasiordering. (cf. [2] 3.6).

15.6. Lemma (1) Let M_1, M_2 be qualitative models (i.e. $\{0, \times, 1\}$ -models) of the type 1^2 and let \sim be an associational operator. If M_2 is a-better than M_1 and if $Asf_{\sim}(M_1) = 1$ then $Asf_{\sim}(M_2) = 1$.

(2) If M_2 is not a-better than M_1 then one can define an associational operator \sim such that $Asf_{\sim}(M_1) = 1$ but $Asf_{\sim}(M_2) \neq 1$.

Proof. (1) Note that the assertion is obvious if M_1, M_2 have complete information (= are $\{0, 1\}$ -models). If N_2 is a completion of M_2 then there is a completion N_1 of M_1 such that N_2 is a-better than N_1 . Since $Asf_{\sim}(M_1) = 1$ and \sim is secured we have $Asf_{\sim}(N_1) = 1$; since \sim is associational and N_1 is a $\{0, 1\}$ -model we have $Asf(N_2) = 1$. Since N_2 was an arbitrary completion of M_2 we have $Asf_{\sim}(M_2) = 1$.

(2) First suppose that M_1, M_2 have complete information. Then put for each $\{0, 1\}$ -model M $Asf_{\sim}(M) = 1$ iff q_M is a-better than q_{M_1} . By the transitivity of "a-better", this is an associational operator; $Asf_{\sim}(M_1) = 1$ and $Asf_{\sim}(M_2) = 0$. In the general case there exists a completion N_2 of M_2 such that for no completion N_1 of M_1 , N_2 is a-better than N_1 . Put for each $\{0, 1\}$ -model M , $Asf_{\sim}(M) = 1$ iff there is a completion N_1 of M_1 such that M is a-better than N_1 . Then $Asf_{\sim}(M_1) = 1$ by securedness but $Asf_{\sim}(M_2) \neq 1$ since $Asf_{\sim}(N_2) = 0$.

15.7. Definition. If $M = \langle M, F_1, f_2 \rangle$ is a qualitative model, if $A \subseteq M$ and if $u = \langle u_1, u_2 \rangle \in \{0, \times, 1\}^2$ then $M(A : u)$ is the model $\langle M, g_1, g_2 \rangle$ where $g_i(a) = f_i(a)$ for $a \notin A$ and $g_i(a) = u_i$ for $a \in A$ (cards of elements of A are changed to be u). In particular, if $a \in M$ then $M(a : u)$ means $M(\{a\} : u)$ and if $v \in \{0, \times, 1\}^2$ then $M(v : u)$ means $M(A : u)$ for $A = \{a \in M; \text{the card of } a \text{ is } v\}$.

15.8. Remark. (1) If $a_1 \neq a_2$ then $M(a_1 : u)(a_2 : v) = M(a_2 : v)(a_1 : u)$.

(2) If $A = \{a_1, \dots, a_n\}$ then $M(A : u) = M(a_1 : u) \dots (a_n : u)$.

15.9. Definition. Let $u, v \in \{0, \times, 1\}^2$. v is a-better than u ($v \geq_{ab} u$) if for each qualitative model M of the type 1^2 and each $a \in M$ we have the following: If the card of a is u then $M(a : v)$ is a-better than M .

15.10. Remark (1) Evidently \geq_{ab} is a quasiordering.

(2) An alternative definition reads as follows: $v \geq_{ab} u$ iff for each qualitative M of the type 1^2 and each $a \in M$ we have: if the card of a is v the M is a-better than $M(a : u)$.

15.11. Theorem. The quasiordering \geq_{ab} is completely described by the following conditions:

(a) $\langle 1, 0 \rangle \equiv_{ab} \langle 1, \times \rangle \equiv_{ab} \langle \times, 0 \rangle$; $\langle 0, 1 \rangle \equiv_{ab} \langle \times, 1 \rangle \equiv_{ab} \langle 0, \times \rangle$.

(b) $\langle \times, \times \rangle <_{ab} \langle 0, 1 \rangle$, $\langle \times \times \rangle <_{ab} \langle 1, 0 \rangle$, $\langle 1, 0 \rangle <_{ab} \langle 1, 1 \rangle$,
 $\langle 1, 0 \rangle <_{ab} \langle 0, 0 \rangle$, $\langle 0, 1 \rangle <_{ab} \langle 1, 1 \rangle$, $\langle 0, 1 \rangle <_{ab} \langle 0, 0 \rangle$.

(c) $\{\langle 1, 1 \rangle, \langle 0, 0 \rangle\}$ and $\{\langle 1, 0 \rangle, \langle 0, 1 \rangle\}$ are incomparable pairs.

We visualize \geq_{ab} in the following diagram where a dashed line means equivalence and a dashed line together with a thick line means that the transition from the side of the dashed line to the side of the thick line makes the pair (strictly) a-better (drives understand).



Proof. $K(a)$ denotes the card of a in an arbitrary fixed model M of the type 1^2 .


(a) Let $K(a) = \langle 1, \times \rangle$ and consider $M' = M(a : \langle 1, 0 \rangle)$. Each completion of M' is a completion of M , hence M' is a-better than M . Conversely, if N is a completion of M then the card of a in N is either $\langle 1, 0 \rangle$ or $\langle 0, 1 \rangle$. In the former case N is a completion of M' ; in the latter case $N' = N(a : \langle 1, 0 \rangle)$ is a completion of M' and N is a-better than N' . Remaining cases are treated similarly.

(c) Let $K(a) = \langle 1, 1 \rangle$ in a $\{0, 1\}$ -model M , put $N = M(a : \langle 0, 0 \rangle)$. Then $q_N = a_M - 1, b_M, c_M, d_M + 1$ so that neither q_M is a-better than q_N nor q_N is a-better than q_M . This shows that $\langle 1, 1 \rangle, \langle 0, 0 \rangle$ are incomparable.

(b) Evidently, $\langle 1, 1 \rangle \geq_{ab} \langle 1, 0 \rangle$. To prove that $\langle 1, 0 \rangle \geq_{ab} \langle 1, 1 \rangle$ does not hold, it suffices to take a $\{0, 1\}$ -model M of the type 1^2 with at least one card $\langle 1, 0 \rangle = K(a)$ and observe that for $N = M(a : \langle 1, 1 \rangle)$ q_M is not a-better than q_N .

If $K(a) = \langle \times, \times \rangle$ then for any u each completion of $M(a : u)$ is a completion of M , hence $M(a : u)$ is a-better than M . To prove that e.g. $\langle \times, \times \rangle \geq_{ab} \langle 1, 1 \rangle$ does not hold suppose that $K(a) = \langle \times, \times \rangle$ in a model M_0 and suppose that all other cards of M_0 are in $\{0, 1\}^2$. Put $M = M_0(a : \langle 1, 1 \rangle)$ and $N = M(a : \langle 0, 0 \rangle)$. We showed in (c) that neither M is a-better than N nor N is a-better than M . This shows that $\langle \times, \times \rangle$ is not a-better than $\langle 1, 1 \rangle$. All remaining cases are treated similarly.

15.12. Theorem. There are four \subseteq -maximal CC-closure sets such that the corresponding closure operator \ll has the following property: For each associational operator \sim, \ll is an improvement operator for \sim, CC and \ll . They are defined by the following tables (i)–(iv):



All of them satisfy 14.16 (i); but only the first one satisfies 14.17 (i). (Hence we prefer the first operator, cf. 14.18.)

Proof. Consider the table in 15.9 and recall that we are interested only in quadruples $\langle u, v \rangle$ where $u \geq_{cc} v$. Since U has to define an improvement operator, $\langle u, v \rangle \in U$ must imply $u \geq_{ab} v$ (cf. 15.6 (2)). Consequently, the following quadruples must not be in U : $\langle 1, 1, 1, \times \rangle, \langle 1, 1, \times, 1 \rangle, \langle 1, \times, \times, \times \rangle, \langle \times, 1, \times, \times \rangle$ (see table (v)).

(v)

The tables (i)–(iv) show four possible maximal equivalence relations in which none of the above quadruples is a pair of equivalent pairs. Cf. 14.13.

15.13. Definition. \ll^a means the closure operator defined by the table (i) of 15.2.

15.14. Theorem. Let $\|Ant^a(\kappa, \lambda, \kappa')\|_M = \|(\kappa, \lambda) \ll^a (\kappa \& \kappa', \lambda)\|_M$ and analogously for Suc^a (cf. 14.19). Then (1) Ant^a is universally defined by the set

$$U_A^a = \{ \langle u, v, \bar{u} \rangle \in \{0, \times, 1\}^3; [(\langle u, v \rangle = \langle 1, 1 \rangle \vee \langle u, v \rangle = \langle 1, \times \rangle) \rightarrow \bar{u} = 1] \& \\ \& [\langle u, v \rangle = \langle \times, 1 \rangle \rightarrow \bar{u} \in \{1, \times\}] \} .$$

(2) Suc^a is universally defined by the set

$$U_S^a = \{ \langle u, v, \bar{v} \rangle \in \{0, \times, 1\}^3; [(\langle u, v \rangle = \langle 1, 1 \rangle \vee \langle u, v \rangle = \langle \times, 1 \rangle) \rightarrow \bar{v} = 1] \& \\ \& [\langle u, v \rangle = \langle 1, \times \rangle \rightarrow \bar{v} \in \{1, \times\}] \} .$$

Proof. In the following table we list all possible tuples of values of $\kappa, \lambda, \kappa \& \kappa'$ such that the quadruple of values of $\kappa, \lambda, \kappa \& \kappa', \lambda$ is in U ; the fourth line contains the corresponding possible values of κ' . “arb” means arbitrary.

κ	1	1	\times	\times	0	arb
λ	1	\times	1	\times	arb	0
$\kappa \& \kappa'$	1	1	\times	$\leq \times$	0	$\leq \ \kappa\ $
κ'	1	1	1, \times	arb	arb	arb

15.15. Theorem. Let κ, λ be psEC's and let M be a model. (1) (a) There is an X such that $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 1$ iff there is no object $a \in M$ such that $[(\|\kappa\|_M(a), \|\lambda\|_M(a)) \text{ is } \langle 1, 1 \rangle \text{ or } \langle 1, \times \rangle] \text{ and } \|F\|_M(a) = \times$. (b) Suppose the last condition holds; call a $u \in \{0, \times, 1\}^2$ critical if $u \in \{ \langle 1, 1 \rangle, \langle 1, \times \rangle, \langle \times, 1 \rangle \}$. The least X such that $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 1$ is $X = \{ \|F\|_M(a); \langle \|\kappa\|_M(a), \|\lambda\|_M(a) \rangle \text{ critical and } \|F\|_M(a) \in V_{reg} \}$.

- (2) (a) There is an X such that $\|Suc^a(\kappa, \lambda, (X)F)\|_M = 1$ iff there is no object $a \in M$ such that $[\langle \|\kappa\|_M(a), \|\lambda\|_M(a) \rangle \text{ is } \langle 1, 1 \rangle \text{ or } \langle \times, 1 \rangle \text{ and } \|F\|_M(a) = \times]$.
 (b) If the last condition holds then the least X such that $\|Suc^a(\kappa, \lambda, (X)F)\|_M = 1$ is the X defined in (1) (b) above.

Proof. Use the table for Ant^a in 15.14. For each $a \in M$ put $u_a = \|\kappa\|_M(a)$, $v_a = \|\lambda\|_M(a)$, $f_a = \|F\|_M(a)$. If there is an object with $\langle u_a, v_a \rangle = \langle 1, 1 \rangle$ or $\langle 1, \times \rangle$ and $f_a = \times$ then, for any X , $\langle u_a, v_a, \|(X)F\|_M(a) \rangle \notin U_A^a$ and hence $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 0$. If there is no such a take the coefficient defined in (1) (b); we show $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 1$. Let $a \in M$. If $\langle u_a, v_a \rangle$ is not critical nothing is to prove. If $\langle u_a, v_a \rangle$ is $\langle 1, 1 \rangle$ or $\langle 1, \times \rangle$ then $f_a \in V_{reg}$, hence $f_a \in X$ and $\|(X)F\|_M(a) = 1$ hence $\langle u_a, v_a, \|(X)F\|_M(a) \rangle \in U_A^a$. If $\langle u_a, v_a \rangle$ is $\langle \times, 1 \rangle$ then either $f_a \in V_{reg}$ and $\|(X)F\|_M(a) = 1$ (as above) or $f_a = \times$, hence $\|(X)F\|_M(a) = \times$ and $\langle u_a, v_a, \|(X)F\|_M(a) \rangle \in U_A^a$. Hence $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 1$. It remains to show that X is minimal. Let $j \in X - Y$ and let a be such that $\langle u_a, v_a \rangle$ is critical and $f_a = j$. Then $\|(Y)F\|_M(a) = 0$ and $\langle u_a, v_a, 0 \rangle \notin U_A^a$; hence $\|Ant^a(\kappa, \lambda, (Y)F)\|_M = 0$.

15.16. Remark. The last theorem yields an effective algorithm for finding $Reg \ll_M^a(\kappa, \lambda)$ (cf. 14.20). *Caution:* it may happen that the only X such that $\|Ant^a(\kappa, \lambda, (X)F)\|_M = 1$ is $X = V_{reg}$; then $(X)F$ is not a pseudoliteral and is not put into $\bar{\kappa}$ (where $Reg \ll_M^a(\kappa, \lambda) = \langle \bar{\kappa}, \bar{\lambda} \rangle$).

15.17. Discussion. We discuss the relation of the above results to GUHA-problems and their solutions.

(i) Recall the general theory of improvement operators (13.9–13.13). Here we have an arbitrary associational operator \sim , the set CC of pairs of psEC's and the ordering \ll . Call \sim a *strict* associational operator if, for each M , κ, λ , $\|\kappa \sim \lambda\|_M = 1$ implies $(\exists a \in M) (\|\kappa \& \lambda\|_M(a) = 1)$. (E.g. the Fisher's operator etc.) If \sim is strict then whenever $\langle \kappa, \lambda \rangle \in CC$ and $\|\kappa \sim \lambda\|_M = 1$ then $Reg \ll_M^a(\kappa, \lambda)$ is a pair of EC's (not only psEC's); denoting it by $\langle \hat{\kappa}, \hat{\lambda} \rangle$ we have $\|\hat{\kappa} \sim \hat{\lambda}\|_M = 1$. Hence suppose \sim to be strict. We have the problem $P = \langle CC, \{1\}, IC \rangle$ where IC is the following rule:

$$\frac{(\kappa \sim \lambda) \& (\kappa, \lambda) \ll (\hat{\kappa}, \hat{\lambda})}{\bar{\kappa} \sim \bar{\lambda}} (\langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle \ll \langle \hat{\kappa}, \hat{\lambda} \rangle).$$

Recall that $\kappa \sim \lambda$ is M -prime if $\langle \kappa, \lambda \rangle$ is the \ll -smallest pair $\langle \kappa', \lambda' \rangle$ such that $\|(\kappa' \sim \lambda') \& (K', \lambda') \ll (\kappa, \lambda)\|_M = 1$.

We know that $X_M = \{(\kappa \sim \lambda) \& (\kappa, \lambda) \ll (\hat{\kappa}, \hat{\lambda}); (\kappa \sim \lambda) \text{ is } M\text{-prime and } \langle \hat{\kappa}, \hat{\lambda} \rangle = Reg \ll_M^a(\kappa, \lambda)\}$ is a solution of P in M . $Reg \ll_M^a(\kappa, \lambda)$ is found using 15.15 and 14.20.

(ii) Note the following fact concerning the present solution: If we want X_M to determine a solution of a hierarchical problem $\langle P, H \rangle$ then, by 13.13, we must

request $R \subseteq R_H$ where R is the ordering described in 13.13. This means that the formula $\kappa \sim \lambda$ should be tested earlier than any $\bar{\kappa} \sim \bar{\lambda}$ with $\langle \kappa, \lambda \rangle \ll \langle \bar{\kappa}, \bar{\lambda} \rangle$. In particular, if κ, λ and $\bar{\kappa}, \bar{\lambda}$ contain the same functors but the coefficients in κ, λ are supersets of the corresponding coefficients in $\bar{\kappa}, \bar{\lambda}$ then $\kappa \sim \lambda$ is to be tested earlier. This is not in accordance with the natural syntactic "simpler-than" relation \triangleleft between EC's (cf. 12.8 (4)). We can ignore this fact when V_{reg} is small (e.g. has four elements); in particular, if $V_{reg} = \{0, 1\}$ then everything is all right and \ll reduces to \subseteq . On the other hand, if V_{reg} is big (e.g. has twenty elements) then one is forced to make some restrictions on the cardinality of coefficients. We are led to the following

15.18. Definition and Theorem. Let $1 \leq k \leq \text{card}(V_{reg})$ and put $F_k = \{\kappa \sim \lambda; \langle \kappa, \lambda \rangle \in CC \text{ and all coefficients in } \kappa \text{ and } \lambda \text{ have cardinality at most } k\}$, $P_k = F_k; \{1, IC\}$. Define $\kappa \sim \lambda$ to be $M - F_k$ -prime if $(\kappa \sim \lambda) \in F_k$, $\|\kappa \sim \lambda\|_M = 1$ and there is no $\langle \kappa_0, \lambda_0 \rangle \ll \langle \kappa, \lambda \rangle$, $\langle \kappa_0, \lambda_0 \rangle \neq \langle \kappa, \lambda \rangle$ such that $(\kappa_0 \sim \lambda_0) \in F_k$, $\|\kappa_0 \sim \lambda_0\|_M = 1$ and $\|(\kappa, \lambda_0) \ll (\kappa, \lambda)\|_M = 1$. Let $Reg \ll_{M,k}^a (\kappa, \lambda) = \langle \bar{\kappa}, \bar{\lambda} \rangle$ result from $Reg \ll_M^a (\kappa, \lambda) = \langle \bar{\kappa}, \bar{\lambda} \rangle$ by omitting in $\bar{\kappa}$ and $\bar{\lambda}$ all literals with coefficients of cardinality bigger than k . Put $X_M^k = \{(\kappa \sim \lambda) \& (\kappa, \lambda) \ll \langle \bar{\kappa}, \bar{\lambda} \rangle; (\kappa \sim \lambda) M - F_k\text{-prime and } \langle \bar{\kappa}, \bar{\lambda} \rangle = Reg \ll_{M,k}^a (\kappa, \lambda)\}$. Then X_M^k is a \subseteq -minimal solution of P_k in M .

The proof is left to the reader as an exercise.

To close the present section, we discuss the possibility of applying 14.22, using a weaker improvement operator and restricting one's interest to formulas $\kappa \sim \lambda$ such that $\|\kappa \sim \lambda\|_M = 1$ and $\kappa \& \lambda$ is M -incompressible.

15.19. Theorem and Definition. There is a uniquely determined CC-closure set satisfying 14.16 (i), 14.17 (i) and 14.22 (i) and such that the corresponding operator is an improvement operator for each associational operator \sim , for CC and \ll . It is given by the following table:

This operator will be denoted by \ll^b .

(2) Introduce Ant^b and Suc^b with the usual meaning; then Ant^b is universally defined by

$$U_a^b = \{\langle u, v, \bar{u} \rangle; [(u = 1 \& v \in \{1, \times\}) \rightarrow \bar{u} = 1] \& [(u = \times \& v \in \{1, \times\}) \rightarrow \bar{u} \in \{1, \times\}]\}.$$

Suc^b is universally defined by

$$U_s^b = \{\langle u, v, \bar{v} \rangle; [(v = 1 \& u \in \{1, \times\}) \rightarrow \bar{v} = 1] \& [(v = \times \& u = \{1, \times\}) \rightarrow \bar{v} \in \{1, \times\}]\}.$$

(3) Theorem 15.15 holds for the present operator \ll^b with the following change: critical pairs are $\langle 1, 1 \rangle$, $\langle 1, \times \rangle$, $\langle \times, 1 \rangle$, and $\langle \times, \times \rangle$.

15.20. Discussion. Let \sim be a strict associational operator. Put

$$IC^* = \left\{ \frac{(\kappa \sim \lambda) \& Inc(\kappa \& \lambda) \& (\kappa, \lambda) \ll^b (\hat{\kappa}, \hat{\lambda})}{(\bar{\kappa} \sim \bar{\lambda}) \& Inc(\bar{\kappa} \& \bar{\lambda})}; \langle \kappa, \lambda \rangle \subseteq \langle \bar{\kappa}, \bar{\lambda} \rangle \subseteq \langle \hat{\kappa}, \hat{\lambda} \rangle \right\},$$

$$F^* = \{(\kappa \sim \lambda) \& Inc(\kappa \& \lambda); \langle \kappa, \lambda \rangle \in CC\}$$

($Inc(\kappa)$ is a formula such that $\|Inc(\kappa)\|_M = 1$ iff κ is M -incompressible, $P^* = \langle F^*, \{1\}, IC^* \rangle$,

$$X_M^* = \{(\kappa \sim \lambda) \& Inc(\kappa \& \lambda) \& (\kappa, \lambda) \ll^b (\hat{\kappa}, \hat{\lambda}); (\kappa \sim \lambda)$$

$$M\text{-prime, } \kappa \& \lambda \text{ } M\text{-incompressible, } \langle \hat{\kappa}, \hat{\lambda} \rangle = Reg \ll_M^b(\kappa, \lambda)\}.$$

Then, by 15.19 and 14.22, IC^* is a sound rule (see also 12.12 (1) and 14.17) and X_M^* is a solution of P^* in M . Put $(\kappa \sim \lambda) \leq_Q (\bar{\kappa} \sim \bar{\lambda})$ iff $\langle \kappa, \lambda \rangle \subseteq \langle \bar{\kappa}, \bar{\lambda} \rangle$; then X_M^* determines a solution of a hierachical problem $\langle P^*, H \rangle$ whenever $Q \subseteq R_H$. (Hence one can go through CC e.g. in the order \leftarrow and obtain a hierarchical solution.) Recall that 15.19 (3) gives an effective algorithm for finding $Reg \ll_M^b(\kappa, \lambda)$. *Caution:* The intuitive adequacy of the restriction to relevance declared by F^* (in particular, to M -incompressible conjunctions) remains to be investigated.

16. IMPLICATION OPERATORS

We shall study some particular associational operators called implicational operators; they have some properties of the operator of logical implication (inclusion).

- 16.1. Definition.** (1) A quadruple $q_2 = \langle a_2, b_2, c_2, d_2 \rangle$ is *i-better* than $q_1 = \langle a_1, b_1, c_1, d_1 \rangle$ if $a_1 + b_1 + c_1 + d_1 = a_2 + b_2 + c_2 + d_2$, $a_2 \geq a_1$, $b_2 \leq b_1$.
 (2) $q_2 = \langle a_2, b_2, c_2, d_2 \rangle$ is *si-better* than $q_1 = \langle a_1, b_1, c_1, d_1 \rangle$ iff $a_1 + b_1 + c_1 + d_1 = a_2 + b_2 + c_2 + d_2$ and $a_2 b_1 \geq a_1 b_2$.

- 16.2. Definition.** (1) A secured operator \sim of the type 1^2 is *implicational* if the following holds for any $\{0, 1\}$ -models M_1, M_2 of the type 1^2 : whenever $Asf_-(M_1) = 1$ and q_{M_2} is i-better than q_{M_1} then $Asf_-(M_2) = 1$.
 (2) \sim is *strictly implicational* if the following holds for any $\{0, 1\}$ -models M_1, M_2 of the type 1^2 : whenever $Asf_-(M_1) = 1$ and q_{M_2} is si-better than q_{M_1} then $Asf_-(M_2) = 1$ and $a_{M_1} > 0$.

- 16.3. Lemma.** (1) If q_2 is a-better than q_1 then q_2 is i-better than q_1 .
 (2) If q_2 is i-better than q_1 then q_2 is si-better than q_1 .

- (3) If $a_1 + b_1 \neq 0 \neq a_2 + b_2$ then q_2 is si-better than q_1 iff $a_2 : (a_2 + b_2) \geq \geq a_1 : (a_1 + b_1)$.
- (4) Each implicational operator is associational.
- (5) Each strictly implicational operator is implicational.

16.4. Remark and Definition. (1) If \sim is implicational M_1, M_2 have the same cardinality and $a_{M_1} = a_{M_2}, b_{M_1} = b_{M_2}$ then $Asf_{\sim}(M_1) = Asf_{\sim}(M_2)$.

(2) Each of the operators defined in [3] 9.18, Examples (1)–(3) is strictly implicational. (Implication, good almost-implication, probable almost-implication.)

(3) One introduces the notion “ M_2 is i-better (si-better) than M_1 ” for qualitative models in the same way as the corresponding a-notion in 15.4. Finally we define in an analogy to 15.9: v is i-better than u ($u, v \in \{0, \times, 1\}^2; v \geq_{ib} u$) if for each qualitative model M of the type 1^2 we have the following: If the card of a is u then $M(a : v)$ is i-better than M (for each $a \in M$). Similarly for \geq_{sib} .

16.5. Theorem. The quasiordering \geq_{ib} is completely described by the following conditions:

- (a) $\langle 1, 0 \rangle \equiv_{ib} \langle 1, \times \rangle \equiv_{ib} \langle \times, 0 \rangle; \langle 0, 0 \rangle \equiv_{ib} \langle 0, \times \rangle \equiv_{ib} \langle 0, 1 \rangle \equiv_{ib} \langle \times, 1 \rangle;$
- (b) $\langle 1, 0 \rangle <_{ib} \langle 0, 0 \rangle <_{ib} \langle 1, 1 \rangle$



Proof. Note that $v \geq_{ab} u$ implies $v \geq_{ib} u$. Hence it suffices to show the following:

- (1) $\langle \times, \times \rangle \geq_{ib} \langle 1, \times \rangle$ (which yields $\langle \times, \times \rangle \geq_{ib} \langle 1, 0 \rangle, \langle 0, \times \rangle$ by transitivity),
- (2) $\langle 0, \times \rangle \geq_{ib} \langle 0, 0 \rangle,$
- (3) $\langle \times, \times \rangle \not\geq_{ib} \langle 0, \times \rangle$ (yields $\langle \times, \times \rangle \not\geq_{ib} \langle 0, 0 \rangle, \langle 1, \times \rangle, \langle 1, 0 \rangle,$
- (4) $\langle 1, \times \rangle \not\geq_{ib} \langle 1, 1 \rangle$ and $\langle \times, 1 \rangle \not\geq_{ib} \langle 1, 1 \rangle.$

(1) Let $a \in M, K_M(a) = \langle 1, \times \rangle;$ put $M_1 = M(a : \langle \times, \times \rangle)$ and let N_1 be a completion of M_1 . If $K_{N_1}(a) = \langle 1, v \rangle$ then N_1 is a completion of M ; if $K_{N_1}(a) = \langle 0, v \rangle$ then put $N = N_1(a : \langle 1, 0 \rangle); a_N = a_{N_1}$ and $b_N = b_{N_1} + 1,$ hence N_1 is i-better than N and hence M_1 is i-better than M .

(3) Let $K_M(a) = \langle 0, \times \rangle,$ put $M_1 = M(a : \langle \times, \times \rangle)$ and suppose that for any $b \neq a, b \in M,$ the card $K_M(b)$ is in $\{0, 1\}^2$. Put $N_1 = M_1(a : \langle 1, 0 \rangle);$ we show that N_1 is not i-better than any completion N of M . We have two possibilities: $N = M(a : \langle 0, 0 \rangle)$ or $N = M(a : \langle 0, 1 \rangle).$ In both cases we have: $a_{N_1} = a_N, b_{N_1} = b_N + 1,$ hence N is i-better than N_1 and N_1 is not i-better than N .

- (2) Let $K_M(a) = \langle 0, 0 \rangle$, put $M_1 = M(a : \langle 0, \times \rangle)$. Let N_1 be a completion of M_1 . If $K_{N_1}(a) = \langle 0, 0 \rangle$ then N_1 is a completion of M ; if $K_{N_1}(a) = \langle 0, 1 \rangle$ then put $N = N_1(a : \langle 0, 0 \rangle)$; it is a completion of M and M, N are i-equivalent.
- (4) Let $K_M(a) = \langle 1, 1 \rangle$, put $M_1 = M(a : \langle 1, \times \rangle)$, suppose M to be a $\{0, 1\}$ -model. Put $N_1 = M_1(a : \langle 1, 0 \rangle)$; then N_1 is a completion of M_1 and N_1 is not i-better than M (M is its only own completion).

16.6. Theorem and Definition. There are two \subseteq -maximal CC-closure sets such that the corresponding closure operator \ll has the following property: For each implicational operator $\rightarrow^* \ll$ is an improvement operator for \rightarrow^* , CC and \ll . They are defined by the following tables (i), (ii):

(i)

(ii)

Both (i) and (ii) satisfy 14.16 (i); only the first one satisfies 14.17 (i). So we prefer (i). The operator defined by (i) will be denoted by \ll^c .

The proof is completely analogous to the proof of the corresponding theorem in Section 15 (Theorem 15.12) and so are the proofs of the following theorems 16.7–16.8. (*Erratum:* In Table (i) the line between $\langle 1, 1 \rangle$ and $\langle \times, 1 \rangle$ should be thick.)

16.7. Theorem. Let $\|Ant^c(\kappa, \lambda, \kappa')\|_M = \|(\kappa, \lambda) \ll^c(\kappa \& \kappa', \lambda)\|_M$ and analogously for Suc^c . Then (1) Ant^c is universally defined by

$$U_A^c = \{u, v, \bar{u}\} \in \{0, \times, 1\}^3; [\langle u, v \rangle = \langle 1, 1 \rangle \rightarrow \bar{v} = 1] \& [\langle u, v \rangle = \langle \times, 1 \rangle \rightarrow \bar{v} \in \{1, \times\}].$$

(2) Suc^c is universally defined by

$$U_S^c = \{\langle u, v, \bar{v} \rangle \in \{0, \times, 1\}^3; [(\langle u, v \rangle = \langle 1, 1 \rangle \vee \langle u, v \rangle = \langle \times, 1 \rangle) \rightarrow \bar{v} = 1]\}.$$

16.8. Theorem. Let κ, λ be psEC's and let M be a model. (1) (a) There is an X such that $\|Ant^c(\kappa, \lambda, (X)F)\|_M = 1$ iff there is no object $a \in M$ such that $\langle \|\kappa\|_M(a), \|\lambda\|_M(a) \rangle$ is $\langle 1, 1 \rangle$ and $\|F\|_M(a) = \times$.

(b) Suppose the last condition holds; call u critical if $u \in \{\langle 1, 1 \rangle, \langle 1, \times \rangle\}$. Then the least X such that $\|Ant^c(\kappa, \lambda, (X)F)\|_M = 1$ is $X = \{\|F\|_M(a); \langle \|\kappa\|_M(a), \|\lambda\|_M(a) \rangle$ critical and $\|F\|_M(a) \in V_{reg}\}$.

(2) (a) There is an X such that $\|Suc^c(\kappa, \lambda, (X)F)\|_M = 1$ iff there is no object $a \in M$ such that $\langle \|\kappa\|_M(a), \|\lambda\|_M(a) \rangle \in \{\langle 1, 1 \rangle, \langle \times, 1 \rangle\}$ and $\|F\|_M(a) = \times$.

(b) If the last condition holds then the least X such that $\|Suc^c(\kappa, \lambda, (X)F)\|_{\mathbf{M}} = 1$ is the X defined in (1) (b).

16.9. Remark. (1) Read 15.16 and 15.17 and make the obvious modifications for the present context.

(2) In the rest of the present section we are going to study strictly implicational operators \rightarrow^* and formulas $\kappa \rightarrow^* \delta$ where κ is a (ps)EC and δ is a (ps)ED. In this case we have a non-trivial deduction rule; this rule is combined with the rule for an improvement operator. In this way we obtain a generalization and improvement of [3] §9(c).

16.10. Definition. (1) We introduce the *empty conjunction* $\wedge \emptyset$; this is an open formula and $\|\wedge \emptyset\|_{\mathbf{M}}(a) = 1$ for each object of each model. *Generalized EC's* (GEC's) are EC's together with the empty conjunction. The same for *generalized psEC's*. *GCD* denotes the set of all pairs $\langle \kappa, \delta \rangle$ where κ is a generalized psEC and δ is a psED. $\langle \kappa, \delta \rangle \in GCD$ is an elementary GCD-pair (consisting of a GEC and an ED) if κ is a GEC, δ is an ED and have no functors in common. *EGCD* is the set of all elementary GCD-pairs.

(2) $\langle \kappa_1, \delta_1 \rangle (\triangleleft \triangleleft) \langle \kappa_2, \delta_2 \rangle$ if $\kappa_1 \triangleleft \kappa_2$ and $\delta_1 \triangleleft \delta_2$ (cf. 14.8)

(3) $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by *specification* if either $\langle \kappa_1, \delta_1 \rangle = \langle \kappa_2, \delta_2 \rangle$ or there is an ED δ_0 such that δ_2 is $\delta_1 \vee \delta_0$ and κ_1 is $\kappa_2 \& neg(\delta_0)$. $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by *reduction* if $\kappa_1 = \kappa_2$ and $\delta_1 \triangleleft \delta_2$. We put $\langle \kappa_1, \delta_1 \rangle SpRd \langle \kappa_2, \delta_2 \rangle$ if there is a $\langle \kappa_3, \delta_3 \rangle$ such that $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_3, \delta_3 \rangle$ by specification and $\langle \kappa_3, \delta_3 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by reduction. (For $\langle \kappa_i, \delta_i \rangle \in EGCD$.)

16.11. Theorem. Let $\langle \kappa_1, \delta_1 \rangle, \langle \kappa_2, \delta_2 \rangle \in EGCD$. If $\langle \kappa_1, \delta_1 \rangle SpRd \langle \kappa_2, \delta_2 \rangle$, if \rightarrow^* is a strictly implicational operator and if $\|\kappa_1 \rightarrow^* \delta_1\|_{\mathbf{M}} = 1$ then $\|\kappa_2 \rightarrow^* \delta_2\|_{\mathbf{M}} = 1$.


Proof. (1) First suppose \mathbf{M} to be a model with complete information. Put $\mathbf{M}_1 = \langle M, \|\kappa_1\|_{\mathbf{M}}, \|\delta_1\|_{\mathbf{M}} \rangle$, $\mathbf{M}_2 = \langle M, \|\kappa_2\|_{\mathbf{M}}, \|\delta_2\|_{\mathbf{M}} \rangle$. One verifies $a_{\mathbf{M}_1} b_{\mathbf{M}_2} \leq a_{\mathbf{M}_2} b_{\mathbf{M}_1}$. This is evident if $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by reduction. If $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by specification denote by m_{ijk} the number of objects a such that $\|\kappa_2\|_{\mathbf{M}}(a) = i$, $\|\delta_1\|_{\mathbf{M}}(a) = j$, $\|\delta_0\|_{\mathbf{M}}(a) = k$. Then $a_{\mathbf{M}_1} = m_{110}$, $b_{\mathbf{M}_1} = m_{100}$, $a_{\mathbf{M}_2} = m_{110} + m_{101} + m_{111}$, $b_{\mathbf{M}_2} = m_{100}$. Consequently, $a_{\mathbf{M}_1} b_{\mathbf{M}_2} \leq a_{\mathbf{M}_2} b_{\mathbf{M}_1}$.

(2) Since \rightarrow^* is secured, the validity of the theorem for arbitrary models follows from its validity for models with complete information. Indeed, if N_2 is a completion of $\langle M, \|\kappa_2\|_{\mathbf{M}}, \|\delta_2\|_{\mathbf{M}} \rangle$ then there is a completion N of \mathbf{M} such that $N_2 = \langle M, \|\kappa_2\|_N, \|\delta_2\|_N \rangle$ (here we use the fact that $\langle \kappa, \delta \rangle$ is an elementary pair). Put $N_1 = \langle M, \|\kappa_1\|_N, \|\delta_1\|_N \rangle$. By the assumption $Asf \rightarrow^*(N_1) = 1$ and hence $Asf \rightarrow^*(N_2) = 1$ by (1).

16.12. Definition. A *GCD-closure set* is a closure set $U \subseteq \{0, \times, 1\}$ for GCD and $(\triangleleft \triangleleft)$ satisfying the following condition: $\langle u, v, \bar{u}, \bar{v} \rangle \in U$ implies $u \geq \bar{u}$ and $v \leq \bar{v}$

for each u, v, \bar{u}, \bar{v} . A GCD-closure operator is an operator defined universally by a GCD-closure set. (Cf. 14.8.)

16.13. Theorem. There are two \subseteq -maximal GCD-closure sets such that the corresponding closure operator \ll is an improvement operator for \rightarrow^* , GCD and $(\ll \triangleleft)$. They are defined by the following tables (i), (ii):



Proof. Consider the table in 16.5 and recall that we are interested in quadruples $\langle u_1, u_2, v_1, v_2 \rangle$ such that $u_1 \geq v_1$ and $u_2 \leq v_2$. Since U has to define an improvement operator, $\langle u, v \rangle \in U$ must imply $u \geq_{ib} v$. Consequently, the quadruple $\langle 1, 1, \times, 1 \rangle$ must not be in U . Theorem 14.13, after obvious modifications, yields the following condition necessary and sufficient for U to be a GCD-closure set: there is an equivalence E on $\{0, \times, 1\}^2$ such that writing $\langle u_1, u_2 \rangle \cong \langle v_1, v_2 \rangle$ for $(u_1 \geq v_1$ and $u_2 \leq v_2)$ and $\langle u_1, u_2 \rangle (\& v) \langle v_1, v_2 \rangle$ for $\langle u_1, \& v_1, u_2 \vee v_2 \rangle$ we have:

- (i) $u \equiv_E w$ and $u \cong v \cong w$ implies $u \equiv_E v \equiv_E w$;
- (ii) $u \equiv_E v$ implies $u \equiv_E [u(\& v) v] \equiv_E v$;
- (iii) $u U v$ iff $(u \cong v$ and $u \equiv_E v)$.

Consequently, the equivalence defined by the tables (i) and (ii) above determine the only two \subseteq -maximal GCD-closure sets not containing the quadruple $\langle 1, 1, \times, 1 \rangle$.

16.14. Remark and Definition. We want now to combine 16.11 with 16.13, so we assume \rightarrow^* to be strictly implicational. Denote for a moment by \ll_1 the operator defined by 16.13 (i) and similarly for \ll_2 . Having a model M and a formula $\kappa \rightarrow^* \delta$ where $\langle \kappa, \delta \rangle \in EGCD$ (say, $\kappa \rightarrow^* \delta$ is an elementary \ast -implication), let $\|\kappa \rightarrow^* \delta\|_M = 1$. Then we can use 16.11 to obtain a formula $\kappa_1 \rightarrow^* \delta_1$ such that $\langle \kappa, \delta \rangle SpRd \langle \kappa_1, \delta_1 \rangle$, hence $\|\kappa_1 \rightarrow^* \delta_1\|_M = 1$, then find a $\langle \kappa_2, \delta_2 \rangle$ such that $\|(\kappa_1, \delta_1) \ll_1 (\kappa_2, \delta_2)\|_M = 1$, hence $\|\kappa_2 \rightarrow^* \delta_2\|_M = 1$, and perhaps iterate this procedure. The following theorem shows that it suffices to use first the improvement operator with a fixed succedent and then to use 16.11; furthermore, that in the present context \ll_1 is better. So we are led to a relation of immediate consequence and to the definition of a pleasant GUA-problem.

16.15. Lemma. (1) Let $i = 1, 2$, let $\langle \kappa_j, \delta_j \rangle \in EGCD$ ($j = 0, 1, 2$) and suppose $\|\kappa_0 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$, $\langle \kappa_0, \delta_0 \rangle SpRd \langle \kappa_1, \delta_1 \rangle$, $\langle \kappa_1, \delta_1 \rangle (\llcorner \lrcorner) \langle \kappa_2, \delta_2 \rangle$ and $\|(\kappa_1, \delta_1) \llcorner_i (\kappa_2, \delta_2)\|_{\mathbf{M}} = 1$. Then there is an EC κ_3 such that $\langle \kappa_0, \delta_0 \rangle (\llcorner \lrcorner) \langle \kappa_3, \delta_0 \rangle$, $\|(\kappa_0, \delta_0) \llcorner_i (\kappa_3, \delta_0)\|_{\mathbf{M}} = 1$ and $\langle \kappa_3, \delta_0 \rangle SpRd \langle \kappa_2, \delta_2 \rangle$.

(2) If $\langle \kappa_0, \delta \rangle, \langle \kappa_1, \delta \rangle \in EGCD$ and if $\kappa_0 \llcorner \kappa_1$ then $\|(\kappa_0, \delta) \llcorner_2 (\kappa_1, \delta)\|_{\mathbf{M}} = 1$ implies $\|(\kappa, \delta) \llcorner_1 (\kappa_1, \delta)\|_{\mathbf{M}} = 1$.

Proof. (1) Consider the following diagram:

$$\begin{array}{ccc}
 \kappa_0, \delta_0 & \xrightarrow{SpRd} & \kappa_1, \delta_1 \\
 \llcorner \downarrow & & \downarrow \llcorner \\
 \kappa^+, \delta_0 & \xrightarrow{SpRd} & \kappa_2, \delta_2
 \end{array}$$

We look for a κ^+ such that the diagram commutes. We have the following relations: $\kappa_1 \subseteq \kappa_0$, $\delta_0 \llcorner \delta_1$, $\kappa_1 \llcorner \kappa_2$, $\delta_1 \llcorner \delta_2$. Hence $\delta_0 \llcorner \delta_2$. The κ^+ we have to find must satisfy the following: $\kappa_2 \subseteq \kappa^+$, $\kappa_0 \llcorner \kappa^+$. Put $\kappa^+ = \kappa_0 \& \kappa_2$ (transformed into an EC: if F occurs both in κ_0 and in κ_2 , take the intersection of coefficients). κ^+ is indeed an EC: if $(X)F$ is in κ_0 then either $(X)F$ is in κ_1 , $(Z)F$ is in κ_2 for some $Z \subseteq X$ and $Z \neq \emptyset$ since \rightarrow^* is strictly implicational and hence $(\exists a \in M) (\|\kappa_0 \& \delta_0\|_{\mathbf{M}}(a) = 1$ (since $\|\kappa_0 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$). Note that $\langle 1, 1, \times, 1 \rangle \notin U$ and cf. the proof of 14.16. Or $(X)F$ is not in κ_1 , then F occurs in δ_1 and hence in δ_2 and consequently F does not occur in κ_2 (since $\langle \kappa_2, \delta_2 \rangle \in EGCD$).

We prove $\|(\kappa_0, \delta_0) \llcorner_i (\kappa^+ \delta_0)\|_{\mathbf{M}} = 1$. Let $i = 1$. We have to verify: If $\|\kappa_0 \& \delta_0\|_{\mathbf{M}}(a) = 1$ then $\|\kappa^+\|_{\mathbf{M}}(a) = 1$. Suppose $\|\kappa_0 \& \delta_0\|_{\mathbf{M}}(a) = 1$; then obviously $\|\kappa_1 \& \delta_1\|_{\mathbf{M}}(a) = 1$ (since $\kappa_1 \subseteq \kappa_0$ and $\delta_0 \llcorner \delta_1$). Then $\|\kappa_2 \& \delta_2\|_{\mathbf{M}}(a) = 1$ (using \llcorner_i) and so $\|\kappa^+\|_{\mathbf{M}}(a) = \|\kappa_0 \& \kappa_2\|_{\mathbf{M}}(a) = 1$. Let $i = 2$. We have to verify: If $\|\kappa_0\|_{\mathbf{M}}(a) = 1$ then $\|\kappa^+\|_{\mathbf{M}}(a) = 1$. Let $\|\kappa_0\|_{\mathbf{M}}(a) = 1$. Then $\|\kappa_1\|_{\mathbf{M}}(a) = 1$ and hence $\|\kappa_2\|_{\mathbf{M}}(a) = 1$ (using \llcorner_2). Hence $\|\kappa^+\|_{\mathbf{M}}(a) = 1$. This completes the proof of (1).

(2) Suppose $\|(\kappa_0, \delta) \llcorner_2 (\kappa_1, \delta)\|_{\mathbf{M}} = 1$ and $\kappa_0 \llcorner \kappa_1$. Consequently, for each a , $\|\kappa_0\|_{\mathbf{M}}(a) = 1$ implies $\|\kappa_1\|_{\mathbf{M}}(a) = 1$. Now, $\|(\kappa_0, \delta) \llcorner_1 (\kappa_1, \delta)\|_{\mathbf{M}} = 1$ means: for each a , if $\|\kappa_0\|_{\mathbf{M}}(a) = 1$ and $\|\delta\|_{\mathbf{M}}(a) = 1$ then $\|\kappa_1\|_{\mathbf{M}}(a) = 1$. So the assertion follows.

16.16. Remark and Definition. In the sequel, \llcorner^d denotes the operator defined by 16.13 (i) If $\langle \kappa, \delta \rangle \in EGCD$ then $Reg^d \llcorner^d(\kappa, \delta)$ is the \llcorner -supremum of all EC's $\bar{\kappa}$ not containing any functor from δ and such that $\kappa \llcorner \bar{\kappa}$ and $\|(\kappa, \delta) \llcorner^d(\bar{\kappa}, \delta)\|_{\mathbf{M}} = 1$. The following is proved in the usual way (cf. 15.14, 15.15 and 16.7, 16.8):

16.17. Theorem. (1) Let $\|Ani^d(\kappa, \delta, \kappa')\|_M = \|(\kappa, \delta) \ll^d (\kappa \& \kappa', \delta)\|_M$. Then Ani^d is universally defined by the set

$$U_A^d = \{\langle u, v, \bar{u} \rangle \in \{0, \times, 1\}^3; [\langle u, v \rangle = \langle 1, 1 \rangle \rightarrow \bar{v} = 1]\}.$$

(2) Let $\langle \kappa, \delta \rangle \in EGCD$ and let M be a model.

(a) There is an X such that $\|Ani^d(\kappa, \delta, (X)F)\|_M = 1$ iff there is no $a \in M$ such that $\|\kappa \& \delta\|_M(a) = 1$ and $\|F\|_M(a) = \times$.

(b) Suppose that the last condition holds; then the least X such that $\|Ani^d(\kappa, \delta, (X)F)\|_M = 1$ is

$$X = \{\|F\|_M(a); \|\kappa \& \delta\|_M(a) = 1\}.$$

(c) Put $\bar{\kappa} = Reg^0 \ll^d_M(\kappa, \delta)$ then $\bar{\kappa}$ is M -incompressible. Moreover, $\bar{\kappa}$ is strongly M_δ -incompressible, where M_δ is the submodel of M formed by all objects $a \in M$ such that $\|\delta\|_M(a) = 1$.

(d) If $(\exists a \in M)(\|\kappa \& \delta\|_M = 1)$ then $\langle \bar{\kappa}, \delta \rangle \in EGCD$.

Proof. (1) and (2) (a), (b) as usual.

(c) The assertion about M_δ -incompressibility is proved as 14.17. (Show: if $\|(\kappa, \delta) \ll^d \ll^d(\kappa \& (X)F, \delta)\|_M = 1$, $\|\kappa \& (X)F\|_{M_\delta} = 1$ and $X_0 \subseteq X$ then $\|(\kappa, \delta) \ll^d(\kappa \& (X_0)F, \delta)\|_M = 1$.)

(d) Assume $(\exists a \in M)(\|\kappa \& \delta\|_M = 1)$. It follows by (b) that $\bar{\kappa}$ is a GEC; by definition $\bar{\kappa}$ does not contain any functor from δ and so $\langle \bar{\kappa}, \delta \rangle \in EGCD$.

16.18. Discussion. We apply the above results to the description of a GUHA-problem and its solution. Put $F = \{\kappa \rightarrow^* \delta; \langle \kappa, \delta \rangle \in EGCD\}$ (relevant questions are elementary *-implications). Call $\kappa \rightarrow^* \delta$ M -prime if (i) $\|\kappa \rightarrow^* \delta\|_M = 1$,

(ii) there is no $\langle \kappa_0, \delta_0 \rangle SpRd \langle \kappa, \delta \rangle$, such that $\langle \kappa_0, \delta_0 \rangle \neq \langle \kappa, \delta \rangle$, $\|\kappa_0 \rightarrow^* \delta_0\|_M = 1$,

(iii) there is no $\kappa_0 \leq \kappa$, $\kappa_0 \neq \kappa$ such that $\|\kappa_0 \rightarrow^* \delta\|_M = 1$ and $\|Ani^d(\kappa_0, \delta, \kappa)\|_M = 1$. Put

$$IC = \left\{ \frac{\kappa \rightarrow^* \delta \& Ani^d(\kappa, \delta, \bar{\kappa})}{\kappa_1 \rightarrow^* \delta_1}; \begin{array}{l} \text{there is a } \kappa_0, \kappa \leq \kappa_0 \leq \bar{\kappa} \text{ such} \\ \text{that } \langle \kappa_0, \delta \rangle SpRd \langle \kappa_1, \delta_1 \rangle \end{array} \right\}.$$

For each M , put $X_M = \{\kappa \rightarrow^* \delta \& Ani^d(\kappa, \delta, \bar{\kappa}); \kappa \rightarrow^* \delta \text{ } M\text{-prime and } \bar{\kappa} = Reg^0 \ll^d_M(\kappa, \delta)\}$.

16.19. Theorem. Under the denotations of 16.18, X_M is a (\subseteq -minimal) solution of $P = \langle F, \{1\}, IC \rangle$ in M .

Proof. Let $\|\kappa \rightarrow^* \delta\|_M = 1$ and let $\kappa \rightarrow^* \delta$ be not M -prime. Let $\langle \kappa_1, \delta_1 \rangle$ be a $SpRd$ -minimal EGCD such that $\|\kappa_1 \rightarrow^* \delta_1\|_M = 1$ and $\langle \kappa_1, \delta_1 \rangle SpRd \langle \kappa, \delta \rangle$.

Let $\langle \kappa_2, \delta_1 \rangle$ be a (\ll -)minimal EGCD such that $\|\kappa_2, \delta_1\|_{\mathbf{M}} \ll^d (\kappa_1, \delta_1)_{\mathbf{M}} = 1$ and $\|\kappa_2 \rightarrow^* \delta_1\|_{\mathbf{M}} = 1$. Then obviously

$$\frac{\kappa_2 \rightarrow^* \delta_1 \& \text{Ant}^d(\kappa_2, \delta_1, \text{Reg}^0 \ll_{\mathbf{M}}^d(\kappa_2, \delta_1))}{\kappa \rightarrow^* \delta} \in IC.$$

We show that $(\kappa_2 \rightarrow^* \delta_1)$ is prime. Evidently, there is no κ_3 such that $\langle \kappa_3, \delta_1 \rangle (\ll \ominus)$ $\langle \kappa_2, \delta_1 \rangle$, $\|\kappa_3 \rightarrow^* \delta_1\|_{\mathbf{M}} = 1$ and $\|\text{Ant}^d(\kappa_3, \delta_1, \kappa_2)\|_{\mathbf{M}} = 1$. Suppose that there is a $\langle \kappa_3, \delta_3 \rangle \text{SPRD} \langle \kappa_2, \delta_2 \rangle$ such that $\|\kappa_3 \rightarrow^* \delta_3\|_{\mathbf{M}} = 1$. Then by 16.15 there is a κ^+ such that $\|\text{Ant}^d(\kappa_3, \delta_3, \kappa^+)\|_{\mathbf{M}} = 1$ and $\langle \kappa^+, \delta_3 \rangle \text{SPRD} \langle \kappa_1, \delta_1 \rangle$, hence $\langle \kappa^+, \delta_3 \rangle \text{SPRD} \langle \kappa, \delta \rangle$ and $\|\kappa^+ \rightarrow^* \delta_3\|_{\mathbf{M}} = 1$. It follows $\langle \kappa^+, \delta_3 \rangle = \langle \kappa_1, \delta_1 \rangle$, and hence $\langle \kappa_2, \delta_3 \rangle = \langle \kappa_2, \delta_1 \rangle$. Hence $\langle \kappa_2, \delta_1 \rangle$ is \mathbf{M} -prime.

16.20. Discussion. We consider the situation of V_{reg} being big. Put $F_k = \{\kappa \rightarrow^* \delta; \langle \kappa, \delta \rangle \in \text{EGCD}$ and all coefficients have at most k elements}. Suppose $2k < \text{card}(V_{\text{reg}})$; under this assumption we have the following: If $\langle \kappa_1, \delta_1 \rangle \neq \langle \kappa_2, \delta_2 \rangle$ and $\langle \kappa_1, \delta_1 \rangle$ results from $\langle \kappa_2, \delta_2 \rangle$ by specification then $\kappa_1 \rightarrow^* \delta_1$ and $\kappa_2 \rightarrow^* \delta_2$ cannot be simultaneously in F_k . Hence we give up any specification and consider a weaker rule

$$IC_1 = \left\{ \frac{(\kappa \rightarrow^* \delta) \& \text{Ant}^d(\kappa, \delta, \bar{\kappa})}{\kappa_1 \rightarrow \delta_1}; \kappa \ll \bar{\kappa}, \delta \ominus \delta_1 \right\}.$$

Define $\kappa \rightarrow^* \delta$ to be $\mathbf{M} - F_k$ -prime if

- (i) $(\kappa \rightarrow^* \delta) \in F_k$ and $\|\kappa \rightarrow^* \delta\|_{\mathbf{M}} = 1$,
- (ii) there is no $\kappa_1 \ll \kappa, \kappa_1 \neq \kappa$ such that $(\kappa_1 \rightarrow^* \delta) \in F_k$ and $\|\kappa_1 \rightarrow^* \delta\|_{\mathbf{M}} = 1$,
- (iii) there is no $\delta_0 \ominus \delta$ such that $\|\kappa_1 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$ (irreducibility).

Let $\text{Reg}^0 \ll_{\mathbf{M},k}^d(\kappa, \delta) = \bar{\kappa}$ if $\bar{\kappa}$ is the conjunction of all $(X) F$ such that $\text{card}(X) \leq k$ and X minimal such that $\|\text{Ant}^d(\kappa, \delta, (X) F)\|_{\mathbf{M}} = 1$. Put

$$X_{\mathbf{M}}^k = \{(\kappa \rightarrow^* \delta) \& \text{Ant}^d(\kappa, \delta, \bar{\kappa}); \kappa \rightarrow^* \delta, \mathbf{M} - F_k\text{-prime and } \bar{\kappa} = \text{Reg}^0 \ll_{\mathbf{M},k}^d(\kappa, \delta)\}.$$

16.21. Theorem. Under the denotations of 16.20, $X_{\mathbf{M}}^k$ is a (\subseteq -minimal solution) of $P_k = \langle F_k, \{1\}, IC_1 \rangle$ in \mathbf{M} .

Proof. Let $(\kappa \rightarrow^* \delta) \in F_k$, $\|\kappa \rightarrow^* \delta\|_{\mathbf{M}} = 1$. Let δ_0 be a \ll -minimal disjunction such that $\|\kappa \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$. Let κ_0 be a \ll -minimal element such that $\kappa_0 \ll \kappa$, $\|\kappa_0 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$, $(\kappa_0 \rightarrow^* \delta_0) \in F_k$ and $\|(\kappa_0, \delta_0) \ll^d(\kappa, \delta_0)\|_{\mathbf{M}} = 1$. Then $(\kappa \rightarrow^* \delta) \in IC_1(\kappa_0 \rightarrow^* \delta_0)$; we prove that for $\bar{\kappa} = \text{Reg}^0 \ll_{\mathbf{M},k}^d(\kappa_0, \delta_0)$ we have: $\kappa_0 \ll \kappa \ll \bar{\kappa}$ and $(\kappa_0 \rightarrow^* \delta_0)$ is $\mathbf{M} - F_k$ -prime. The assertion concerning \ll is obvious. $(\kappa_0 \rightarrow^* \delta_0) \in F_k$, $\|\kappa_0 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$ by the definition of κ_0 ; evidently, there is no $\kappa_1 \ll \kappa_0$, $\kappa_1 \neq \kappa_0$ such that $\|\kappa_1 \rightarrow^* \delta_0\|_{\mathbf{M}} = 1$ ($\kappa_1 \rightarrow^* \delta_0) \in F_k$ and $\|(\kappa_1, \delta_0) \ll^d(\kappa_0, \delta_0)\|_{\mathbf{M}} = 1$. If there is a δ_1 such that $\delta_1 \ominus \delta_0$, $\delta_1 \neq \delta_0$ and $\|\kappa_0 \rightarrow^* \delta_1\|_{\mathbf{M}} = 1$ then one shows as in 16.15 (1) such that $\langle \kappa, \delta_1 \rangle$ results from $\langle \kappa, \delta_0 \rangle$ by reduction and that $\|(\kappa_0, \delta_1) \ll^d(\kappa, \delta_1)\|_{\mathbf{M}} = 1$. Hence $(\kappa_0 \rightarrow^* \delta_0)$ is $\mathbf{M} - F_k$ -prime. This completes the proof.

16.22. Remark and Definition. As an analogy to 15.19 (2) we consider the following diagram:

123



This is the operator (ii) from 16.13; it will be denoted by \ll^e . Define $Reg^0 \ll_M^e(\kappa, \delta)$ as in 16.16 as the \ll -supremum of all $\bar{\kappa}$ such that $\kappa \ll \bar{\kappa}$ and $\|(\kappa, \delta) \ll^e (\bar{\kappa}, \delta)\|_M = 1$. We have the following

16.23. Theorem. (1) If Ant^e has the usual meaning then Ant^e is universally defined by

$$U_A^e = \{\langle u, v, \bar{u} \rangle; u = 1 \rightarrow \bar{\kappa} = 1\}.$$

(2) There is an X such that $\|Ant^e(\kappa, \delta, (X)F)\|_M = 1$ iff there is no $a \in M$ such that $\|\kappa\|_M(a) = 1$ and $\|F\|_M(a) = \times$. If the last condition holds the least X such that $\|Ant^e(\kappa, \delta, (X)F)\|_M = 1$ is $X = \{\|F\|_M(a); \|\kappa\|_M(a) = 1\}$.

(3) If $\langle \kappa, \delta \rangle \in EGCD$ and $\bar{\kappa} = Reg^0 \ll_M^e(\kappa, \delta)$ then $\bar{\kappa}$ is strongly M -incompressible and $\|\kappa \approx \bar{\kappa}\|_M = 1$; if $(\exists a \in M)(\|\kappa\|_M(a) = 1)$ then $\bar{\kappa}$ is an EC and $\langle \kappa, \delta \rangle \in EGCD$.

Proof. The proofs of (1) and (2) are routine and completely analogous to previous proofs of analogous theorems. Then it follows by (2) that if $\langle \kappa, \delta \rangle \in EGCD$ then $\bar{\kappa}$ is an GEC and hence $\langle \bar{\kappa}, \delta \rangle \in EGCD$. To prove the strong incompressibility of $\bar{\kappa}$ one observes the following: If $\|(\kappa, \delta) \ll^e (\kappa \& (X)F, \delta)\|_M = 1$ and if $X_0 \subseteq X$, $\|\kappa \& (X)F \approx \kappa \& (X_0)F\|_M = 1$ then $\|(\kappa, \delta) \ll^e (\kappa \& (X_0)F, \delta)\|_M = 1$. The assertion concerning \approx is obvious.

16.24. Corollary. If κ is strongly incompressible and if $\bar{\kappa} = Reg^0 \ll_M^e(\kappa, \delta)$ then $\kappa \subseteq \bar{\kappa}$.

Proof. We have $\kappa \ll \bar{\kappa}$ and $\|\kappa \approx \bar{\kappa}\|_M = 1$, hence if κ_0 is the subconjunction of κ having the same functors as κ then κ_0 is poorer than κ and $\|\kappa \approx \kappa_0\|_M = 1$, hence $\kappa = \kappa_0$ by strong incompressibility.

16.25. Definition and Theorem. Let $SInc(\kappa)$ be a formula such that $\|SInc(\kappa)\|_M = 1$ iff κ is strongly M -incompressible. Let $F^* = \{(\kappa \rightarrow^* \delta) \& SInc(\kappa); \langle \kappa, \delta \rangle \in EGCD\}$ and

$$IC^* = \left\{ \frac{(\kappa \rightarrow^* \delta) \& SInc(\kappa) \& (\kappa, \delta) \ll^e (\bar{\kappa}, \delta)}{\kappa_1 \rightarrow^* \delta_1 \& SInc(\kappa_1)}; \kappa \subseteq \kappa_1 \subseteq \kappa, \delta \triangleleft \delta_1 \right\}.$$

For each M let X_M^* be the set of all formulas $(\kappa \rightarrow^* \delta) \& SInc(\kappa) \& (\kappa, \delta) \ll^e (\bar{\kappa}, \delta)$ such that $\kappa \rightarrow^* \delta$ is M -prime (in the sense of 16.18 but with the present meaning

124 of \ll and Ant), κ strongly M -incompressible and $\bar{\kappa} = Reg^0 \ll_M^e(\kappa, \delta)$. Then X_M^* is a (\subseteq -minimal) solution of $P^* = \langle F^*, \{1\}, IC^* \rangle$ in M .

(Received October 8, 1973.)

REFERENCES

- [1] J. P. Cleave: The notion of logical consequence in the logic of inexact predicates. *Zeitschr. f. Math. Logik* (to appear).
- [2] P. Hájek: Automatic Listing of Important Observational Statements I. *Kybernetika* 9 (1973), 187–205.
- [3] P. Hájek: Automatic Listing of Important Observational Statements II. *Kybernetika* 9 (1973), 251–271.
- [4] P. Hájek: Some logical problems of automated research. In: *Math. Foundations of Computer Science (Proceedings)*, High Tatras 1973, 85–93.
- [5] P. Hájek, K. Bendová, Z. Renc: The GUHA method and the three valued logic. *Kybernetika* 7 (1971), 421–435.
- [6] S. Körner: *Experience and theory*. London 1966.
- [7] C. R. Rao: *Linear statistical inference and its applications*. New York 1965.

Dr. Petr Hájek, CSc.; Matematický ústav ČSAV (Mathematical Institute — Czechoslovak Academy of Sciences), Žitná 25, 115 67 Praha 1, Czechoslovakia.