

Bohdan Zelinka

Signed 2-domination in caterpillars

Mathematica Bohemica, Vol. 129 (2004), No. 4, 393–398

Persistent URL: <http://dml.cz/dmlcz/134049>

Terms of use:

© Institute of Mathematics AS CR, 2004

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

SIGNED 2-DOMINATION IN CATERPILLARS

BOHDAN ZELINKA, Liberec

(Received December 19, 2003)

Abstract. A caterpillar is a tree with the property that after deleting all its vertices of degree 1 a simple path is obtained. The signed 2-domination number $\gamma_s^2(G)$ and the signed total 2-domination number $\gamma_{st}^2(G)$ of a graph G are variants of the signed domination number $\gamma_s(G)$ and the signed total domination number $\gamma_{st}(G)$. Their values for caterpillars are studied.

Keywords: caterpillar, signed 2-domination number, signed total 2-domination number

MSC 2000: 05C69, 05C05

This paper concerns caterpillars. A caterpillar [1] is a tree with the property that after deleting all its vertices of degree 1 a simple path is obtained. According to this definition a caterpillar has at least three vertices. But we need not care about graphs with one or two vertices. For such graphs our considerations are trivial.

Let G be a caterpillar. The mentioned simple path will be denoted by B and called the body of the caterpillar G . Let the number of vertices of B be m . Let a_1, \dots, a_m be these vertices and let $a_i a_{i+1}$ for $i = 1, \dots, m - 1$ be the edges of B . By $[m]$ we shall denote the set of integers i such that $1 \leq i \leq m$. For each $i \in [m]$ let s_i be the degree of a_i in G . The vector $\vec{s} = (s_1, \dots, s_m)$ will be called the degree vector of the caterpillar G .

Now we shall define variants of the signed domination number and of the signed total domination number [2] of a graph. For a vertex $u \in V(G)$ the symbol $N(u)$ denotes the open neighbourhood of u in G , i.e. the set of all vertices which are adjacent to u in G . The closed neighbourhood of u is $N[u] = N(u) \cup \{u\}$. Similarly the open 2-neighbourhood $N^2(u)$ is the set of all vertices having the distance 2 from u in G . The closed 2-neighbourhood of u is $N^2[u] = N[u] \cup N^2(u)$. If f is a mapping

of $V(G)$ into some set of numbers and $S \subseteq V(G)$, then $f(S) = \sum_{x \in S} f(x)$ and the weight of f is $w(f) = f(V(G)) = \sum_{x \in V(G)} f(x)$.

Let $f: V(G) \rightarrow \{-1, 1\}$. If $f(N^2[u]) \geq 1$ (or $f(N^2(u)) \geq 1$) for each $u \in V(G)$, then f is called a signed 2-dominating (or signed total 2-dominating, respectively) function on G . The minimum of weights $w(f)$ taken over all signed 2-dominating (or all signed total 2-dominating) functions f is the signed 2-dominating number $\gamma_s^2(G)$ (or the signed total 2-domination number $\gamma_{st}^2(G)$, respectively) of G .

For each $i \in [m]$ let $t_i \in \{1, 2\}$ and $t_i \equiv s_i + 1 \pmod{2}$.

We shall prove a theorem concerning $\gamma_s^2(G)$.

Theorem 1. *Let G be a caterpillar with the degree vector $\vec{s} = (s_1, \dots, s_m)$ such that $n \geq 2$ and $s_i \geq 3$ for all $i \in [m]$. Then*

$$\gamma_s^2(G) = \sum_{i=1}^m t_i - 2m + 2.$$

Proof. Consider a vertex a_i with $i \in [m]$. As $s_i \geq 3$, there exists at least one vertex $u \in N(a_i)$ which does not belong to B and has degree 1. Then $N^2[u] = N[a_i]$. Let f be a signed 2-dominating function on G . Then $f(N^2[u]) = f(N[a_i]) \geq 1$. The set $N[a_i]$ has $s_i + 1$ vertices. If s_i is even, then $s_i + 1$ is odd. At least $\frac{1}{2}(s_i + 2) = \frac{1}{2}s_i + 1$ vertices of $N[a_i]$ must have the value 1 in f and at most $\frac{1}{2}s_i$ of them may have the value -1 . Then $f(N^2[u]) \geq (\frac{1}{2}s_i + 1) - \frac{1}{2}s_i = 1 = t_i$. If s_i is odd, then $s_i + 1$ is even and at least $\frac{1}{2}(s_i + 1) + 1$ vertices of $N[a_i]$ must have the value 1 in f and at most $\frac{1}{2}(s_i + 1) - 1$ of them may have the value -1 . Then $f(N^2[u]) \geq 2 = t_i$. We may easily construct the function f such that it has the value -1 in exactly $\frac{1}{2}s_i$ vertices of degree 1 in $N[a_i]$ with i even and in exactly $\frac{1}{2}(s_i + 1) - 1 = \frac{1}{2}(s_i - 1)$ vertices of degree 1 in $N[a_i]$ with i odd. In all other vertices (including all vertices of the body) the function f has the value 1.

We have $\bigcup_{i=1}^m N[a_i] = V(G)$. The vertex a_1 is contained in exactly two sets $N[a_i]$, namely in $N[a_1]$ and $N[a_2]$. Similarly a_m is contained in exactly two sets $N[a_{m-1}]$, $N[a_m]$. For $i \in [m] - \{1, m\}$ the vertex a_i is contained in exactly three sets $N[a_{i-1}]$, $N[a_i]$, $N[a_{i+1}]$. Each vertex outside the body is contained in exactly one of these sets. By the Inclusion-Exclusion Principle we have

$$\begin{aligned} w(f) &= f(V(G)) = \sum_{i=1}^m f(N[a_i]) - 2 \sum_{i=2}^{m-1} f(a_i) - f(a_1) - f(a_m) \\ &= \sum_{i=1}^m t_i - 2(m-2) - 1 - 1 = \sum_{i=1}^m t_i - 2m + 2. \end{aligned}$$

As f is the minimum function satisfying the requirements, we have

$$\gamma_s^2(G) = w(f) = \sum_{i=1}^m t_i - 2m + 2.$$

□

An analogous theorem concerns $\gamma_{st}^2(G)$.

Theorem 2. *Let G be a caterpillar with the degree vector $\vec{s} = (s_1, \dots, s_m)$ such that $m \geq 2$ and $s_i \geq 4$ for all $i \in [m]$. Then*

$$\gamma_{st}^2(G) = \sum_{i=1}^m t_i + 2.$$

Proof. Consider a vertex a_i with $i \in [m]$. As $s_i \geq 5$, there exists at least one vertex $u \in N(a_i)$ which does not belong to B and has degree 1. Then $N^2(u) = N(a_i) - \{u\}$. Let f be a signed total 2-dominating function on G . Then $f(N^2(u)) = f(N(a_i) - \{u\}) \geq 1$. The set $N(a_i) - \{u\}$ has $s_i - 1$ vertices. If s_i is even, then $s_i - 1$ is odd. At least $\frac{1}{2}s_i$ vertices of $N(a_i) - \{u\}$ must have the value 1 in f and at most $\frac{1}{2}(s_i - 2) = \frac{1}{2}s_i - 1$ of them may have the value -1 . Then $f(N^2(u)) \geq \frac{1}{2}s_i - (\frac{1}{2}s_i - 1) = 1 = t_i$. If s_i is odd, then $s_i - 1$ is even and at least $\frac{1}{2}(s_i - 1) + 1$ vertices of $N(a_i) - \{u\}$ must have the value 1 in f and at most $\frac{1}{2}(s_i - 1) - 1$ of them may have the value -1 . Then $f(N^2(u)) \geq 2 = t_i$. As $s_i \geq 5$ for $i \in [m]$, in both these cases we must admit the possibility $f(u) = 1$. Then in the case of s_i even we have $f(N(a_i)) \geq 2 = t_i + 1$ and in the case of s_i odd we have $f(N(a_i)) \geq 3 = t_i + 1$. We may easily construct the function f such that it has the value -1 in $\frac{1}{2}s_i - 1$ vertices of degree 1 in $N(a_i)$ for s_i even, in $\frac{1}{2}(s_i - 1) - 1 = \frac{1}{2}(s_i - 3)$ vertices of degree 1 in $S(a_i)$ for s_i odd and the value 1 for all other vertices (including all vertices of B). Each vertex a_j for $j \in [m] - \{1, m\}$ is contained in two sets $N(a_i)$, namely in $N(a_{j-1})$ and $N(a_{j+1})$. Each other vertex is contained in exactly one set $N(a_i)$. Again by the Inclusion-Exclusion Principle we have

$$\begin{aligned} w(f) &= f(V(G)) = \sum_{i=1}^m f(N(a_i)) - \sum_{i=2}^{m-1} f(a_i) \\ &= \sum_{i=1}^m (t_i + 1) - (m - 2) = \sum_{i=1}^m t_i + m - (m - 2) = \sum_{i=1}^m t_i + 2. \end{aligned}$$

As f is the minimum function satisfying the requirements, we have

$$\gamma_{st}^2(G) = w(f) = \sum_{i=1}^m t_i + 2.$$

□

In Figs. 1 and 2 a caterpillar G with the degree vector $(5, 6, 7)$ is depicted. We have $t_1 = t_3 = 2, t_2 = 1$ and therefore $\gamma_{st}^2(G) = 7$ and $\gamma_s^2(G) = 1$. In Fig. 1 the values of the corresponding signed total 2-dominating function are illustrated; in the vertices denoted by $+$ the value is 1 and in the vertices denoted by $-$ it is -1 . Similarly in Fig. 2 the corresponding signed 2-dominating function is illustrated.




Fig. 1.




Fig. 2.

In Theorems 1 and 2 we had the assumption $m \geq 2$. The following proposition concerns the singular case $m = 1$.

Proposition 1. *Let G be a caterpillar with the body consisting of one vertex, i.e. a star with the central vertex a_1 and with $s_1 \geq 2$ vertices of degree 1. Then $\gamma_{st}^2(G)$ is undefined and $\gamma_s^2(G) = t_1$.*

Proof. The open 2-neighbourhood $N^2(a_1) = \emptyset$ and thus $f(N^2(a_1)) = 0$ for any function $f: V(G) \rightarrow \{-1, 1\}$, hence none of such functions might be signed total 2-dominating in G . On the other hand, $N^2[a_1] = V(G)$ and $|V(G)| = s_1 + 1$. Analogously as in the proofs of Theorems 1 and 2 we prove that for s_1 even we have $\gamma_s^2(G) = 1 = t_1$ and for s_1 odd we have $\gamma_s^2(G) = 2 = t_1$. \square

Proposition 2. *Let G be a caterpillar with $m \equiv 2 \pmod{5}, m \geq 5, s_i = 3$ for all $i \in [m]$. Then $\gamma_{st}^2(G) \leq \frac{4}{3}(m + 3) + 2$, while $\sum_{i=1}^m t_i + 2 = 2(m + 1)$.*

Proof. As $s_i = 3$ for each $i \in [m]$, we have $t_i = 2$ for each $i \in [m]$. Each vertex a_i for $i \in [m] - \{1, m\}$ is adjacent to exactly one vertex v_i of degree 1. The vertex a_1 is adjacent to two such vertices v_1, w_1 and similarly a_m to v_m, w_m . Let $f: V(G) \rightarrow \{-1, 1\}$ be defined so that $f(v_i) = -1$ for $i \equiv 0 \pmod{3}$ and $f(u) = 1$ for all other vertices u . This is a signed total 2-dominating function on G (this can be easily verified by the reader) and $w(f) = \frac{1}{3}(4m + 10)$. Therefore $\gamma_{st}^2(G) \leq \frac{1}{3}(4m + 10)$, while $\sum_{i=1}^m t_i + 2 = 2(m + 1)$. For $m \geq 3$ we have $\frac{1}{3}(4m + 10) < 2(m + 1)$.

In Fig. 3 we see such a caterpillar for $m = 8$ with the corresponding function f . In this case $\gamma_{st}^2(G) = 14, \sum_{i=1}^m t_i + 2 = 18$. For the signed 2-domination number




Fig. 3.

here Theorem 1 holds. In Fig. 4 the same caterpillar is depicted with the function f realizing the signed domination number $\gamma_s^2(G) = \sum_{i=1}^m t_i - 2m + 2 = 2$. \square




Fig. 4.

Proposition 3. *Let G be a caterpillar with $m \geq 2$ and $s_i = 2$ for each $i \in [m]$. Then $\sum_{i=1}^m t_i - 2m + 2 < \gamma_s^2(G)$, but $\sum_{i=1}^m t_i + 2 = \gamma_{st}^2(G)$.*

Proof. The caterpillar thus described is a simple path of length $m + 1$. It has $m + 2$ vertices. The inequality $\gamma_s^2(G) \leq \sum_{i=1}^m t_i - 2m + 2$ would imply that there exists a signed 2-dominating function f which has the value -1 in m vertices, while the value 1 only in two vertices. This is evidently impossible. On the other hand the open 2-neighbourhood of any vertex consists of at most two vertices and therefore the unique signed total 2-dominating function is the constant function equal to 1 in the whole set $V(G)$. Then

$$\gamma_{st}^2(G) = w(f) = \sum_{i=1}^m t_i + 2 = m + 2.$$

\square

Now we shall study the signed 2-domination number of a simple path P_n with n vertices (i.e. of length $n - 1$). We shall not use the notation for caterpillars used above, but we shall denote the vertices by u_1, \dots, u_n and edges by $u_i u_{i+1}$ for $i = 1, \dots, n - 1$.

Theorem 3. *Let P_n be a path with n vertices. If $n \equiv 0 \pmod{5}$, then $\gamma_s^2(P_n) = \frac{1}{5}n$. In general, asymptotically $\gamma_s^2(P_n) \approx \lfloor \frac{1}{5}n \rfloor$.*

Proof. If $n \equiv 0 \pmod{5}$, then the closed neighbourhood $N^2[u_i] = \{u_{i-2}, u_{i-1}, u_i, u_{i+1}, u_{i+2}\}$ for $i \equiv 3 \pmod{5}$, $3 \leq i \leq n - 2$, form a partition of $V(P_n)$. Let f

be a signed 2-dominating function on P_n . Then f must have the value 1 in at least three vertices and may have the value -1 in at most two vertices of each class of this partition. Then $w(f) \geq \frac{3}{5}n = \frac{1}{5}n$. A function f for which the equality occurs may be defined so that $f(u_i) = -1$ for $i \equiv 0 \pmod{5}$ and $i \equiv 1 \pmod{5}$ and $f(u_i) = 1$ for $i \equiv 2 \pmod{5}$, $i \equiv 3 \pmod{5}$ and $i \equiv 4 \pmod{5}$. Therefore $\gamma_s^2(P_n) = w(f) = \frac{1}{5}n$.

Now let $m \equiv r \pmod{5}$, $r \leq 4$. Let $q = n - r$. We have $q \equiv 0 \pmod{5}$ and thus $\gamma_s^2(P_q) = \frac{1}{5}q$. The path P_n is obtained from P_q by adding a path with r vertices. Let g be a minimum signed 2-dominating function on P_n , let g_0 be its restriction to P_q . We have $w(g_0) = \frac{1}{5}q$. Now the vertices of P_n not in P_q may have values 1 or -1 in g and thus $\frac{1}{5}q - r \leq w(g) \leq \frac{1}{5}q + r$. In general, $\frac{1}{5}q - 4 \leq \gamma_s^2(P_n) \leq \frac{1}{5}q + 4$. This implies

$$\frac{9}{5n} - \frac{4}{n} \leq \frac{\gamma_s^2(P_n)}{n} \leq \frac{9}{5n} + \frac{4}{n}.$$

Therefore $\lim_{n \rightarrow \infty} \frac{\gamma_s^2(P_n)}{n} = \frac{9}{5n}$ and thus $\gamma_s^2(P_n) \approx \frac{9}{5} = \lfloor \frac{n}{5} \rfloor$. □

In Fig. 5 we see a path P_{15} (with $\gamma_s^2(P_{15}) = 3$) in which the corresponding signed 2-dominating function is illustrated.




Fig. 5.

As has already been mentioned, $\gamma_{st}^2(P_n) = n$ for each positive integer n .

Without a proof we shall state the values of $\gamma_s^2(P_n)$ for $n \leq 4$. We have $\gamma_s^2(P_1) = 1$, $\gamma_s^2(P_2) = 2$, $\gamma_s^2(P_3) = 1$, $\gamma_s^2(P_4) = 2$.

References

- [1] *A. Recski*: Maximal results and polynomial algorithms in VLSI routing. Combinatorics, Graphs, Complexity. Proc. Symp. Prachatice, 1990. JČMF Praha, 1990.
- [2] *T. W. Haynes, S. T. Hedetniemi, P. J. Slater*: Fundamentals of Domination in Graphs. Marcel Dekker, New York, 1998.