

Henry Escudro; Ping Zhang  
On detectable colorings of graphs

*Mathematica Bohemica*, Vol. 130 (2005), No. 4, 427–445

Persistent URL: <http://dml.cz/dmlcz/134214>

## Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ON DETECTABLE COLORINGS OF GRAPHS

HENRY ESCUADRO, PING ZHANG, Kalamazoo

(Received July 25, 2005)

*Abstract.* Let  $G$  be a connected graph of order  $n \geq 3$  and let  $c: E(G) \rightarrow \{1, 2, \dots, k\}$  be a coloring of the edges of  $G$  (where adjacent edges may be colored the same). For each vertex  $v$  of  $G$ , the color code of  $v$  with respect to  $c$  is the  $k$ -tuple  $c(v) = (a_1, a_2, \dots, a_k)$ , where  $a_i$  is the number of edges incident with  $v$  that are colored  $i$  ( $1 \leq i \leq k$ ). The coloring  $c$  is detectable if distinct vertices have distinct color codes. The detection number  $\det(G)$  of  $G$  is the minimum positive integer  $k$  for which  $G$  has a detectable  $k$ -coloring. We establish a formula for the detection number of a path in terms of its order. For each integer  $n \geq 3$ , let  $D_u(n)$  be the maximum detection number among all unicyclic graphs of order  $n$  and  $d_u(n)$  the minimum detection number among all unicyclic graphs of order  $n$ . The numbers  $D_u(n)$  and  $d_u(n)$  are determined for all integers  $n \geq 3$ . Furthermore, it is shown that for integers  $k \geq 2$  and  $n \geq 3$ , there exists a unicyclic graph  $G$  of order  $n$  having  $\det(G) = k$  if and only if  $d_u(n) \leq k \leq D_u(n)$ .

*Keywords:* detectable coloring, detection number

*MSC 2000:* 05C15, 05C70

## 1. INTRODUCTION

Let  $G$  be a connected graph of order  $n \geq 3$  and let  $c: E(G) \rightarrow \{1, 2, \dots, k\}$  be a coloring of the edges of  $G$  for some positive integer  $k$  (where adjacent edges may be colored the same). The *color code* of a vertex  $v$  of  $G$  (with respect to  $c$ ) is the ordered  $k$ -tuple

$$c(v) = (a_1, a_2, \dots, a_k) \text{ (or simply, } c(v) = a_1 a_2 \dots a_k),$$

where  $a_i$  is the number of edges incident with  $v$  that are colored  $i$  for  $1 \leq i \leq k$ . Therefore,  $\sum_{i=1}^k a_i = \deg_G v$ . The coloring  $c$  is called *detectable* if distinct vertices have distinct color codes; that is, for every two vertices of  $G$ , there exists a color such

that the number of incident edges with that color is different for these two vertices. The *detection number*  $\det(G)$  of  $G$  is the minimum positive integer  $k$  for which  $G$  has a detectable  $k$ -coloring. Such a coloring is called a *minimum detectable coloring*. Since every nontrivial graph contains at least two vertices having the same degree, the vertices of a nontrivial connected graph cannot be distinguished by their degrees alone. Therefore, every connected graph of order 3 or more has detection number at least 2.

To illustrate these concepts, consider the graph  $G$  shown in Figure 1(a). A coloring of the edges of  $G$  is shown in Figure 1(b). For this 3-coloring  $c$ , the color codes of its vertices are

$$\begin{aligned} c(u) &= 110, & c(v) &= 021, & c(w) &= 210, \\ c(x) &= 201, & c(y) &= 101, & c(z) &= 001. \end{aligned}$$

Since the vertices of  $G$  have distinct color codes,  $c$  is a detectable coloring. Figure 1(c) shows yet another detectable coloring  $c'$  of the graph  $G$  of Figure 1(a). For this coloring,

$$c'(u) = 20, \quad c'(v) = 30, \quad c'(w) = 21, \quad c'(x) = 12, \quad c'(y) = 02, \quad c'(z) = 01.$$

The coloring  $c'$  uses only two colors. Once a detectable 2-coloring for the graph  $G$  of Figure 1(c) was obtained, we can immediately conclude that  $\det(G) = 2$  as every connected graph of order 3 or more has detection number at least 2.




Figure 1. A detectable coloring of a graph

The concept of detectable coloring was studied in [1], [2], [3], [4], [5], inspired by the basic problem in graph theory that concerns finding means to distinguish the vertices of a connected graph. The following results were stated in [2], [5].

**Theorem A.** Let  $c$  be a  $k$ -coloring of the edges of a graph  $G$ . The maximum number of different color codes of the vertices of degree  $r$  in  $G$  is  $\binom{r+k-1}{r}$ .

**Theorem B.** If  $c$  is a detectable  $k$ -coloring of a connected graph  $G$  of order at least 3, then  $G$  contains at most  $\binom{r+k-1}{r}$  vertices of degree  $r$ .

Since vertices with distinct degrees in a connected graph always have distinct color codes, it is most challenging to find minimum detectable colorings of graphs having many vertices of the same degree. The detection numbers of complete graphs and complete bipartite graphs have been determined and detectable colorings of connected  $r$ -regular graphs and trees have been studied as well (see [2], [3], [4], [5]). The detection number of the cycle  $C_n$  of order  $n$  was established in [5].

**Theorem C.** Let  $n \geq 3$  be an integer and let  $l = \lceil \sqrt{n/2} \rceil$ . Then

$$\det(C_n) = \begin{cases} 2l & \text{if } 2l^2 - l + 1 \leq n \leq 2l^2, \\ 2l - 1 & \text{if } 2(l - 1)^2 + 1 \leq n \leq 2l^2 - l. \end{cases}$$

In this work, we first establish a formula for the detection number of paths in Section 2 and then study some extremal problems concerning detection numbers of unicyclic graphs in Section 3. We refer to the book [6] for graph theory notation and terminology not described in this paper.

## 2. DETECTABLE COLORING OF PATHS

In this section, we determine the detection numbers of all paths. In order to do this, we first present four results, the first of which is a consequence of Theorem B, the next two are well-known results in graph theory, and the fourth has a straightforward proof.

**Corollary 2.1.** Let  $k \geq 2$  be an integer. If  $n > \binom{k}{2} + 2$ , then  $\det(P_n) \geq k$ .

**Theorem D.** For each positive integer  $k$ , the complete graph  $K_{2k}$  can be factored into  $k - 1$  Hamiltonian cycles and a 1-factor.

**Theorem E.** For each positive integer  $k$ , the complete graph  $K_{2k+1}$  is Hamiltonian factorable (into  $k$  Hamiltonian cycles).

**Lemma 2.2.** For each integer  $n \geq 3$ , there exists a unique positive integer  $l$  such that

$$\binom{2l}{2} + 2 = 2l^2 - l + 2 \leq n \leq 2l^2 + 3l + 2 = \binom{2l+2}{2} + 1.$$

Furthermore,  $l = \lceil \frac{1}{4}(-3 + \sqrt{8n-7}) \rceil$ .

**Theorem 2.3.** Let  $n \geq 3$  and let  $l = \lceil \frac{1}{4}(-3 + \sqrt{8n-7}) \rceil$ . Then

$$\det(P_n) = \begin{cases} 2l & \text{if } 2l^2 - l + 2 \leq n \leq 2l^2 + 3, \\ 2l + 1 & \text{if } 2l^2 + 4 \leq n \leq 2l^2 + 3l + 2. \end{cases}$$

*Proof.* Observe that  $2l^2 - l + 2 \leq n \leq 2l^2 + 3l + 2$  by Lemma 2.2. It is easy to see that  $\det(P_n) = 2$  for  $3 \leq n \leq 5$  and so the result holds for  $3 \leq n \leq 5$ . Hence, we may restrict our attention to  $n \geq 6$ . We consider two cases, according to whether  $2l^2 - l + 2 \leq n \leq 2l^2 + 3$  or  $2l^2 + 4 \leq n \leq 2l^2 + 3l + 2$ .

*Case 1:*  $2l^2 - l + 2 \leq n \leq 2l^2 + 3$ . By Corollary 2.1, if  $n \geq \binom{2l}{2} + 3 = 2l^2 - l + 3$ , then  $\det(P_n) \geq 2l$ . We now show that if  $n = 2l^2 - l + 2$ , then  $\det(P_n) \geq 2l$ . Since  $n > \binom{2l-1}{2} + 2 = 2l^2 - 3l + 3$  for every  $l \geq 1$ , it follows that  $\det(P_n) \geq 2l - 1$ . Suppose that there exist a detectable  $(2l-1)$ -coloring  $c$  of  $P_n$  where  $n = 2l^2 - l + 2$ . Since the maximum number of vertices in a path with detection number  $2l-1$  is  $\binom{(2l-1)+1}{2} + 2 = \binom{2l}{2} + 2 = 2l^2 - l + 2$ , it follows that all possible color codes for the vertices of degree 2 are used in the coloring  $c$ . Observe that among the possible color codes for vertices of degree 2, there is a total of  $2l-2$  codes starting with 1. Indeed, among the codes containing exactly two 1's, there is a total of  $2l-2$  codes having 1 in the  $j$ th position for every  $j = 1, 2, \dots, 2l-1$ . Since the code of each end-vertex of  $P_n = P_{2l^2-l+2}$  contains exactly one 1, it follows that in the corresponding detectable  $(2l-1)$ -tuple factorization of  $P_n = P_{2l^2-l+2}$ , two of the factors have an odd number of vertices of degree 1, which is not possible. Hence,  $\det(P_n) = \det(P_{2l^2-l+2}) \neq 2l-1$ . Consequently,  $\det(P_n) = \det(P_{2l^2-l+2}) \geq 2l$ . This shows that  $\det(P_n) \geq 2l$  if  $2l^2 - l + 2 \leq n \leq 2l^2 + 3$ .

We now show that  $\det(P_n) \leq 2l$  if  $2l^2 - l + 2 \leq n \leq 2l^2 + 3$  by considering two subcases, depending on whether  $n = 2l^2 + 3$  or  $2l^2 - l + 2 \leq n \leq 2l^2 + 2$ .

*Subcase 1.1:*  $n = 2l^2 + 3$ . Let  $V(K_{2l}) = \{1, 2, \dots, 2l\}$ . We now describe a method to assign a detectable coloring of the edges of  $P_{2l^2+3}$  with the elements of  $V(K_{2l}) = \{1, 2, \dots, 2l\}$ . By Theorem D, there exists a factorization of  $K_{2l}$  into  $l-1$  Hamiltonian cycles

$$H_1, H_2, \dots, H_{l-1}$$

and a 1-factor  $F$ . For each integer  $i$  with  $1 \leq i \leq l-1$ , suppose that

$$H_i: 1 = a_{i,1}, a_{i,2}, \dots, a_{i,2l}, 1,$$

where  $a_{i,j}$  ( $1 \leq j \leq 2l$ ) is the  $j$ th vertex of  $H_i$ . We may assume, without loss of generality, that

$$H_1: 1, 2, \dots, 2l, 1.$$

Therefore,  $a_{1,j} = j$  for  $1 \leq j \leq 2l$ . Also, let  $b_1$  be the neighbor of 1 in the 1-factor  $F$  of  $K_{2l}$ . Note that  $b_1 \neq a_{i,2}$  and  $b_1 \neq a_{i,2l}$  for every  $i$  with  $1 \leq i \leq l-1$ . Suppose that the edges of  $P_{2l^2+3}$  are encountered in the order

$$e_1, e_2, \dots, e_{2l^2+2}$$

as we proceed along the path. For each integer  $k$  with  $1 \leq k \leq 2l^2$ , either  $1 \leq k \leq 4l$  or  $k = i(2l) + j$  for some integers  $i$  and  $j$  with  $2 \leq i \leq l-1$  and  $1 \leq j \leq 2l$ . We now define a coloring  $c: E(P_{2l^2+3}) \rightarrow V(K_{2l})$  of the edges of  $P_{2l^2+3}$  by

$$c(e_k) = \begin{cases} a_{1, \lceil k/2 \rceil} = \lceil k/2 \rceil & \text{if } 1 \leq k \leq 4l, \\ a_{i,j} & \text{if } k = i(2l) + j, 2 \leq i \leq l-1, 1 \leq j \leq 2l, \\ 1 & \text{if } k = 2l^2 + 1, \\ b_1 & \text{if } k = 2l^2 + 2. \end{cases}$$

In other words, we assign the color  $\lceil k/2 \rceil$  to the edge  $e_k$  for  $1 \leq k \leq 4l$ , color the next  $2l$  edges  $e_{2(2l)+j}$  ( $1 \leq j \leq 2l$ ) of  $P_{2l^2+3}$  by  $a_{2,j}$ , color the next  $2l$  edges  $e_{3(2l)+j}$  ( $1 \leq j \leq 2l$ ) by  $a_{3,j}$  and so on. We continue this process until we have gone through all the Hamiltonian cycles  $H_1, H_2, \dots, H_{l-1}$ . We have now assigned colors to the first  $2l^2$  edges of  $P_{2l^2+3}$ . We assign the colors 1 and  $b_1$  to the last two edges in that order. (Figure 2 illustrates a detectable  $2l$ -coloring for  $P_{2l^2+3} = P_{21}$  for  $l = 3$ .) Since every vertex of degree 2 of  $P_{2l^2+3}$  is incident with two edges having a unique pair of colors and the edges incident with the end-vertices are colored 1 and  $b_1 (\neq 1)$ ,  $c$  is a detectable  $2l$ -coloring of  $P_{2l^2+3}$  and so  $\det(P_{2l^2+3}) \leq 2l$ .




Figure 2. The detectable coloring of  $P_{21}$  in Subcase 1.1.

Subcase 1.2:  $n = 2l^2 + 3 - p$ , for some integer  $p$  with  $1 \leq p \leq l + 1$ . For each integer  $q$  with  $1 \leq q \leq p$ , let  $v_q$  be the vertex incident with  $e_{2q-1}$  and  $e_{2q}$  on  $P_{2l^2+3}$ . Suppressing the vertex  $v_q$  ( $1 \leq q \leq p$ ) so that  $e_{2q-1}$  and  $e_{2q}$  become the single edge  $f_q$ , we obtain a path  $P_{2l^2+3-p}$  of order  $2l^2 + 3 - p$ . Let  $c$  be the detectable  $2l$ -coloring of  $P_{2l^2+3}$  defined in Subcase 1.1. Define an edge coloring  $c^* : E(P_{2l^2+3-p}) \rightarrow V(K_{2l})$  of  $P_{2l^2+3-p}$  by

$$c^*(e) = \begin{cases} c(e_{2q-1}) & \text{if } e = f_q \text{ for some } q \text{ with } 1 \leq q \leq p, \\ c(e) & \text{otherwise.} \end{cases}$$

The codes of the vertices of  $P_{2l^2+3-p}$  are all those of  $P_{2l^2+3}$  except those  $p$   $2l$ -tuples for which 2 occurs in the  $q$ th coordinate for  $1 \leq q \leq p$ . This is a detectable  $2l$ -coloring of  $P_{2l^2+3-p}$  and so  $\det(P_{2l^2+3-p}) \leq 2l$ . Figure 3 illustrates a detectable  $2l$ -coloring of  $P_{2l^2+3-p} = P_{17}$  for  $l = 3$  and  $p = 4 = l + 1$ .




Figure 3. The detectable coloring of  $P_{17}$  in Subcase 1.2.

Case 2:  $2l^2 + 4 \leq n \leq 2l^2 + 3l + 2$ . By Corollary 2.1, if  $n > \binom{2l+1}{2} + 2 = 2l^2 + l + 2$ , then  $\det(P_n) \geq 2l + 1$ . Thus, if  $n \geq 2l^2 + l + 3$ , then  $\det(P_n) \geq 2l + 1$ . Now, let  $n = 2l^2 + l + 2$ . Since  $n > \binom{2l}{2} = 2l^2 - l + 2$ , it follows that  $\det(P_n) = \det(P_{2l^2+l+2}) \geq 2l$ . Suppose now that there exists a detectable  $2l$ -coloring  $c$  of  $P_n = P_{2l^2+l+2}$ . Because the largest possible number of vertices in a path with detection number  $2l$  is  $\binom{2l+1}{2} + 2 = 2l^2 + l + 2$ , all possible color codes for the vertices of degree 2 are used in the coloring  $c$ . Observe that among the codes containing exactly two 1's, there is a total of  $2l - 1$  codes having 1 in the  $j$ th position for every  $j = 1, 2, \dots, 2l$ . The code of each end-vertex of  $P_n = P_{2l^2+l+2}$  contains exactly one 1. This implies that in the corresponding detectable  $2l$ -tuple factorization of  $P_n = P_{2l^2+l+2}$ , all but two of the factors have an odd number of vertices of degree 1, which is not possible. Hence,  $\det(P_n) = \det(P_{2l^2+l+2}) \geq 2l + 1$ . Suppose now that  $1 \leq p \leq l - 2$ . Then  $2l^2 + l + 2 - p \geq 2l^2 + l + 2 - (l - 2) = 2l^2 + 4$ . But  $2l^2 + 4 > 2l^2 - l + 2$ . It follows that  $\det(P_{2l^2+l+2-p}) \geq 2l$  for every  $p = 1, 2, \dots, l - 2$ . If  $c$  is a detectable  $2l$ -coloring of  $P_{2l^2+l+2-p}$ , then in the corresponding  $2l$ -tuple factorization of  $P_{2l^2+l+2-p}$ , there would be at least  $2l - (2 + 2(l - 2)) = 2$  factors having an odd number of vertices of degree 1 which is not possible. It follows then that  $\det(P_n) \geq 2l + 1$  if  $n \geq 2l^2 + 4$ . Since  $2l^2 + 3l + 2 > 2l^2 + 4$  for all  $l \geq 1$ , we have  $\det(P_n) \geq 2l + 1$  if  $2l^2 + 4 \leq n \leq 2l^2 + 3l + 2$ .

It remains to show that  $\det(P_n) \leq 2l + 1$  whenever  $2l^2 + 4 \leq n \leq 2l^2 + 3l + 2$ . This is accomplished by finding a detectable  $(2l + 1)$ -coloring of  $P_n$ . We consider two subcases.

**Subcase 2.1.**  $n = 2l^2 + 3l + 2$ . Let  $V(K_{2l+1}) = \{1, 2, \dots, 2l, 2l + 1\}$ . We now describe a method to assign a detectable coloring of  $P_{2l^2+3l+2}$  with the elements of  $V(K_{2l+1})$ . By Theorem E,  $K_{2l+1}$  can be factored into  $l$  Hamiltonian cycles

$$H_1, H_2, \dots, H_l.$$

For each integer  $i$  with  $1 \leq i \leq l$ , suppose that

$$H_i: 1 = a_{i,1}, a_{i,2}, \dots, a_{i,2l+1}, 1,$$

where  $a_{i,j}$  ( $1 \leq j \leq 2l + 1$ ) is the  $j$ th vertex of  $H_i$ . We may assume, without loss of generality, that

$$H_1: 1, 2, \dots, 2l + 1, 1.$$

Therefore,  $a_{1,j} = j$  for  $1 \leq j \leq 2l + 1$ . Suppose that the edges of  $P_{2l^2+3l+2}$  are encountered in the order

$$e_1, e_2, \dots, e_{2l^2+3l+1}$$

as we proceed along the path. We now define a coloring  $c: E(P_{2l^2+3l+2}) \rightarrow V(K_{2l+1})$  of the edges of  $P_{2l^2+3l+2}$  by

$$c(e_k) = \begin{cases} a_{1, \lceil k/2 \rceil} = \lceil k/2 \rceil & \text{if } 1 \leq k \leq 4l + 2, \\ a_{i,j} & \text{if } k = i(2l + 1) + j, 2 \leq i \leq l, 1 \leq j \leq 2l + 1. \end{cases}$$

That is, we color the first  $4l + 2$  edges  $e_k$  ( $1 \leq k \leq 4l + 2$ ) of  $P_{2l^2+3l+2}$  by  $\lceil k/2 \rceil$ , color the next  $2l + 1$  edges  $e_{2(2l+1)+j}$  ( $1 \leq j \leq 2l + 1$ ) of  $P_{2l^2+3l+2}$  by  $a_{2,j}$ , color the next  $2l + 1$  edges  $e_{3(2l+1)+j}$  ( $1 \leq j \leq 2l + 1$ ) by  $a_{3,j}$  and so on. (Figure 4 illustrates the detectable  $(2l + 1)$ -coloring of  $P_{2l^2+3l+2} = P_{29}$  for  $l = 3$ .) The last  $2l + 1$  edges  $e_{l(2l+1)+j}$  ( $1 \leq j \leq 2l + 1$ ) are then colored by  $a_{l,j}$ . Since every vertex of degree 2 of  $P_{2l^2+3l+2}$  is incident with two edges having a unique pair of colors and the edges incident with the two end-vertices are assigned 1 and  $a_{l,2l+1} \neq 1$ , it follows that  $c$  is a detectable  $(2l + 1)$ -coloring of  $P_{2l^2+3l+2}$  and so  $\det(P_{2l^2+3l+2}) \leq 2l + 1$ . Hence,  $\det(P_{2l^2+3l+2}) = 2l + 1$ .

**Subcase 2.2:**  $2l^2 + 4 \leq n \leq 2l^2 + 3l + 1$ . Let  $n = (2l^2 + 3l + 2) - p$ , where  $1 \leq p \leq 3l - 2$ . We consider two subcases, according to whether  $1 \leq p \leq 2l + 1$  or  $2l + 2 \leq p \leq 3l - 2$ .

**Subcase 2.2.1:**  $1 \leq p \leq 2l + 1$ . For each integer  $q$  with  $1 \leq q \leq p$ , let  $v_q$  be the vertex incident with  $e_{2q-1}$  and  $e_{2q}$  on  $P_{2l^2+3l+2}$ . Suppressing the vertex  $v_q$  ( $1 \leq q \leq$







Figure 4. The detectable coloring of  $P_{29}$  in Subcase 2.1.

$p$ ) so that  $e_{2q-1}$  and  $e_{2q}$  become the single edge  $f_q$ , we obtain a path  $P_{2l^2+3l+2-p}$  of order  $2l^2+3l+2-p$ . Let  $c$  be the detectable  $(2l+1)$ -coloring of  $P_{2l^2+3l+2}$  defined in Subcase 2.1. Define an edge coloring  $c^* : E(P_{2l^2+3l+2-p}) \rightarrow V(K_{2l+1})$  of  $P_{2l^2+3l+2-p}$  by

$$c^*(e) = \begin{cases} c(e_{2q-1}) & \text{if } e = f_q \text{ for some } q \text{ with } 1 \leq q \leq p, \\ c(e) & \text{otherwise.} \end{cases}$$

The codes of the vertices of  $P_{2l^2+3l+2-p}$  are all those of  $P_{2l^2+3l+2}$  except those  $(2l+1)$ -tuples for which 2 occurs in the  $q$ th coordinate for  $1 \leq q \leq p$ . Since this is a detectable  $(2l+1)$ -coloring of  $P_{2l^2+3l+2-p}$ , it follows that  $\det(P_{2l^2+3l+2-p}) \leq 2l+1$ . Figure 5 illustrates the detectable  $(2l+1)$ -coloring of  $P_{2l^2+3l+2-p} = P_{25}$  when  $l = 3$  and  $p = 4$ .




Figure 5. The detectable coloring of  $P_{25}$  in Subcase 2.2.1.

**Subcase 2.2.2:**  $2l+2 \leq p \leq 3l-2$ . Note that this subcase can only occur when  $l \geq 4$ . Let  $p = (2l+1) + h$  where  $1 \leq h \leq l-3$ . Observe that  $l-3 < 2l+1$  for all positive integers  $l$ . Recall that the edges of  $P_{2l^2+3l+2}$  are encountered in the order

$$e_1, e_2, \dots, e_{2l^2+3l}, e_{2l^2+3l+1}$$

as we proceed along the path. Let  $v_i$  denote the vertex of  $P_{2l^2+3l+2}$  incident with  $e_i$  and  $e_{i+1}$  for  $1 \leq i \leq 6l+3$ . First, we construct a path  $P_{(2l^2+3l+2)-(2l+1)}$  from  $P_{2l^2+3l+2}$  by

- (1) deleting the vertices  $v_{4l+3}, v_{4l+4}, \dots, v_{6l+2}$  and therefore, deleting the  $2l + 1$  edges  $e_{4l+3}, e_{4l+4}, \dots, e_{6l+3}$  (which correspond to the Hamiltonian cycle  $H_2$ ), and
- (2) identifying the vertices  $v_{4l+2}$  and  $v_{6l+3}$ .

This produces a path  $P_{(2l^2+3l+2)-(2l+1)}$  of order  $(2l^2 + 3l + 2) - (2l + 1)$ . Next, we suppress the vertex  $v_{2j-1}$  for  $1 \leq j \leq h$ , where the two edges  $e_{2j-1}$  and  $e_{2j}$  become the single edge  $f_j$ . This produces a path  $P_{(2l^2+l+2)-(2l+1+h)} = P_n$ . Let  $c$  be the detectable  $(2l + 1)$ -coloring of  $P_{2l^2+3l+2}$  defined in Subcase 2.1. Define an edge coloring  $c': E(C_n) \rightarrow V(K_{2l+1})$  by

$$c'(e) = \begin{cases} c(e_{2j-1}) & \text{if } e = f_j \text{ for } 1 \leq j \leq h, \\ c(e) & \text{otherwise.} \end{cases}$$

Figure 6 illustrates a detectable  $(2l + 1)$ -coloring of  $P_{2l^2+3l+2} = P_{46}$  where  $l = 4$  and a detectable  $(2l + 1)$ -coloring of  $P_{(2l^2+3l+2)-(2l+1+h)} = P_{36}$  (for  $l = 4$  and  $h = 1$ ) obtained from the coloring of  $P_{46}$ .




Figure 6. Detectable colorings of  $P_{46}$  and  $P_{36}$  in Subcase 2.2.2 in the proof of Theorem 2.3.

The codes of the vertices of  $P_n$  are all these of  $P_{2l^2+3l+2}$  except

- (a) those  $(2l + 1)$ -tuples for which 2 occurs in the  $j$ th coordinate for  $1 \leq j \leq h$  (there are  $h$  such  $(2l + 1)$ -tuples) and
- (b) those  $(2l + 1)$ -tuples that are produced from the Hamiltonian cycle  $H_2$ ; that is, the codes of the vertices  $v_{4l+3}, v_{4l+4}, \dots, v_{6l+3}$  in the path  $P_{2l^2+3l+2}$  (there are  $2l + 1$  such  $(2l + 1)$ -tuples).

Since  $c'$  is a detectable  $(2l + 1)$ -coloring of  $P_n$ , it follows that  $\det(P_n) \leq 2l + 1$ .  $\square$

### 3. EXTREMAL PROBLEMS ON UNICYCLIC GRAPHS

A connected graph with exactly one cycle is called a *unicyclic graph*. A graph  $G$  of order  $n$  and size  $m$  is unicyclic if and only if  $G$  is connected and  $m = n$ . In this section, we study some extremal problems concerning detection numbers of unicyclic graphs, in particular, the problems of determining how large and how small the detection number of a unicyclic graph of a fixed order can be.

Observe that if  $n_i$  is the number of vertices of degree  $i$  in a unicyclic graph  $G$  with maximum degree  $\Delta$ , then

$$(1) \quad n_1 = n_3 + 2n_4 + 3n_5 + \dots + (\Delta - 2)n_\Delta.$$

For each integer  $n \geq 3$ , let  $D_u(n)$  denote the maximum detection number among all unicyclic graphs of order  $n$  and  $d_u(n)$  the minimum detection number among all unicyclic graphs of order  $n$ . That is, if  $\mathcal{U}_n$  is the set of all unicyclic graphs of order  $n$ , then

$$D_u(n) = \max\{\det(G) : G \in \mathcal{U}_n\}$$

$$d_u(n) = \min\{\det(G) : G \in \mathcal{U}_n\}.$$

Figure 7 shows all the unicyclic graphs of order  $n$  for  $3 \leq n \leq 5$  together with a minimum detectable coloring for each. Hence  $D_u(3) = d_u(3) = 3$ , and  $D_u(n) = 3$  and  $d_u(n) = 2$  for  $n = 4, 5$ .




Figure 7. Minimum detectable colorings of unicyclic graphs of order  $n = 3, 4, 5$ .

In order to determine  $D_u(n)$  for  $n \geq 6$ , we first present a lemma. For a graph  $F$ , let  $m(F)$  denote the size of  $F$ .

**Lemma 3.1.** *Let  $G$  be connected graph of order  $n \geq 3$ . If  $H$  is a connected subgraph of  $G$ , then*

$$\det(G) - \det(H) \leq m(G) - m(H).$$

*Proof.* Color the  $m(H)$  edges of  $H$  using  $k = \det(H)$  colors and color the remaining  $m(G) - m(H)$  edges of  $G$  using the colors  $k+1, k+2, \dots, k+(m(G) - m(H))$ . This gives us a detectable  $(m(G) - m(H) + k)$ -coloring of  $G$ . It follows that  $\det(G) \leq m(G) - m(H) + \det(H)$ .  $\square$

The following is an immediate consequence of Lemma 3.1

**Corollary 3.2.** *Let  $G$  be a connected graph of order  $n \geq 3$  and size  $m$ . If  $g$  is the girth of  $G$ , then*

$$\det(G) \leq m - g + \det(C_g).$$

**Proposition 3.3.** *For  $n \geq 6$ ,  $D_u(n) = n - 3$ .*

*Proof.* It is easy to verify that  $\det(K_{1,n-1} + e) = n - 3$  for  $n \geq 6$  and so  $D_u(n) \geq n - 3$  for  $n \geq 6$ . It remains to show that  $D_u(n) \leq n - 3$  for  $n \geq 6$ . Let  $G$  be a unicyclic graph of order  $n \geq 6$  and let  $g$  be the girth of  $G$ . If  $3 \leq g \leq 5$ , then  $G$  contains a subgraph  $F$  such that  $F \in \{F_1, F_2, F_3, F_4, F_5\}$ , where  $F_i$  ( $1 \leq i \leq 4$ ) is shown in Figure 7 and  $F_5$  is the graph obtained from  $C_5$ :  $v_1, v_2, v_3, v_4, v_5, v_1$  by adding a pendant edge  $vv_1$ . We have seen that  $\det(F_i) = m(F_i) - 3$  for  $1 \leq i \leq 4$ . For the graph  $F_5$ , the 3-coloring  $c$  defined by  $c(v_1v_2) = 1$ ,  $c(v_2v_3) = c(v_3v_4) = 2$ , and  $c(v_4v_5) = c(v_1v_5) = c(vv_1) = 3$  is a minimum detectable coloring of  $F_5$  and so  $\det(F_5) = 3 = m(F_5) - 3$ . Therefore,  $\det(F) = m(F) - 3$  for each  $F \in \{F_1, F_2, F_3, F_4, F_5\}$ . It then follows by Lemma 3.1 that  $\det(G) \leq m(G) + \det(F) - m(F) = n + (m(F) - 3) - m(F) = n - 3$  for  $3 \leq g \leq 5$ . If  $g \geq 6$ , then  $\det(C_g) \leq g - 3$  by Theorem C. It then follows by Corollary 3.2 that  $\det(G) \leq n - g + \det(C_g) \leq n - g + (g - 3) = n - 3$ . Thus,  $D_u(n) \leq n - 3$  for all  $n \geq 6$ .  $\square$

Next, we determine the minimum detection number among all unicyclic graphs of order  $n$ . According to Theorem B, every unicyclic graph of order  $n \geq 3$  having detection number  $k$  contains at most  $k$  end-vertices and at most  $\frac{1}{2}k(k+1)$  vertices of degree 2. It then follows by (1) that

$$n \leq k + \frac{k(k+1)}{2} + k = \frac{k^2 + 5k}{2}.$$

Furthermore, if  $G$  is a unicyclic graph of order  $n = \frac{1}{2}(k^2 + 5k)$  with  $\det(G) = k$ , then  $G$  must contain exactly  $k$  end-vertices, exactly  $\frac{1}{2}k(k+1)$  vertices of degree 2, and

exactly  $k$  vertices of degree 3. We first determine  $d_u(n)$  for the values of  $n$  mentioned above.

**Theorem 3.4.** *Let  $k \geq 2$  be an integer. If  $n = \frac{1}{2}(k^2 + 5k)$ , then  $d_u(n) = k$ .*

**Proof.** First, we show that if  $n = \frac{1}{2}(k^2 + 5k)$ , then  $d_u(n) \geq k$ . Assume, to the contrary, that there exists a unicyclic graph  $G$  of order  $\frac{1}{2}(k^2 + 5k)$  such that  $\det(G) \leq k - 1$ . By Theorem B,  $G$  has at most  $k - 1$  end-vertices and at most  $\frac{1}{2}k(k - 1)$  vertices of degree 2. Therefore,  $G$  contains at least

$$\frac{k^2 + 5k}{2} - (k - 1) - \frac{k(k - 1)}{2} = 2k + 1$$

vertices of degree 3 or more. It then follows by (1) that  $G$  contains at least  $2k + 1$  end-vertices, which is impossible. Thus,  $d_u(n) \geq k$ .

To show that  $d_u(n) \leq k$ , we construct a unicyclic graph  $G_k$  of order  $n = \frac{1}{2}(k^2 + 5k)$  having detection number  $k$  such that  $G_k$  has exactly  $k$  end-vertices, exactly  $\frac{1}{2}(k^2 + k)$  vertices of degree 2, and exactly  $k$  vertices of degree 3. We consider two cases, according as to whether  $k$  is odd or even.

**Case 1.  $k$  is odd.** Then  $k = 2l - 1$  for some integer  $l \geq 2$ . We now construct  $G_k$ . Let

$$C_{2l^2-l}: v_1, v_2, \dots, v_{2l^2-l}, v_1$$

be a cycle of length  $2l^2 - l$  and for  $1 \leq i \leq k$ , let  $Q_i$  be a copy of  $K_2$  with  $V(Q_i) = \{u_{i,1}, u_{i,2}\}$ . Then the graph  $G_k$  is obtained from  $C_{2l^2-l}$  and  $Q_i$  ( $1 \leq i \leq k$ ) by adding the edges  $v_{2i}u_{i,1}$  ( $1 \leq i \leq k$ ). Observe that  $G_k$  is a unicyclic graph of order  $n = (2l^2 - l) + 2(2l - 1) = \frac{1}{2}(k^2 + 5k)$ .

We now define a  $k$ -coloring  $c$  for the edges of  $G_k$ . First, we color the  $2l^2 - l$  edges of  $C_{2l^2-l}$  with the elements of  $V(K_{2l-1}) = \{1, 2, \dots, 2l, 2l - 1\}$  as follows. Let  $H_1, H_2, \dots, H_{l-1}$  be  $l-1$  pairwise edge-disjoint Hamiltonian cycles of  $K_{2l-1}$ . For each integer  $i$  with  $1 \leq i \leq l - 1$ , suppose that  $H_i: 1 = a_{i,1}, a_{i,2}, \dots, a_{i,2l-1}, 1$ , where  $a_{i,j}$  ( $1 \leq j \leq 2l - 1$ ) is the  $j$ th vertex of  $H_i$  and we assume that  $H_1: 1, 2, \dots, 2l - 1, 1$ . Therefore,  $a_{1,j} = j$  for  $1 \leq j \leq 2l - 1$ . Suppose that the edges of  $C_{2l^2-l}$  are encountered in the order

$$e_1, e_2, \dots, e_{2l^2-l}, e_{2l^2-l+1} = e_1$$

as we proceed about the cycle in some direction. Then we define

$$c(e_k) = \begin{cases} a_{1, \lceil k/2 \rceil} = \lceil k/2 \rceil & \text{if } 1 \leq k \leq 4l - 2, \\ a_{i,j} & \text{if } k = i(2l - 1) + j, 2 \leq i \leq l - 1, \text{ and } 1 \leq j \leq 2l - 1. \end{cases}$$

It was shown in the proof of Theorem C that this coloring of the cycle  $C_{2l^2-l}$  is detectable. Furthermore, let  $c(v_{2i}u_{i,1}) = c(u_{i,1}u_{i,2}) = i$  for  $1 \leq i \leq k$ . Thus  $c$  uses  $k$  colors. It remains to show that  $c$  is detectable. Note that the color codes of the vertices of  $G_k$  consist of all possible color codes for vertices of degrees 1 and 2 together with all the  $k$ -tuples whose only nonzero entry is 3 occurring in the  $i$ th coordinate for  $1 \leq i \leq k$ . Since each of the color codes described above occurs exactly once,  $c$  is a detectable  $k$ -coloring for  $G_k$ . Therefore,  $\det(G_k) \leq k$  and consequently,  $\det(G_k) = k$ .

**Case 2.  $k$  is even.** Then  $k = 2l$  for some positive integer  $l$ . If  $k = 2$ , then  $n = 7$ . Since the unicyclic graph  $G_2$  of order 7 in Figure 8 has detection number 2 (as shown in that figure), the result holds for  $k = 2$ . Thus we may assume that  $k \geq 4$  and so  $l \geq 2$ .




Figure 8. A detectable 2-coloring of  $G_2$  in Case 2.

Let  $C_{2l^2} : v_1, v_2, \dots, v_{2l^2}, v_1$  be a cycle of length of  $2l^2$ . For  $1 \leq i \leq l$ , let  $Q_i$  be a copy of  $K_2$  with  $V(Q_i) = \{u_{i,1}, u_{i,2}\}$  and for  $l + 1 \leq i \leq 2l$ , let  $Q_i : u_{i,1}, u_{i,2}, u_{i,3}$  be a copy of a path of length 2. Then the graph  $G_k$  is obtained from  $C_{2l^2}$  and  $Q_i$  ( $1 \leq i \leq k$ ) by adding the edges  $v_{2i}u_{i,1}$  ( $1 \leq i \leq k$ ). Observe that  $G_k$  is a unicyclic graph of order  $n = 2l^2 + 2l + 3l = \frac{1}{2}(k^2 + 5k)$ .

We now define a  $k$ -coloring  $c$  for the edges of  $G_k$ . First, we color the  $2l^2$  edges of the cycle  $C_{2l^2}$  with the elements of  $V(K_{2l}) = \{1, 2, \dots, 2l\}$  as follows. Let  $H_1, H_2, \dots, H_{l-1}$  be  $l - 1$  pairwise edge-disjoint Hamiltonian cycles of  $K_{2l}$  and let  $F$  be the 1-factor of  $K_{2l}$  with  $E(F) = \{x_i y_i : 1 \leq i \leq l\}$ , where  $x_l = 2l = k$ . For each integer  $i$  with  $1 \leq i \leq l - 1$ , suppose that  $H_i : 1 = a_{i,1}, a_{i,2}, \dots, a_{i,2l}, 1$ , where  $a_{i,j}$  ( $1 \leq j \leq 2l$ ) is the  $j$ th vertex of  $H_i$  and  $H_1 : 1, 2, \dots, 2l, 1$ , say. Therefore,  $a_{1,j} = j$  for  $1 \leq j \leq 2l$ . Suppose that the edges of  $C_{2l^2}$  are encountered in the order

$$e_1, e_2, \dots, e_{2l^2}, e_{2l^2+1} = e_1,$$

as we proceed about the cycle in some direction. For each integer  $k$  with  $1 \leq k \leq 2l^2$ , either  $1 \leq k \leq 4l$  or  $k = i(2l) + j$  for some integers  $i$  and  $j$  with  $2 \leq i \leq l - 1$  and  $1 \leq j \leq 2l$ . We now define

$$c(e_k) = \begin{cases} a_{1, \lceil k/2 \rceil} = \lceil k/2 \rceil & \text{if } 1 \leq k \leq 4l, \\ a_{i,j} & \text{if } k = i(2l) + j, 2 \leq i \leq l - 1 \text{ and } 1 \leq j \leq 2l. \end{cases}$$

This coloring of  $C_{2l2}$  is detectable, which was shown in the proof of Theorem C. Furthermore, for  $1 \leq i \leq l$ , let  $c(v_{2i}u_{i,1}) = c(u_{i,1}u_{i,2}) = x_i$  and for  $l+1 \leq i \leq 2l$ , let  $c(v_{2i}u_{i,1}) = x_{i-l}$  and let  $c(u_{i,1}u_{i,2}) = c(u_{i,2}u_{i,3}) = y_{i-l}$ . Thus  $c$  uses  $k$  colors. It remains to show that  $c$  is detectable. Note that in the coloring  $c$  all possible color codes for vertices of degrees 1 and 2 are used exactly once. The vertices of degree 3, namely  $v_{2i}$  ( $i = 1, 2, \dots, l$ ), also have distinct color codes since  $v_{2i}$  is the only vertex whose code has an entry that is at least 2 in the  $i$ th position. Therefore,  $\det(G_k) \leq k$  and consequently,  $\det(G_k) = k$ .  $\square$

With the aid of Theorem 3.4, we are now able to establish the following.

**Theorem 3.5.** *For each integer  $k \geq 2$ , if*

$$\frac{(k-1)^2 + 5(k-1)}{2} + 1 \leq n \leq \frac{k^2 + 5k}{2}$$

then  $d_u(n) = k$ .

*Proof.* First, we show that if  $\frac{1}{2}(k^2 + 3k - 2) = \frac{1}{2}((k-1)^2 + 5(k-1)) + 1 \leq n \leq \frac{1}{2}(k^2 + 5k)$ , then  $d_u(n) \geq k$ . Assume, to the contrary, that there exists a unicyclic graph  $G$  of order  $n \geq \frac{1}{2}(k^2 + 3k - 2)$  such that  $\det(G) \leq k - 1$ . By Theorem B,  $G$  has at most  $k - 1$  end-vertices and at most  $\frac{1}{2}(k^2 - k)$  vertices of degree 2. Therefore,  $G$  contains at least  $n - (k - 1) - \frac{1}{2}(k^2 - k) \geq \frac{1}{2}(k^2 + 3k - 2) - (k - 1) - \frac{1}{2}(k^2 - k) = k$  vertices of degree 3 or more. It then follows by (1) that  $G$  has at least  $k$  end-vertices which is impossible.

We next show that  $d_u(n) \leq k$  if  $\frac{1}{2}(k^2 + 3k - 2) \leq n \leq \frac{1}{2}(k^2 + 5k)$ . Theorem 3.4 shows that this is true when  $n = \frac{1}{2}(k^2 + 5k)$ . Assume therefore that  $n = \frac{1}{2}(k^2 + 5k) - p$  where  $1 \leq p \leq k + 1$ . We consider two cases, according to whether  $k$  is odd or even.

*Case 1.  $k$  is odd.* Then  $k = 2l - 1$  for some integer  $l \geq 2$ . There are two subcases, depending on whether  $1 \leq p \leq k$  or  $p = k + 1$ .

*Subcase 1.1.  $1 \leq p \leq k$ .* Construct the unicyclic graph  $G$  of order  $\frac{1}{2}(k^2 + 5k) - p$  from the unicyclic graph  $G_k$  described in Theorem 3.4 by suppressing the vertices  $u_{i,1}$  so that the edges  $v_{2i}u_{i,1}$  and  $u_{i,1}u_{i,2}$  become the single edge  $v_{2i}u_{i,2}$  where  $1 \leq i \leq p$ . Define a  $k$ -coloring  $c^*$  of  $G$  by

$$c^*(e) = \begin{cases} c(v_{2i}u_{i,2}) & \text{if } e = v_{2i}u_{i,2} \text{ for some } i \text{ with } 1 \leq i \leq p, \\ c(e) & \text{otherwise.} \end{cases}$$

The color codes for the vertices of  $G$  are all those of  $G_k$  except those  $(2l - 1)$ -tuples for which 2 occurs in the  $i$ th coordinate (and 0 occurs everywhere else) for  $1 \leq i \leq p$ .

Thus  $c^*$  is a detectable  $k$ -coloring of  $G$ . Since  $G$  has  $k$  end-vertices, it follows that  $\det(G) = k$ . Consequently,  $d_u(n) \leq k$ .

**Subcase 1.2.**  $p = k + 1$  and so  $n = \frac{1}{2}(k^2 + 3k - 2)$ . Consider the unicyclic graph  $G$  of order  $\frac{1}{2}(k^2 + 3k - 2) + 1 = \frac{1}{2}(k^2 + 3k)$  together with the edge coloring described in Subcase 1.1 (that is, when  $p = k$  in Subcase 1.1.). We delete the edge  $v_{2k}v_{2k-1}$ , identify the vertices  $v_{2k}$  and  $v_{2k-1}$ , and label this new vertex by  $v$ . This gives us a unicyclic graph  $G'$  of order  $n = \frac{1}{2}(k^2 + 3k - 2)$ . Observe that the color codes of the vertices of  $G'$  are those of  $G$  except for those of  $v_{2k}$  and  $v_{2k-1}$ , and that  $v$  is the only vertex of  $G'$  of degree 3 whose color code has 2 as the  $k$ th coordinate. Hence, we have a detectable  $k$ -coloring of  $G'$ . Since  $G'$  has  $k$  end-vertices, it follows that  $\det(G') \geq k$  and consequently,  $\det(G') = k$ . Therefore,  $d_u(n) \leq k$ .

**Case 2.**  $k$  is even. Then  $k = 2l$  for some positive integer  $l$ . The result holds for  $k = 2$  (that is,  $l = 1$ ) as the graphs in Figure 9 show. For  $k \geq 4$  (and so  $l \geq 2$ ), we consider three subcases, according to whether  $1 \leq p \leq k/2$ ,  $k/2 + 1 \leq p \leq k$ , or  $p = k + 1$ .




Figure 9. Detectable colorings when  $k = 2$  in Case 2.

**Subcase 2.1.**  $1 \leq p \leq k/2$ . Construct the unicyclic graph  $G$  of order  $\frac{1}{2}(k^2 + 5k) - p$  from the unicyclic graph  $G_k$  described in Theorem 3.4 by suppressing the vertices  $u_{i,1}$  so that the edges  $v_{2i}u_{i,1}$  and  $u_{i,1}u_{i,2}$  become the single edge  $v_{2i}u_{i,2}$  where  $1 \leq i \leq p$ . Define a  $k$ -coloring  $c^*$  of  $G$  by

$$c^*(e) = \begin{cases} c(v_{2i}u_{i,2}) & \text{if } e = v_{2i}u_{i,2} \text{ for some } i \text{ with } 1 \leq i \leq p, \\ c(e) & \text{otherwise.} \end{cases}$$

The color codes for the vertices of  $G$  are all those of  $G_k$  except those  $(2l)$ -tuples for which 2 occurs in the  $x_i$ th coordinate (and 0 occurs everywhere else) for  $1 \leq i \leq p$ . Thus  $c^*$  is a detectable  $k$ -coloring of  $G$ . Since  $G$  has  $k$  end-vertices, it follows that  $\det(G) \leq k$ . Consequently,  $d_u(n) \leq k$ .

**Subcase 2.2.**  $k/2 + 1 \leq p \leq k$ . Construct the unicyclic graph  $G'$  of order  $\frac{1}{2}(k^2 + 5k) - p$  from the unicyclic graph  $G$  of order  $\frac{1}{2}(k^2 + 5k) - k/2 = \frac{1}{2}(k^2 + 4k)$  described in Subcase 2.1 (that is, when  $p = k/2$  in Subcase 2.1) by suppressing the vertices  $u_{i,2}$  so that the edges  $u_{i,1}u_{i,2}$  and  $u_{i,2}u_{i,3}$  become the single edge  $u_{i,1}u_{i,3}$



where  $k/2 + 1 \leq i \leq p$ . Define a  $k$ -coloring  $c'$  for  $G'$  by

$$c'(e) = \begin{cases} c^*(u_{i,2}u_{i,3}) & \text{if } e = u_{i,1}u_{i,3} \text{ for some } i \text{ with } k/2 + 1 \leq i \leq p, \\ c^*(e) & \text{otherwise.} \end{cases}$$

The color codes for the vertices of  $G'$  are all those of  $G$  except those  $(2l)$ -tuples for which 2 occurs in the  $y_{i-k/2}$ th coordinate (and 0 occurs everywhere else) for  $k/2 + 1 \leq i \leq p$ . Thus  $c'$  is a detectable  $k$ -coloring of  $G'$ . Since  $G'$  has  $k$  end-vertices, it follows that  $\det(G') = k$ . Consequently,  $d_u(n) \leq k$ .

**Subcase 2.3.**  $p = k + 1$ . That is,  $n = \frac{1}{2}(k^2 + 3k - 2)$ . Consider the unicyclic graph  $G'$  of order  $\frac{1}{2}(k^2 + 3k - 2) + 1 = \frac{1}{2}(k^2 + 3k)$  together with the edge coloring described in Subcase 2.2 (that is, when  $p = k$  in Subcase 2.2). We delete the edge  $v_{2k}v_{2k-1}$ , identify the vertices  $v_{2k}$  and  $v_{2k-1}$ , and label this new vertex by  $v$ . This gives us a unicyclic graph  $G''$  of order  $n = \frac{1}{2}(k^2 + 3k - 2)$ . Observe that the color codes of the vertices of  $G''$  are those of  $G'$  except for those of  $v_{2k}$  and  $v_{2k-1}$ , and that  $v$  is the only vertex of  $G''$  of degree 3 whose color code has 2 as the  $k$ th coordinate. Hence, we have a detectable  $k$ -coloring of  $G''$ . Since  $G''$  has  $k$  end-vertices, it follows that  $\det(G'') = k$ . Consequently,  $d_u(n) \leq k$ .  $\square$

Solving for the smallest integer  $k$  for which  $n \leq \frac{1}{2}(k^2 + 5k)$ , we obtain the following.

**Theorem 3.6.** For each integer  $n \geq 4$ ,

$$d_u(n) = \left\lceil \frac{-5 + \sqrt{8n + 25}}{2} \right\rceil.$$

By Theorem 3.6,  $d_u(n) \approx \sqrt{2n}$  for large values of  $n$ . We now determine all pairs  $k, n$  of integers for which there exists a unicyclic graph of order  $n$  having detection number  $k$ .

**Theorem 3.7.** Let  $k \geq 2$  and  $n \geq 3$  be integers. There exists a unicyclic graph  $G$  of order  $n$  such that  $\det(G) = k$  if and only if  $d_u(n) \leq k \leq D_u(n)$ .

**Proof.** By definition, if  $G$  is a unicyclic graph of order  $n$  such that  $\det(G) = k$ , then  $d_u(n) \leq k \leq D_u(n)$ . It remains to verify the converse. The result holds for  $3 \leq n \leq 5$  as the graphs in Figure 7 show. Furthermore, the graphs in Figure 10 show that the result holds for  $n = 6, 7$  as well.

We now assume that  $n \geq 8$  and so  $k \geq 3$ . In this case, we show that if

$$d_u(n) = \left\lceil \frac{-5 + \sqrt{8n + 25}}{2} \right\rceil \leq k \leq n - 3 = D_u(n),$$




Figure 10. Minimum detectable colorings for graphs of order  $n = 6, 7$ .

then there is a unicyclic graph  $G$  of order  $n$  such that  $\det(G) = k$ . For each integer  $i$  with  $i = 0, 1, \dots, n - d_u(n) - 3$ , we construct a unicyclic graph  $H_i$  such that  $H_i$  has order  $n$  and  $\det(H_i) = d_u(n) + i$ . Let  $H$  be the unicyclic graph of order  $n$  described in the proof of Theorem 3.5 and  $c$  the  $d_u(n)$ -coloring described in the proof of Theorem 3.5 as well. We first construct a unicyclic graph  $H_0$  from  $H$  as follows.

- (a) If vertex  $u_{1,1} \in V(H)$ , then delete the vertex  $u_{1,2}$ ; while if  $u_{1,1} \notin V(H)$ , then delete the vertex  $v_3$  and join the vertices  $v_2$  and  $v_4$ .
- (b) Delete the edge  $v_1v_2$ , add the vertex  $v$ , and join  $v$  to  $v_1$  and  $v_2$ .

Then  $H_0$  has exactly  $d_u(n)$  end-vertices and so  $\det(H_0) \geq d_u(n)$ . Define the coloring  $c_0: E(H_0) \rightarrow \{1, 2, \dots, d_u(n)\}$  by

$$c_0(e) = \begin{cases} c(e) & \text{if } e \in E(H), \\ 1 & \text{if } e \notin E(H). \end{cases}$$

Then  $c_0$  is a detectable  $d_u(n)$ -coloring of  $H_0$ . Thus  $\det(H_0) \leq d_u(n)$  and so  $\det(H_0) = d_u(n)$ .

Observe that if  $l = \lceil d_u(n)/2 \rceil$ , then the girth of  $H_0$  is  $2l^2 - l$ ,  $2l^2$ ,  $2l^2 - l + 1$ , or  $2l^2 + 1$ , depending on (1) the parity of  $d_u(n)$  and (2) whether the vertex  $u_{1,1}$  is in  $H$  or not. In each case, if we denote the girth of  $H_0$  by  $g(l)$ , then  $g(l) > 3$  and so  $H_0 \neq K_{1,n-1} + e$ . Note that the vertices  $v_1, v$  and  $v_2$ , in this order, are consecutive vertices in the cycle of  $H_0$ . For the purpose of notation, we relabel the vertex  $v_2$  as  $w_0$ . Since  $H_0 \neq K_{1,n-1} + e$  (as  $g(l) > 3$ ), it follows that there exists a vertex  $x_0$  in  $H_0$  adjacent to  $w_0$  such that  $\deg_{H_0} x_0 \neq 1$  and  $x_0 \notin \{v, v_1\}$ .

We now construct a unicyclic graph  $H_1$  from  $H_0$  by deleting the edge  $w_0x_0$ , identifying the vertices  $w_0$  and  $x_0$ , labeling the new vertex by  $w_1$ , introducing a new vertex  $y_1$ , and joining  $y_1$  to  $w_1$ . We note that  $H_1$  has order  $n$  and has  $d_u(n) + 1$  end-vertices. Thus,  $\det(H_1) \geq d_u(n) + 1$ . To show that  $\det(H_1) \leq d_u(n) + 1$ , we provide

a detectable  $(d_u(n) + 1)$ -coloring of  $H_1$ . Define  $c_1: E(H_1) \rightarrow \{1, 2, \dots, d_u(n) + 1\}$  by

$$c_1(e) = \begin{cases} c_0(e) & \text{if } e \in E(H_0), \\ d_u(n) + 1 & \text{if } e = w_1y_1. \end{cases}$$

Then  $c_1$  is detectable  $(d_u(n) + 1)$ -coloring of  $H_1$ . This implies that  $\det(H_1) = d_u(n) + 1$ .

In general, we construct  $H_{i+1}$  from  $H_i$  and obtain the edge coloring  $c_{i+1}$  from  $c_i$ , where  $0 \leq i \leq n - d_u(n) - 4$ , as follows:

- (1) Let  $x_i$  be a vertex in  $H_i$  that is adjacent to  $w_i$  such that  $\deg_{H_i} x_i \neq 1$  and  $x_i \notin \{v, v_1\}$ .
- (2) Construct  $H_{i+1}$  by deleting the edge  $w_ix_i$ , identifying the vertices  $w_i$  and  $x_i$ , labeling the new vertex by  $w_{i+1}$ , introducing a new vertex  $y_{i+1}$ , and joining  $y_{i+1}$  to  $w_{i+1}$ .
- (3) Define  $c_{i+1}: E(H_{i+1}) \rightarrow \{1, 2, \dots, d_u(n) + i + 1\}$  by

$$c_{i+1}(e) = \begin{cases} c_i(e) & \text{if } e \in E(H_i), \\ d_u(n) + i + 1 & \text{if } e = w_{i+1}y_{i+1}. \end{cases}$$

Observe that for every integer  $i = 0, 1, \dots, n - d_u(n) - 3$ :

- (i)  $H_i$  is a unicyclic graph of order  $n$  with  $d_u(n) + i$  end-vertices;
- (ii)  $c_i$  is a detectable  $(d_u(n) + i)$ -coloring of  $H_i$ ;
- (iii) Parts (i) and (ii) imply that  $\det(H_i) = d_u(n) + i$ .

Figure 11 illustrates how to construct the unicyclic graphs  $H_i$  ( $0 \leq i \leq 6$ ) for  $n = 12$ . In this case,  $d_u(12) = 3$ ,  $D_u(12) = 9$ , and  $\det(H_i) = d_u(n) + i = 3 + i$  for  $0 \leq i \leq 6$ .

□




Figure 11. Constructing unicyclic graphs in the proof of Theorem 3.7 for  $n = 12$ .

### References

- [1] *M. Aigner, E. Triesch*: Irregular assignments and two problems á la Ringel. Topics in Combinatorics and Graph Theory. (R. Bodendiek and R. Henn, eds.). Physica, Heidelberg (1990), 29–36.
- [2] *M. Aigner, E. Triesch, Z. Tuza*: Irregular assignments and vertex-distinguishing edge-colorings of graphs. Combinatorics '90, Proc. Conf., Gaeta/Italy 1990, Elsevier Science Pub., New York (1992), 1–9.
- [3] *A. C. Burris*: On graphs with irregular coloring number 2. Congr. Numerantium 100 (1994), 129–140.
- [4] *A. C. Burris*: The irregular coloring number of a tree. Discrete Math. 141 (1995), 279–283.
- [5] *G. Chartrand, H. Escudro, F. Okamoto, P. Zhang*: Detectable colorings of graphs. To appear in Util. Math.
- [6] *G. Chartrand, P. Zhang*: Introduction to Graph Theory. McGraw-Hill, Boston, 2005.

*Authors' addresses:* Henry Escudro, Ping Zhang, Department of Mathematics, Western Michigan University, Kalamazoo, MI 49008, USA, e-mail: ping.zhang@wmich.edu.