

Jiří Anděl; Pavel Ranocha

Stationary distribution of absolute autoregression

*Kybernetika*, Vol. 41 (2005), No. 6, [735]--742

Persistent URL: <http://dml.cz/dmlcz/135689>

## Terms of use:

© Institute of Information Theory and Automation AS CR, 2005

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these

*Terms of use.*



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*  
<http://project.dml.cz>

## STATIONARY DISTRIBUTION OF ABSOLUTE AUTOREGRESSION

JIŘÍ ANDĚL AND PAVEL RANOCHA

A procedure for computation of stationary density of the absolute autoregression (AAR) model driven by white noise with symmetrical density is described. This method is used for deriving explicit formulas for stationary distribution and further characteristics of AAR models with given distribution of white noise. The cases of Gaussian, Cauchy, Laplace and discrete rectangular distribution are investigated in detail.

*Keywords:* absolute autoregression, stationary distribution, marginal distribution

*AMS Subject Classification:* 60G10

### 1. INTRODUCTION

Let  $\{X_t\}$  be an ergodic Markov process with discrete time. Its stationary distribution  $\pi$  is given by integral equation

$$\pi(A) = \int_{-\infty}^{\infty} P(A|x) d\pi(x), \quad A \in \mathcal{B} \quad (1)$$

where  $\mathcal{B}$  is the  $\sigma$ -algebra of Borel sets and  $P$  is the conditional (i. e. transition) probability. Even in the simplest linear models such as AR(1) it is not easy to find a closed form solution  $\pi$  of (1). If the model for  $\{X_t\}$  is non-linear, it is even more difficult to solve (1). One of the rare exceptions where a solution of (1) was found is so called absolute autoregression (AAR)

$$X_t = a|X_{t-1}| + \varepsilon_t \quad (2)$$

where  $a \in (-1, 1)$  and  $\varepsilon_t$  is a strict white noise (i. e., a sequence of i.i.d. random variables). Anděl et al. [4] proved that for  $a \in (-1, 0)$  and for  $\varepsilon_t \sim N(0, 1)$  the stationary density of (2) is

$$h(x) = \sqrt{\frac{2(1-a^2)}{\pi}} \exp\{-(1-a^2)x^2/2\} \Phi(ax) \quad (3)$$

where  $\Phi$  is the distribution function of  $N(0, 1)$ . It was derived that in this case

$$EX_t = \sqrt{\frac{2}{\pi}} \frac{a}{\sqrt{1-a^2}}, \quad \text{var } X_t = \frac{\pi - 2a^2}{\pi(1-a^2)} \quad (4)$$

and the correlation coefficient between  $X_t$  and  $X_{t-1}$  is

$$\rho(a) = \frac{|a|\pi + 2a^2\sqrt{1-a^2} - 2a^2 - 2|a|\arctg\sqrt{a^{-2}-1}}{\pi - 2a^2}. \tag{5}$$

Let  $C(\alpha, \beta)$  be the Cauchy distribution with the density

$$f(x) = \frac{1}{\pi} \frac{\beta}{\beta^2 + (x - \alpha)^2}.$$

Consider the model (2) with  $a \in (-1, 0)$  and  $\varepsilon_t \sim C(0, 1)$ . Define  $A = |a|/(1 - |a|)$ . Anděl and Bartoň [3] proved that  $X_t$  in (2) has the stationary density

$$h(x) = \frac{2A}{\pi^2} \left\{ \frac{(1+A)\pi}{2A[(1+A)^2 + x^2]} - \frac{x \ln[A^{-2}(1+x^2)] + (A^2 - 1 + x^2) \arctg x}{4A^2x^2 + (1 - A^2 + x^2)^2} \right\}. \tag{6}$$

Chan and Tong [5] and Tong [7, p. 141] simplified the methods used for derivation of (3) and (6). Their procedure can be summarized as follows. Let  $\varepsilon_t$  in (2) have a symmetric density  $f$ . Let  $g$  be the stationary density of the AR(1) process  $\xi_t$  given by

$$\xi_t = a\xi_{t-1} + \varepsilon_t. \tag{7}$$

Then the stationary density  $h$  of  $X_t$  in (2) is

$$h(y) = 2 \int_0^\infty g(x)f(y - ax) dx. \tag{8}$$

(The authors overlooked that the factor 2 must be introduced in the last formula.) Let us remark that if we have a guess that a function  $h$  could be a stationary density of  $X_t$  then it is easy to verify it from (2).

Problems in non-linear time series models are usually quite complicated and must be solved numerically or by simulations. It is important to have a few explicit solutions because they allow to compare accuracy of numerical methods with the exact results. One such solution for the first-order threshold autoregression with Laplace white noise has been deduced quite recently by Loges [6]. In this paper we derive some new stationary distributions of the AAR process  $\{X_t\}$  given by (2).

## 2. NORMAL DISTRIBUTION

We mentioned above that formulas (3), (4), and (5) were derived under the assumptions that  $\varepsilon_t \sim N(0, 1)$  and  $a \in (-1, 0)$ . We generalize the results to  $a \in (-1, 1)$ .

If  $\varepsilon_t \sim N(0, 1)$  then  $\xi_t$  in (7) has the distribution  $N(0, \frac{1}{1-a^2})$ . From (8) we get that the stationary density  $h$  of  $X_t$  is

$$h(y) = 2 \int_0^\infty \sqrt{\frac{1-a^2}{2\pi}} \exp\{-(1-a^2)x^2/2\} \frac{1}{\sqrt{2\pi}} \exp\{-(y-ax)^2/2\} dx.$$

Direct integration leads to formula (3) and we can see that (3) and (4) are valid for  $a \in (-1, 1)$ .

Using the same procedures as in Theorem 4.3 in Anděl et al. [4], we can derive for  $a \in (0, 1)$  that the correlation coefficient  $\rho(a)$  between  $X_t$  and  $X_{t-1}$  is also given by formula (5). This means that  $\rho(-a) = \rho(a)$ ,  $a \in (-1, 1)$ .

The density  $h$  is plotted in Figure 1 (for  $a = -0.8$ ) and in Figure 2 (for  $a = 0.8$ ). Expectation  $EX_t$  and variance  $\text{var } X_t$  as functions of  $a$  given by (4) are introduced in Figure 3 and Figure 4, respectively. In Figure 5 we can see  $\rho(a)$ , which is defined by (5).




Fig. 1. Function  $h$  for  $a = -0.8$ .




Fig. 2. Function  $h$  for  $a = 0.8$ .




Fig. 3. Expectation  $EX_t$ .




Fig. 4. Variance  $\text{var } X_t$ .

The joint stationary density of  $(X_s, X_{s-1})$  is

$$p_2(x_s, x_{s-1}) = \begin{cases} \frac{\sqrt{1-a^2}}{\pi} \exp\left\{-\frac{1-a^2}{2}x_{s-1}^2\right\} \Phi(ax_{s-1}) \exp\left\{-\frac{(x_s+ax_{s-1})^2}{2}\right\} & \text{for } x_{s-1} < 0, \\ \frac{\sqrt{1-a^2}}{\pi} \exp\left\{-\frac{1-a^2}{2}x_{s-1}^2\right\} \Phi(ax_{s-1}) \exp\left\{-\frac{(x_s-ax_{s-1})^2}{2}\right\} & \text{for } x_{s-1} > 0. \end{cases}$$

The joint stationary density of  $(X_s, X_{s-2})$  is

$$\begin{aligned}
 p_3(x_s, x_{s-2}) = & \frac{1}{\pi} \sqrt{\frac{1-a^2}{1+a^2}} \Phi(ax_{s-2}) \\
 & \times \left( \exp \left\{ -\frac{x_s^2 - 2a^2 x_s x_{s-2} + x_{s-2}^2}{2(1+a^2)} \right\} \Phi \left[ \frac{a(x_s + x_{s-2})}{\sqrt{1+a^2}} \right] \right. \\
 & \left. + \exp \left\{ -\frac{x_s^2 + 2a^2 x_s x_{s-2} + x_{s-2}^2}{2(1+a^2)} \right\} \Phi \left[ \frac{a(x_s - x_{s-2})}{\sqrt{1+a^2}} \right] \right).
 \end{aligned}$$




Fig. 5. Correlation coefficient  $\rho(a)$ .

The functions  $p_2$  and  $p_3$  for  $a = 0.8$  are introduced in Figure 6 and Figure 7, respectively.




Fig. 6. Function  $p_2(x_s, x_{s-1})$ .




Fig. 7. Function  $p_3(x_s, x_{s-2})$ .

### 3. CAUCHY DISTRIBUTION

If  $\varepsilon_t \sim C(0, 1)$  then the stationary distribution of the process  $\xi_t = a\xi_{t-1} + \varepsilon_t$  with  $|a| < 1$  is  $C(0, Q)$  where  $Q = 1/(1 - |a|)$ . The corresponding density is

$$g(x) = \frac{1}{\pi} \frac{Q}{Q^2 + x^2}.$$

The stationary density of  $X_t$  can be calculated from (8). We obtain

$$h(y) = \frac{2Q}{\pi^2} \int_0^\infty \frac{1}{Q^2 + x^2} \frac{1}{1 + (y - ax)^2} dx.$$

Again define  $A = |a|/(1 - |a|)$ . Let  $a \in (0, 1)$ . After some computations, which can be simplified using program package Mathematica, we get

$$h(y) = \frac{2A}{\pi^2} \left\{ \frac{(1 + A)\pi}{2A[(1 + A)^2 + y^2]} + \frac{y \ln[A^{-2}(1 + y^2)] + (A^2 - 1 + y^2) \operatorname{arctg} y}{4A^2y^2 + (1 - A^2 + y^2)^2} \right\}. \quad (9)$$

The density  $h$  for  $a = -0.8$  defined by (6) is plotted in Figure 8. If  $a = 0.8$  then  $h$  is defined by (9) and its graph can be found in Figure 9.




Fig. 8. Function  $h$  for  $a = -0.8$ .




Fig. 9. Function  $h$  for  $a = 0.8$ .

### 4. A DISCRETE WHITE NOISE

Assume that  $a = \frac{1}{2n}$  and

$$\varepsilon_t = \frac{2i - 1}{2n} b \text{ with probability } \frac{1}{2n}$$

for  $i = -n+1, \dots, n$  where  $b > 0$  and  $n = 1, 2, \dots$ . Then the rectangular distribution  $R(-b, b)$  is the stationary distribution of the process  $\xi_t$  in (7) (see Anděl [1, 2]). Let  $\chi_B$  be the characteristic function of the set  $B$  and let  $\delta_c(x)$  be the Dirac  $\delta$ -function, i.e.  $\int_{-\infty}^\infty \delta_c(x) dx = 1$  where

$$\delta_c(x) = \begin{cases} \infty & \text{for } x = c, \\ 0 & \text{otherwise.} \end{cases}$$

The distribution of  $\varepsilon_t$  can be described by the generalized density

$$f(x) = \frac{1}{2n} \sum_{i=-n+1}^n \delta_{\frac{2i-1}{2n}b}(x).$$

A random variable with the density  $f$  is discrete and it reaches values  $\frac{2i-1}{2n}b$ ,  $i = -n+1, \dots, n$ , each with probability  $\frac{1}{2n}$ . A straightforward calculation gives

$$h(y) = \frac{1}{b} \int_0^\infty \chi_{[-b,b]}(x) f\left(y - \frac{1}{2n}x\right) dx = \frac{1}{b} \sum_{i=-n+1}^n \chi_{[\frac{b}{n}i - \frac{b}{2n}, \frac{b}{n}i]}(y).$$

It is easy to verify that  $h$  is really the stationary density of the AAR process (2). Further we obtain

$$EX_t = \frac{b}{4n}, \quad EX_t^2 = \frac{b^2}{3}, \quad \text{var } X_t = \frac{b^2(16n^2 - 3)}{48n^2}.$$

Since

$$X_t = \frac{1}{2n}|X_{t-1}| + \varepsilon_t, \quad E\varepsilon_t = 0,$$

we have

$$EX_t X_{t-1} = \frac{1}{2n} E|X_{t-1}| X_{t-1} = -\frac{1}{2n} \int_{-\infty}^0 x^2 h(x) dx + \frac{1}{2n} \int_0^\infty x^2 h(x) dx = \frac{b^2}{8n^2}$$

and thus

$$\rho = \text{corr}(X_t, X_{t-1}) = \frac{EX_t X_{t-1} - (EX_t)^2}{\text{var } X_t} = \frac{3}{16n^2 - 3}.$$

A simple case arises for  $n = 1$  when we have the process  $X_t = \frac{1}{2}|X_{t-1}| + \varepsilon_t$  where

$$\varepsilon_t = \begin{cases} -\frac{b}{2} & \text{with probability } \frac{1}{2}, \\ \frac{b}{2} & \text{with probability } \frac{1}{2}. \end{cases}$$

The stationary density of  $X_t$  is

$$h(y) = \frac{1}{b} \left\{ \chi_{[-\frac{b}{2}, 0]}(y) + \chi_{[\frac{b}{2}, b]}(y) \right\}$$

and  $\rho = 3/13 = 0.231$ .

Now, we consider the case  $a = \frac{1}{2n+1}$  and

$$\varepsilon_t = \frac{2i}{2n+1}b \text{ with probability } \frac{1}{2n+1}$$

for  $i = -n, \dots, n$  where  $b > 0$  and  $n = 1, 2, \dots$ . The stationary distribution of the process  $\xi_t$  in (7) is again the rectangular distribution  $R(-b, b)$  (see Anděl [1, 2]). The generalized density of  $\varepsilon_t$  is

$$f(x) = \frac{1}{2n+1} \sum_{i=-n}^n \delta_{\frac{2i}{2n+1}b}(x)$$

and we obtain

$$h(y) = \frac{1}{b} \int_0^\infty \chi_{[-b,b]}(x) f\left(y - \frac{1}{2n+1}x\right) dx = \frac{1}{b} \sum_{i=-n}^n \chi_{\left[\frac{2i}{2n+1}b, \frac{2i+1}{2n+1}b\right]}(y).$$

It can be verified that  $h$  is the stationary density of the process  $\{X_t\}$  in (2). Further we obtain

$$EX_t = \frac{b}{2(2n+1)}, \quad EX_t^2 = \frac{b^2}{3}, \quad \text{var } X_t = b^2 \frac{16n^2 + 16n + 1}{12(2n+1)^2}$$

and

$$EX_t X_{t-1} = b^2 \frac{6n^2 + 6n + 1}{3(2n+1)^4}, \quad \rho = \frac{12n^2 + 12n + 1}{(2n+1)^2(16n^2 + 16n + 1)}.$$

### 5. LAPLACE DISTRIBUTION

Laplace distribution  $\text{La}(b)$  has the density

$$p(x) = \frac{1}{2b} \exp\left\{-\frac{|x|}{b}\right\}$$

where  $b > 0$  is a parameter. Assume that  $a \in (-1, 1)$  and that  $\{Z_t\}$  are i.i.d.  $\text{La}(b)$  random variables. Let the strict white noise be defined by

$$\varepsilon_t = \begin{cases} 0 & \text{with probability } a^2, \\ Z_t & \text{with probability } 1 - a^2. \end{cases}$$

Then  $\xi_t = a\xi_{t-1} + \varepsilon_t$  has the stationary density  $p(x)$  (see Anděl [1, 2]) and

$$h(y) = 2 \int_0^\infty p(x)[a^2 \delta_0(y - ax) + (1 - a^2)p(y - ax)] dx.$$

From here we obtain that for  $a \in (-1, 1)$  the stationary density of the process  $\{X_t\}$  is given by

$$h(y) = \begin{cases} \frac{1+a}{2b} \exp\left\{-\frac{y}{b}\right\} & \text{for } y > 0, \\ \frac{1-a}{2b} \exp\left\{\frac{y}{b}\right\} & \text{for } y < 0. \end{cases}$$

Further we get

$$EX_t = ab, \quad EX_t^2 = 2b^2, \quad \text{var } X_t = b^2(2 - a^2)$$

and

$$EX_t X_{t-1} = 2a^2 b^2, \quad \rho = \frac{a^2}{2 - a^2}.$$

If  $a = 0$  then  $\rho = 0$  and  $\rho \rightarrow 1$  as  $a \rightarrow \pm 1$ .



## ACKNOWLEDGMENT

The work is a part of the research project MSM 113200008 supported by the Ministry of Education, Youth and Sports of the Czech Republic. It was also partially supported by Grant 201/05/H007 of the Grant Agency of the Czech Republic.

(Received December 23, 2004.)

## REFERENCES

- 
- [1] J. Anděl: Dependent random variables with a given marginal distribution. *Acta Univ. Carolin. – Math. Phys.* 24 (1983), 3–12.
  - [2] J. Anděl: Marginal distributions of autoregressive processes. In: *Trans. 9th Prague Conference Inform. Theory, Statist. Dec. Functions, Random Processes*. Academia, Praha 1983.
  - [3] J. Anděl and T. Bartoň: A note on the threshold AR(1) model with Cauchy innovations. *J. Time Ser. Anal.* 7 (1986), 1–5.
  - [4] J. Anděl, I. Netuka, and K. Zvára: On threshold autoregressive processes. *Kybernetika* 20 (1984), 89–106.
  - [5] K. S. Chan and H. Tong: A note on certain integral equations associated with non-linear time series analysis. *Probab. Theory Related Fields* 73 (1986), 153–159.
  - [6] W. Loges: The stationary marginal distribution of a threshold AR(1) process. *J. Time Ser. Anal.* 25 (2004), 103–125.
  - [7] H. Tong: *Non-Linear Time Series*. Clarendon Press, Oxford 1990.

*Jiří Anděl and Pavel Ranocha, Department of Probability and Mathematical Statistics, Faculty of Mathematics and Physics – Charles University, Sokolovská 83, 186 75 Praha 8, Czech Republic.*

*e-mails: andel@karlin.mff.cuni.cz, ranocha@karlin.mff.cuni.cz*