

Robert Černý

Sharp generalized Trudinger inequalities via truncation for embedding into multiple exponential spaces

Commentationes Mathematicae Universitatis Carolinae, Vol. 51 (2010), No. 4, 577--593

Persistent URL: <http://dml.cz/dmlcz/140839>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2010

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Sharp generalized Trudinger inequalities via truncation for embedding into multiple exponential spaces

ROBERT ČERNÝ

Abstract. We prove that the generalized Trudinger inequality for Orlicz-Sobolev spaces embedded into multiple exponential spaces implies a version of an inequality due to Brézis and Wainger.

Keywords: Orlicz spaces, Sobolev inequalities

Classification: 46E35, 46E30

1. Introduction

Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain. The classical Sobolev embedding theorem asserts that $W_0^{1,p}(\Omega)$ is continuously embedded into $L^{p^*}(\Omega)$ if $1 \leq p < n$ and $p^* = \frac{pn}{n-p}$. Further $W_0^{1,p}(\Omega)$, $p > n$, is embedded to $L^\infty(\Omega)$. Even though p^* tends to infinity as $p \rightarrow n-$, there are unbounded functions in $W_0^{1,n}(\Omega)$.

A famous result by Trudinger [25] (see also [12], [22], [24] and [26]) states that the space $W_0^{1,n}(\Omega)$ is continuously embedded in the Orlicz space $\exp L^{\frac{n}{n-1}}(\Omega)$ (see Preliminaries for the definition of Orlicz spaces), i.e. there exist $C_1 = C_1(n)$ and $C_2 = C_2(n)$ such that

$$(1.1) \quad \int_{\Omega} \exp\left(\left(\frac{|u(x)|}{C_1 \|\nabla u\|_{L^n(\Omega)}}\right)^{\frac{n}{n-1}}\right) dx \leq C_2 \mathcal{L}_n(\Omega)$$

for every non-trivial function $u \in W_0^{1,n}(\Omega)$.

It is known (see [13], [7] and [3]) that $\exp L^{\frac{n}{n-1}}(\Omega)$ is the smallest Orlicz space with this property. However, even sharper inequalities exist in other scales. By a result of Brézis and Wainger [1] and independently Hansson [11] (see also [19] for a simple proof) we know that

$$(1.2) \quad \int_0^{\mathcal{L}_n(\Omega)} \frac{(u^*(t))^n}{\log^n\left(\frac{e\mathcal{L}_n(\Omega)}{t}\right)} \frac{dt}{t} \leq C \|\nabla u\|_{L^n(\Omega)}^n$$

for every $u \in W_0^{1,n}(\Omega)$. This inequality can be also derived from capacity estimates by Maz'ya [17]. The results in [8] and [4] tell us that this inequality gives us the smallest rearrangement invariant Banach function space $Y(\Omega)$ such

that $W_0^{1,n}(\Omega)$ is continuously embedded into $Y(\Omega)$. From [1, Proof of Theorem 3(b)] one can easily see that equality (1.2) is stronger than (1.1).

Next we would like to have a version of (1.2) which is suitable for Orlicz-Sobolev spaces embedded into multiple exponential Orlicz spaces. Recall that for $s > 0$, a measure μ on Ω , $f : \Omega \mapsto \mathbb{R}$ μ -measurable and for $\psi : [0, \mathcal{L}_n(\Omega)] \mapsto [0, \infty)$ non-decreasing and continuous on $[0, \mathcal{L}_n(\Omega)]$, differentiable on $(0, \mathcal{L}_n(\Omega))$ and satisfying $\psi(0) = 0$, we have the following well-known identity

$$(1.3) \quad \int_0^{\mathcal{L}_n(\Omega)} (f_\mu^*(t))^s \psi'(t) dt = \int_0^\infty \psi(\mu(\{x \in \Omega : |f(x)| > r\})) sr^{s-1} dr$$

(f_μ^* denotes the non-increasing rearrangement of f with respect to the measure μ). Using (1.3) and some easy estimates we obtain that (1.2) is equivalent to

$$(1.4) \quad \int_0^\infty \frac{t^{n-1}}{\log^{n-1} \left(\frac{e\mathcal{L}_n(\Omega)}{\mathcal{L}_n(\{x \in \Omega : |u(x)| \geq t\})} \right)} dt \leq C \|\nabla u\|_{L^n(\Omega)}^n$$

with the convention that we integrate only over $t \in (0, \infty)$ such that $\mathcal{L}_n(\{|u| \geq t\}) > 0$ (we define $\psi(t) = \log^{1-n}(\frac{e\mathcal{L}_n(\Omega)}{t})$ for $t \in (0, \mathcal{L}_n(\Omega)]$ and $\psi(0) = 0$). We use this convention throughout the paper.

When Ω is sufficiently nice, (1.1) turns to the following inequality for functions that do not have a zero trace on the boundary: there are $C_1 = C_1(n)$ and $C_2 = C_2(n)$ so that for every non-trivial $u \in W^{1,n}(\Omega)$ we have

$$(1.5) \quad \inf_{c \in \mathbb{R}} \int_\Omega \exp\left(\left(\frac{|u(x) - c|}{C_1 \|\nabla u\|_{L^n(\Omega)}}\right)^{\frac{n}{n-1}}\right) dx \leq C_2 \mathcal{L}_n(\Omega)$$

and (1.4) turns to

$$(1.6) \quad \inf_{c \in \mathbb{R}} \int_0^\infty \frac{t^{n-1}}{\log^{n-1} \left(\frac{e\mathcal{L}_n(\Omega)}{\mathcal{L}_n(\{x \in \Omega : |u(x) - c| \geq t\})} \right)} dt \leq C \|\nabla u\|_{L^n(\Omega)}^n$$

for every $u \in W^{1,n}(\Omega)$.

It is a surprising result by Koskela and Onninen [16] that if Ω is such that (1.5) is valid for every $u \in W^{1,n}(\Omega)$, then (1.6) is also valid for every $u \in W^{1,n}(\Omega)$. That is, with no additional requirement on Ω we have that the validity of the embedding (1.5) implies the validity of the sharper embedding (1.6). It is also proved in [16] that the Sobolev inequality for $W^{1,p}(\Omega) \hookrightarrow L^{p^*}(\Omega)$, $1 \leq p < n$, improves the same way into an inequality by O’Neil [20] and Peetre [21].

In recent paper [15], Hencl proves a version of the result from [16] for Orlicz-Sobolev spaces embedded into single and double exponential spaces.

The aim of this note is to show that the same phenomenon occurs in all Orlicz-Sobolev spaces embedded into multiple exponential Orlicz spaces.

Let us give some information concerning the spaces we are interested in. The space $W_0 L^n \log^\alpha L(\Omega)$, $\alpha < n - 1$, of the (first order) Sobolev type, modeled on

the Zygmund space $L^n \log^\alpha L(\Omega)$, is continuously embedded into the Orlicz space with the Young function that behaves like $\exp(t^{\frac{n}{n-1-\alpha}})$ for large t . These results are due to Fusco, Lions, Sbordone [9] for $\alpha < 0$ and Edmunds, Gurka, Opic [5] in general. Moreover it is shown in [5] (see also [3] and [7]) that in the limiting case $\alpha = n - 1$ we have the embedding into a double exponential space, i.e. the space $W_0L^n \log^{n-1} L \log^\alpha \log L(\Omega)$, $\alpha < n - 1$, is continuously embedded into the Orlicz space with the Young function that behaves like $\exp(\exp(t^{\frac{n}{n-1-\alpha}}))$ for large t . Further in the limiting case $\alpha = n - 1$ we have the embedding into triple exponential space and so on. The borderline case is always $\alpha = n - 1$ and for $\alpha > n - 1$ we have the embedding into $L^\infty(\Omega)$. It is well-known that the Zygmund space $L^n \log^\alpha L(\Omega)$ coincides with the Orlicz space $L^\Phi(\Omega)$, where

$$\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t^n \log^\alpha(t)} = 1,$$

the space $L^n \log^{n-1} L \log^\alpha \log L(\Omega)$ coincides with $L^\Phi(\Omega)$ where

$$\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t^n \log^{n-1}(t) \log^\alpha(\log(t))} = 1,$$

and so on. For a further discussion about the limiting cases $\alpha = n - 1$ see [6].

To simplify our notation when working with the multiple exponential spaces, let us write for $\ell \in \mathbb{N}$, $\ell \geq 2$

$$\log_{[\ell]}(t) = \log(\log_{[\ell-1]}(t)), \quad \text{where} \quad \log_{[1]}(t) = \log(t)$$

and

$$\exp_{[\ell]}(t) = \exp(\exp_{[\ell-1]}(t)), \quad \text{where} \quad \exp_{[1]}(t) = \exp(t).$$

Next, let us recall the version of (1.1) for embedding into multiple exponential spaces. Let $\Omega \subset \mathbb{R}^n$, $n \geq 2$, be a bounded domain, let $\ell \in \mathbb{N}$, $\ell \geq 2$, let $\alpha < n - 1$ and let Φ be a Young function satisfying

$$\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t^n \left(\prod_{i=1}^{\ell-1} \log_{[i]}^{n-1}(t) \right) \log_{[\ell]}^\alpha(t)} = 1.$$

Then it is shown in [5] and [9] (see also [3], [14] and [2]) that there are constants C_1 and C_2 such that

$$\int_{\Omega} \exp_{[\ell]} \left(\left(\frac{|u(x)|}{C_1 \|\nabla u\|_{L^\Phi(\Omega)}} \right)^{\frac{n}{n-1-\alpha}} \right) dx \leq C_2$$

for every non-trivial $u \in W_0L^\Phi(\Omega)$.

Following [16] and [15] we state our results in the generality which can be applied in the context of analysis on metric measure spaces. In what follows X is always a metric space equipped with a Borel measure μ and Ω is a measurable subset of X .

In the sequel we consider differentiable Young functions Φ such that

$$(1.7) \quad \lim_{t \rightarrow \infty} \frac{\Phi(t)}{t^s \left(\prod_{i=1}^{\ell-1} \log_{[i]}^{s-1}(t) \right) \log_{[\ell]}^\alpha(t)} = 1$$

with $\ell \in \mathbb{N}$, $\ell \geq 2$, $s > 1$ and $\alpha < s - 1$. We further suppose that there are $C, \delta > 0$ satisfying

$$(1.8) \quad \frac{1}{C} t^s \leq \Phi(t) \leq C t^s \quad \text{for } t \in [0, \delta).$$

Theorem 1.1. *Let $\Omega \subset X$ be a domain with $\mu(\Omega) < \infty$ and let $u, g : \Omega \rightarrow \mathbb{R}$. Fix $\ell \in \mathbb{N}$, $\ell \geq 2$, $s \in (1, \infty)$ and $\alpha \in \mathbb{R}$, $\alpha < s - 1$. Set $E = \exp_{[\ell]}(1)$. Suppose that Φ is a Young function satisfying (1.7) and (1.8). Assume that the inequality*

$$(1.9) \quad \inf_{c \in \mathbb{R}} \int_{\Omega} \exp_{[\ell]} \left(\left(\frac{|u(y) - c|}{C_1 \|g\|_{L^\Phi(\Omega)}} \right)^{\frac{s}{s-1-\alpha}} \right) d\mu(y) \leq C_2$$

is stable under truncation. Then

$$(1.10) \quad \inf_{c \in \mathbb{R}} \int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{x \in \Omega : |u(x) - c| \geq t\})} \right)} dt < \infty.$$

The requirement that the inequality (1.9) is stable under truncation means that for every $d \in \mathbb{R}$, $0 < t_1 < t_2 < \infty$ and $z \in \{-1, 1\}$ the pairs $v_{t_1}^{t_2}, g_{t_1}^{t_2} = g\chi_{\{t_1 < v \leq t_2\}}$, where $v = z(u - d)$ and $v_{t_1}^{t_2} = \min\{\max\{0, v - t_1\}, t_2 - t_1\}$, also satisfy (1.9):

$$\inf_{c \in \mathbb{R}} \int_{\Omega} \exp_{[\ell]} \left(\left(\frac{|v_{t_1}^{t_2}(y) - c|}{C_1 \|g_{t_1}^{t_2}\|_{L^\Phi(\Omega)}} \right)^{\frac{s}{s-1-\alpha}} \right) d\mu(y) \leq C_2.$$

Notice that the function u clearly satisfies the truncation property if $\Omega \subset \mathbb{R}^n$, $s = n$, $\mu = \mathcal{L}_n$ and $g = |\nabla u|$. For further applications of the powerful truncation technique which was first used in [18] we refer the reader to [17], [10] and references given there.

The validity of (1.10) is known in the Euclidean setting if we deal only with functions with zero traces (see [5], [8] and [4]). Again these spaces serve as the best rearrangement invariant target space of the embedding of $W_0L^\Phi(\Omega)$. Our approach gives a new proof of these embeddings and we have additional information if we deal with functions that do not have a zero trace on the boundary.

The paper is organized the following way. In the third section we study some properties of the functions $\exp_{[j]}$ and $\log_{[j]}$, $j \in \mathbb{N}$. The fourth section is devoted to the proof of Theorem 1.1.

2. Preliminaries

We denote by \mathcal{L}_n the n -dimensional Lebesgue measure. For two functions $h, g : I \mapsto \mathbb{R}$ we write $h \sim g$ on I if there is a constant $C > 1$ such that $\frac{1}{C}h(t) \leq g(t) \leq Ch(t)$ for every $t \in I$. When $I = [0, \infty)$ we simply write $h \sim g$.

A function $\Phi : [0, \infty) \mapsto [0, \infty)$ is a Young function if $\Phi(0) = 0$, Φ is increasing, convex and $\lim_{t \rightarrow \infty} \frac{\Phi(t)}{t} = \infty$. For a fixed measure μ , we denote by $L^\Phi(\Omega)$ the Orlicz space corresponding to a Young function Φ on a set Ω with a measure μ . This space is equipped with the Luxemburg norm

$$\|f\|_{L^\Phi(\Omega)} = \inf \left\{ \lambda > 0 : \int_{\Omega} \Phi \left(\frac{|f(x)|}{\lambda} \right) d\mu(x) \leq 1 \right\}.$$

For an introduction to Orlicz spaces see [23]. By $WL^\Phi(\Omega)$ we denote the set of functions f such that $f, |\nabla f| \in L^\Phi(\Omega)$ and by $W_0L^\Phi(\Omega)$ we denote the closure of $C_0^\infty(\Omega)$ in $WL^\Phi(\Omega)$.

Let $\ell \in \mathbb{N}$, $\ell \geq 2$, $s > 1$ and $\alpha < s - 1$. Suppose that the Young function Φ satisfies (1.7) and (1.8). Let us define auxiliary functions $\varphi_1, \Phi_1 : [0, \infty) \mapsto [0, \infty)$ by

$$\varphi_1(t) = \left(\prod_{j=1}^{\ell} \log_{[j]}^{s-1}(E+t) \right) \log_{[j]}^{\alpha}(E+t), \quad \Phi_1(t) = t^s \varphi_1(t), \quad t \geq 0.$$

From conditions (1.7), (1.8) we see that for any fixed $t_0 > 0$ we have

$$(2.1) \quad \Phi_1(t) \geq \frac{1}{C}t^s, \quad \Phi \sim \Phi_1, \quad \varphi_1 \sim 1 \text{ on } [0, t_0] \text{ and } \Phi_1(t) \sim t^s \text{ on } [0, t_0].$$

We say that a function Φ satisfies the Δ_2 -condition if there is $C_\Delta > 0$ such that $\Phi(2t) \leq C_\Delta \Phi(t)$ for every $t \geq 0$. If Φ satisfies the Δ_2 -condition then (see [23, Proposition 6, p. 77])

$$(2.2) \quad \int_{\Omega} \Phi \left(\frac{|f(x)|}{\|f\|_{L^\Phi(\Omega)}} \right) d\mu(x) = 1 \quad \text{provided } \|f\|_{L^\Phi(\Omega)} > 0.$$

Notice that our function Φ satisfies Δ_2 -condition thanks to (1.7) and (1.8). And so do φ_1 and Φ_1 .

Let $\Psi : [0, \infty) \mapsto [0, \infty)$ be an increasing convex function and let $h : S \rightarrow \mathbb{R}$ be a non-negative function. Then we can use the following version of Jensen’s inequality:

$$(2.3) \quad \frac{1}{\mu(S)} \int_S h(x) dx \leq \Psi^{-1} \left(\frac{1}{\mu(S)} \int_S \Psi(h(x)) dx \right).$$

We also use a simple lemma from [16].

Lemma 2.1. *Let ν be a finite measure on a set Y . If $w : Y \mapsto [0, \infty)$ is a ν -measurable function such that $\nu(\{y \in Y : w(y) = 0\}) \geq \frac{\nu(Y)}{2}$, then, for every $t > 0$ we have*

$$\nu(\{y \in Y : w(y) \geq t\}) \leq 2 \inf_{c \in \mathbb{R}} \nu\left(\left\{y \in Y : |w(y) - c| \geq \frac{t}{2}\right\}\right).$$

By C we denote a generic positive constant that may depend on $\ell, s, \alpha, C_1, K, \|g\|_{L^\Phi(\Omega)}$ and $\|f\|_{L^\Phi(\Omega)}$. This constant may vary from expression to expression as usual.

3. Some properties of the functions $\exp_{[j]}$ and $\log_{[j]}$

Lemma 3.1. *Let $a, b, d \geq 1$. Then for every $j \in \mathbb{N}, j \leq \ell$ we have*

$$(3.1) \quad \log_{[j]}(E + ab) \leq 2 \log(E + b) \log_{[j]}(E + a)$$

and

$$(3.2) \quad \log_{[j]}(E + a^d) \leq C \log_{[j]}(E + a).$$

PROOF: Let us prove (3.1). Using the fact that for $x, y \geq 1$ we have $x + y \leq 2xy$ we obtain

$$\begin{aligned} \log(E + ab) &\leq \log(Eb + ab) = \log(b) + \log(E + a) \\ &\leq \log(E + b) + \log(E + a) \leq 2 \log(E + b) \log(E + a). \end{aligned}$$

Similarly we use the inequality $2 \log(E + b) \leq E + b$ and above estimate to obtain

$$\begin{aligned} \log_{[2]}(E + ab) &\leq \log\left(2 \log(E + b) \log(E + a)\right) \leq \log\left((E + b) \log(E + a)\right) \\ &= \log(E + b) + \log_{[2]}(E + a) \leq 2 \log(E + b) \log_{[2]}(E + a) \end{aligned}$$

and we continue by induction.

Now, let us prove (3.2). We have

$$\log(E + a^d) \leq \log((E + a)^d) = d \log(E + a)$$

and thus

$$\log_{[2]}(E + a^d) \leq \log(C \log(E + a)) = \log(C) + \log_{[2]}(E + a) \leq C \log_{[2]}(E + a).$$

We continue by induction. □

Lemma 3.2. *If $t \geq 0$, then*

$$t^{k_\ell} \leq \frac{\prod_{i=1}^\ell k_i!}{\prod_{i=1}^{\ell-1} k_i^{k_{i+1}}} \exp_{[\ell]}(t)$$

whenever $k_i \in \mathbb{N}, i = 1, \dots, \ell$.

PROOF: We have $\exp(t) = \sum_{k_1=0}^{\infty} \frac{t^{k_1}}{k_1!}$,

$$\exp_{[2]}(t) = \sum_{k_1=0}^{\infty} \frac{\exp^{k_1}(t)}{k_1!} = \sum_{k_1=0}^{\infty} \frac{\exp(k_1 t)}{k_1!} = \sum_{k_1, k_2=0}^{\infty} \frac{k_1^{k_2} t^{k_2}}{k_1! k_2!}$$

and by induction

$$\exp_{[\ell]}(t) = \sum_{k_1, \dots, k_{\ell}=0}^{\infty} \frac{\prod_{i=1}^{\ell-1} k_i^{k_{i+1}}}{\prod_{i=1}^{\ell} k_i!} t^{k_{\ell}}.$$

Each summand on the right hand side is estimated by $\exp_{[\ell]}(t)$ and we are done. \square

Lemma 3.3. *Suppose that $\xi, \psi > 0$ satisfy*

$$\xi^{\frac{1}{k_{\ell}}} \leq C \frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{k_{\ell}}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{k_{\ell}}}} \psi \quad \text{where } k_i \in \mathbb{N}, k_i \leq k_{\ell}, i = 1, \dots, \ell.$$

Then

$$\xi^{\frac{1}{a_{\ell}}} \leq C \frac{\prod_{i=1}^{\ell} a_i^{\frac{a_i}{a_{\ell}}}}{\prod_{i=1}^{\ell-1} a_i^{\frac{a_{i+1}}{a_{\ell}}}} \psi \quad \text{for every } a_i \in [1, \infty), a_i \leq a_{\ell}, i = 1, \dots, \ell.$$

PROOF: First let us show that we have

$$(3.3) \quad \xi^{\frac{1}{b}} \leq C \frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{b}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{b}}} \psi \quad \text{for every } b \in [1, \infty), k_i \leq b + 1, i = 1, \dots, \ell.$$

Let $m \in \mathbb{N}$ be the integer part of b . Then by assumption we have

$$(3.4) \quad \begin{aligned} \xi^{\frac{1}{b}} &\leq \max(\xi^{\frac{1}{m+1}}, \xi^{\frac{1}{m}}) \leq C\psi \max\left(\frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{m}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{m}}}, \frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{m+1}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{m+1}}}\right) \\ &\leq C\psi \frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{m}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{m+1}}}. \end{aligned}$$

Next let us prove

$$(3.5) \quad k_i^{\frac{k_i}{m}} \leq C k_i^{\frac{k_i}{b}}, \quad i = 1, \dots, \ell \quad \text{and} \quad k_i^{\frac{k_{i+1}}{b}} \leq C k_i^{\frac{k_{i+1}}{m+1}}, \quad i = 1, \dots, \ell - 1.$$

The first inequality in (3.5) follows from

$$k_i^{\frac{k_i}{m} - \frac{k_i}{b}} = k_i^{\frac{k_i(b-m)}{bm}} \leq k_i^{\frac{k_i}{b}} \leq (b+1)^{\frac{b+1}{bm}} \leq (3m)^{\frac{3m}{m^2}} = (3m)^{\frac{3}{m}} \leq C.$$

The second inequality in (3.5) is proved by

$$k_i \frac{k_{i+1}}{b} - \frac{k_{i+1}}{m+1} = k_i \frac{k_{i+1}(m+1-b)}{b(m+1)} \leq k_i \frac{k_{i+1}}{bm} \leq (b+1) \frac{b+1}{bm} \leq (3m) \frac{3m}{m^2} = (3m) \frac{3}{m} \leq C.$$

Now, (3.3) follows from (3.4) and (3.5).

Next, we are going to prove assertion of the lemma applying inequality (3.3) with k_i being the integer parts of a_i , $i = 1, \dots, \ell$. For $i = 1, \dots, \ell - 1$ we observe that

$$a_i \frac{a_{i+1}}{b} = \left(\frac{a_i}{k_i}\right) \frac{a_{i+1}}{b} k_i \frac{a_{i+1}}{b} - \frac{k_{i+1}}{b} k_i \frac{k_{i+1}}{b} \leq 2^2 k_i^{\frac{1}{b}} k_i^{\frac{k_{i+1}}{b}} \leq 2^2 (2b)^{\frac{1}{b}} k_i^{\frac{k_{i+1}}{b}} \leq C k_i^{\frac{k_{i+1}}{b}}.$$

Therefore

$$(3.6) \quad \frac{\prod_{i=1}^{\ell} k_i^{\frac{k_i}{b}}}{\prod_{i=1}^{\ell-1} k_i^{\frac{k_{i+1}}{b}}} \leq C \frac{\prod_{i=1}^{\ell} a_i^{\frac{a_i}{b}}}{\prod_{i=1}^{\ell-1} a_i^{\frac{a_{i+1}}{b}}} \text{ for every } b \in [1, \infty), a_i \leq b+1, i = 1, \dots, \ell.$$

Now, we set $a_\ell = b$ and (3.3) together with (3.6) conclude the proof. □

Lemma 3.4. *Let Ψ be a non-negative increasing function satisfying $\Psi(t) \sim t\varphi_1(t)$ for $t \geq 0$. Then there is $C_\Psi > 0$ such that the inverse function Ψ^{-1} satisfies on $[0, \infty)$*

$$\Psi^{-1}(t) \leq C_\Psi t \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{1-s}(E+t) \right) \log_{[\ell]}^{-\alpha}(E+t) = C_\Psi \frac{t}{\varphi_1(t)} =: \tilde{\Psi}(t).$$

PROOF: First, let us prove that there is $t_1 > 0$ such that

$$(3.7) \quad \log_{[j]}(E+t^{\frac{1}{2}}) \geq \frac{1}{2} \log_{[j]}(E+t) \text{ for } t \geq t_1, j \in \mathbb{N}, j \leq \ell.$$

For $j = 1$ it is obvious. For $j = 2$ we have

$$\log_{[2]}(E+t^{\frac{1}{2}}) \geq \log\left(\frac{1}{2} \log(E+t)\right) = \log_{[2]}(E+t) - \log(2) \geq \frac{1}{2} \log_{[2]}(E+t)$$

provided t is large enough. And we continue by induction.

Further, we see that for $\alpha \geq 0$ there is $t_2 \geq t_1$ such that for $t \geq t_2$ we have from (3.7)

$$(3.8) \quad \log_{[\ell]}^\alpha(E + \tilde{\Psi}(t)) \geq \log_{[\ell]}^\alpha(E + t^{\frac{1}{2}}) \geq \frac{1}{2^\alpha} \log_{[\ell]}^\alpha(E + t)$$

while for $\alpha < 0$ we find $t_2 \geq t_1$ so that for every $t \geq t_2$ we obtain

$$(3.9) \quad \log_{[\ell]}^\alpha(E + \tilde{\Psi}(t)) \geq \log_{[\ell]}^\alpha(E + t) > \frac{1}{2^{|\alpha|}} \log_{[\ell]}^\alpha(E + t).$$

Therefore by (3.7), (3.8) and (3.9) we have for $t \geq t_2$

$$\begin{aligned} \Psi(\tilde{\Psi}(t)) &\geq \frac{1}{C} \tilde{\Psi}(t) \varphi_1(\tilde{\Psi}(t)) \\ &= \frac{C_\Psi}{C} t \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{1-s}(E+t) \right) \log_{[\ell]}^{-\alpha}(E+t) \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{s-1}(E+\tilde{\Psi}(t)) \right) \\ &\quad \times \log_{[\ell]}^\alpha(E+\tilde{\Psi}(t)) \\ &\geq \frac{C_\Psi}{C} t \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{1-s}(E+t) \right) \log_{[\ell]}^{-\alpha}(E+t) \\ &\quad \times \frac{1}{2^{(s-1)(\ell-1)}} \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{s-1}(E+t) \right) \frac{1}{2^{|\alpha|}} \log_{[\ell]}^\alpha(E+t) \\ &\geq \frac{C_\Psi}{C} t. \end{aligned}$$

Thus $\Psi^{-1}(t) \leq \tilde{\Psi}(t)$ on $[t_2, \infty)$ provided C_Ψ is large enough. On the other hand we have $\Psi(t) \sim t$ on every bounded interval by (2.1) and thus $\Psi^{-1}(t) \sim t$ on every bounded interval. As $\frac{1}{\varphi_1}$ is bounded away from zero on any bounded interval, we have $\tilde{\Psi}(t) \sim t$ there and we are done. □

4. Proof of Theorem 1.1

In this section we prove Theorem 1.1. Our proof is very similar to the proofs from [15] (thanks to our auxiliary lemmata from the previous section).

Lemma 4.1. *Suppose that the functions $f_k : \Omega \rightarrow \mathbb{R}$ have pairwise disjoint supports and that $f = \sum_{k=1}^\infty f_k \in L^\Phi(\Omega)$. We further assume that for every $k \in \mathbb{N}$ such that $\|f_k\|_{L^\Phi(\Omega)} > 0$ we have*

$$(4.1) \quad (s+2) \log\left(\frac{1}{\|f_k\|_{L^\Phi(\Omega)}}\right) < \log\left(\frac{E\mu(\Omega)}{\mu(\{f_k \neq 0\})}\right) + C.$$

Then

$$\sum_{k=1}^\infty \|f_k\|_{L^\Phi(\Omega)}^s < \infty.$$

PROOF: Denote $\lambda_k = \|f_k\|_{L^\Phi(\Omega)}$. Without loss of generality we can suppose that $\lambda_k > 0$ for every $k \in \mathbb{N}$. We can further suppose that $\|f\|_{L^\Phi(\Omega)} = 1$. Indeed, otherwise we replace f_k with $\frac{f_k}{\|f\|_{L^\Phi(\Omega)}}$, $k \in \mathbb{N}$, which are functions satisfying the

following version of (4.1)

$$\begin{aligned}
 (s + 2) \log\left(\frac{1}{\left\|\frac{f_k}{\|f\|_{L^\Phi(\Omega)}}\right\|_{L^\Phi(\Omega)}}\right) &= (s + 2) \log\left(\frac{1}{\|f_k\|_{L^\Phi(\Omega)}}\right) + (s + 2) \log(\|f\|_{L^\Phi(\Omega)}) \\
 &\leq \log\left(\frac{E\mu(\Omega)}{\mu(\{f_k \neq 0\})}\right) + C + (s + 2) \max(0, \log(\|f\|_{L^\Phi(\Omega)})) \\
 &= \log\left(\frac{E\mu(\Omega)}{\mu(\{\frac{f_k}{\|f\|_{L^\Phi(\Omega)}} \neq 0\})}\right) + C.
 \end{aligned}$$

Hence we have $\lambda_k \in (0, 1]$, for every $k \in \mathbb{N}$. Notice that (4.1) implies

$$(4.2) \quad (s + 2) \log\left(E + \frac{1}{\lambda_k}\right) < \log\left(\frac{E\mu(\Omega)}{\mu(\{f_k \neq 0\})}\right) + C.$$

Let $k_0 \in \mathbb{N}$ be fixed (value of k_0 is given bellow, we need (4.8) to be satisfied). The function φ_1 is increasing for t large and satisfies the Δ_2 -condition. Hence by (3.2) from Lemma 3.1 and the inequality $ab \leq a^2 + b^2$, $a, b \in \mathbb{R}$, we have

$$\begin{aligned}
 \varphi_1\left(\frac{|f_k|}{\lambda_k}\right) &\leq C + \varphi_1\left(|f_k|^2 + \frac{1}{\lambda_k^2}\right) \leq C + C\varphi_1\left(|f_k|^2\right) + C\varphi_1\left(\frac{1}{\lambda_k^2}\right) \\
 &\leq C + C\varphi_1(|f_k|) + C\varphi_1\left(\frac{1}{\lambda_k}\right).
 \end{aligned}$$

Therefore (2.1) and (2.2) give

$$\begin{aligned}
 \sum_{k=1}^{\infty} \lambda_k^s &= \sum_{k=1}^{k_0} \lambda_k^s + \sum_{k=k_0+1}^{\infty} \int_{\Omega} \lambda_k^s \Phi\left(\frac{|f_k|}{\lambda_k}\right) d\mu \\
 &\leq \sum_{k=1}^{k_0} \|f\|_{L^\Phi(\Omega)}^s + C \sum_{k=k_0+1}^{\infty} \int_{\Omega} \lambda_k^s \Phi_1\left(\frac{|f_k|}{\lambda_k}\right) d\mu \\
 (4.3) \quad &= C + C \sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s \varphi_1\left(\frac{|f_k|}{\lambda_k}\right) d\mu \\
 &\leq C + C \left(\sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s d\mu + \sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s \varphi_1(|f_k|) d\mu \right. \\
 &\quad \left. + \sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s \varphi_1\left(\frac{1}{\lambda_k}\right) d\mu \right) \\
 &= C + C(S_1 + S_2 + S_3).
 \end{aligned}$$

Notice that we have by (2.1) and (2.2)

$$(4.4) \quad \sum_{k=1}^{\infty} \int_{\Omega} \Phi_1(|f_k|) d\mu = \int_{\Omega} \Phi_1(|f|) d\mu \leq C \int_{\Omega} \Phi(|f|) d\mu = C$$

and

$$(4.5) \quad \sum_{k=1}^{\infty} \int_{\Omega} |f_k|^s d\mu \leq C \sum_{k=1}^{\infty} \int_{\Omega} \Phi_1(|f_k|) d\mu \leq C.$$

From (4.5) we obtain

$$(4.6) \quad S_1 = \sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s d\mu \leq C$$

and (4.4) implies

$$(4.7) \quad S_2 = \sum_{k=k_0+1}^{\infty} \int_{\Omega} \Phi_1(|f_k|) d\mu \leq C.$$

It remains to estimate S_3 . First, we claim that there is $k_0 \in \mathbb{N}$ such that

$$(4.8) \quad \log\left(E + \frac{1}{\lambda_k}\right) \leq C \log\left(E + \frac{1}{\mu(\{f_k \neq 0\})}\right) \int_{\Omega} \Phi(|f_k|) d\mu$$

for every $k \geq k_0$. Let us prove this claim. From (2.2), $\lambda_k \leq 1$ and inequality (3.1) from Lemma 3.1 we obtain

$$\begin{aligned} \lambda_k^s &= \int_{\Omega} \lambda_k^s \Phi\left(\frac{|f_k|}{\lambda_k}\right) d\mu \leq C \int_{\Omega} \lambda_k^s \Phi_1\left(\frac{|f_k|}{\lambda_k}\right) d\mu = C \int_{\Omega} |f_k|^s \varphi_1\left(\frac{|f_k|}{\lambda_k}\right) d\mu \\ &= \int_{\Omega} |f_k|^s \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{s-1}\left(E + \frac{|f_k|}{\lambda_k}\right)\right) \log_{[\ell]}^{\alpha}\left(E + \frac{|f_k|}{\lambda_k}\right) d\mu \\ &\leq C \log^{(\ell-1)(s-1)+|\alpha|}\left(E + \frac{1}{\lambda_k}\right) \int_{\Omega} |f_k|^s \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{s-1}(E + |f_k|)\right) \log_{[\ell]}^{\alpha}(E + |f_k|) d\mu \\ &\leq C \frac{1}{\lambda_k} \int_{\Omega} |f_k|^s \left(\prod_{j=1}^{\ell-1} \log_{[j]}^{s-1}(E + |f_k|)\right) \log_{[\ell]}^{\alpha}(E + |f_k|) d\mu \\ &= C \frac{1}{\lambda_k} \int_{\Omega} \Phi_1(|f_k|) d\mu \leq C \frac{1}{\lambda_k} \int_{\Omega} \Phi(|f_k|) d\mu. \end{aligned}$$

This implies

$$-(s+1) \log\left(E + \frac{1}{\lambda_k}\right) \leq C + \log\left(\int_{\Omega} \Phi(|f_k|) d\mu\right).$$

Summing up this inequality and (4.2) we obtain

$$\log\left(E + \frac{1}{\lambda_k}\right) \leq \log\left(E + \frac{1}{\mu(\{f_k \neq 0\})} \int_{\Omega} \Phi(|f_k|) d\mu\right) + C.$$

Therefore, since $\lambda_k \rightarrow 0$ we easily find $k_0 \in \mathbb{N}$ large enough so that (4.8) is satisfied for every $k \geq k_0$.

Now, we can start estimating S_3 . From the definition of φ_1 , the fact that $\varphi_1(t)$ is increasing for large t and from (4.8) we obtain

$$\varphi_1\left(\frac{1}{\lambda_k}\right) \leq C + C\varphi_1\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\Omega} \Phi(|f_k|) d\mu\right).$$

Hence

(4.9)

$$\begin{aligned} S_3 &= \sum_{k=k_0+1}^{\infty} \varphi_1\left(\frac{1}{\lambda_k}\right) \int_{\Omega} |f_k|^s d\mu \\ &\leq C \sum_{k=k_0+1}^{\infty} \int_{\Omega} |f_k|^s d\mu + C \sum_{k=k_0+1}^{\infty} \varphi_1\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\Omega} \Phi(|f_k|) d\mu\right) \int_{\Omega} |f_k|^s d\mu. \end{aligned}$$

Thus we need a suitable estimate of $\int_{\Omega} |f_k|^s d\mu$.

Fix an increasing convex function $\Psi : [0, \infty) \mapsto [0, \infty)$ such that $\Psi(t) \sim t\varphi_1(t)$. Therefore Ψ and Ψ^{-1} satisfy the Δ_2 -condition and Ψ^{-1} can be estimated by $\tilde{\Psi}$ from Lemma 3.4. Thus from Jensen's inequality (2.3) for the function $h = |f_k|^s$ and $S = \{f_k \neq 0\}$ we obtain

$$\begin{aligned} \frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} |f_k|^s d\mu &\leq \Psi^{-1}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} \Psi(|f_k|^s) d\mu\right) \\ &\leq \Psi^{-1}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} C|f_k|^s \varphi_1(|f_k|^s) d\mu\right). \end{aligned}$$

Next we use the fact that $\varphi_1(t^s) \leq C\varphi_1(t)$ (see (3.2)), (2.1) and Lemma 3.4

$$\begin{aligned} \frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} |f_k|^s d\mu &\leq \Psi^{-1}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} C|f_k|^s \varphi_1(|f_k|) d\mu\right) \\ &= \Psi^{-1}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} C\Phi_1(|f_k|) d\mu\right) \\ &\leq \Psi^{-1}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} C\Phi(|f_k|) d\mu\right) \\ &\leq \tilde{\Psi}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} C\Phi(|f_k|) d\mu\right). \end{aligned}$$

Now, we can plainly suppose that the constant C on the last line satisfies $C \geq 1$. Therefore, as $\varphi_1(t)$ is non-decreasing for large t and bounded away from zero on

$[0, \infty)$, we have $\frac{1}{\varphi_1(Ct)} \leq \frac{C}{\varphi_1(t)}$ and thus $\tilde{\Psi}(Ct) \leq C\tilde{\Psi}(t)$ on $[0, \infty)$. Hence we obtain

$$\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} |f_k|^s d\mu \leq C\tilde{\Psi}\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} \Phi(|f_k|) d\mu\right).$$

Therefore we have

$$\int_{\{f_k \neq 0\}} |f_k|^s d\mu \leq C \int_{\{f_k \neq 0\}} \Phi(|f_k|) d\mu \frac{1}{\varphi_1\left(\frac{1}{\mu(\{f_k \neq 0\})} \int_{\{f_k \neq 0\}} \Phi(|f_k|) d\mu\right)}.$$

This estimate, (2.1), (4.4), (4.5) and (4.9) imply

$$(4.10) \quad S_3 \leq C.$$

Now (4.3), (4.6), (4.7) and (4.10) conclude the proof. □

PROOF OF THEOREM 1.1: Let us choose $d \in \mathbb{R}$ such that

$$\mu(\{u \geq d\}) \geq \frac{\mu(\Omega)}{2} \quad \text{and} \quad \mu(\{u \leq d\}) \geq \frac{\mu(\Omega)}{2}.$$

Set $v_+ = \max\{u - d, 0\}$ and $v_- = -\min\{u - d, 0\}$. In the sequel v stands for v_+ and v_- , respectively. Our aim is to prove

$$(4.11) \quad \int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha}\left(\frac{E\mu(\Omega)}{\mu(\{v \geq t\})}\right)} dt < \infty \quad \text{for } v = v_+, v = v_-.$$

First, let us show how (4.11) concludes the proof. Since $\{|u - d| \geq t\} = \{v_+ \geq t\} \cup \{v_- \geq t\}$, we have

$$\mu(\{|u - d| \geq t\}) \leq 2 \max\{\mu(\{v_+ \geq t\}), \mu(\{v_- \geq t\})\}.$$

Moreover we have for all $s \in [1, \infty)$

$$\frac{1}{\log_{[\ell]}(Es)} \leq C \frac{1}{\log_{[\ell]}(2Es)}.$$

From this estimate and (4.11) we obtain

$$(4.12) \quad \begin{aligned} & \inf_{c \in \mathbb{R}} \int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha}\left(\frac{E\mu(\Omega)}{\mu(\{x \in \Omega: |u(x) - c| \geq t\})}\right)} dt \\ & \leq \int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha}\left(\frac{E\mu(\Omega)}{\mu(\{x \in \Omega: |u(x) - d| \geq t\})}\right)} dt \\ & \leq C \left(\int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha}\left(\frac{E\mu(\Omega)}{\mu(\{v_+ \geq t\})}\right)} dt + \int_0^\infty \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha}\left(\frac{E\mu(\Omega)}{\mu(\{v_- \geq t\})}\right)} dt \right) < \infty \end{aligned}$$

which is the assertion of the theorem.

In the rest of the proof we establish (4.11). We distinguish two cases.

If $v \in L^\infty(\Omega)$, then inequality (4.11) is obviously satisfied (recall the convention that we integrate over $t \in (0, \infty)$ such that $\mu(\{v \geq t\}) > 0$ only) and thus we are done.

Hence we can suppose that $v \notin L^\infty(\Omega)$ in the rest of the proof.

STEP 1.

Fix $0 < t_1 < t_2 < \infty$. From (1.9), the truncation property and Lemma 3.2 we have

$$(4.13) \quad \inf_{c \in \mathbb{R}} \left(\int_{\Omega} |v_{t_1}^{t_2} - c|^{\frac{sk_\ell}{s-1-\alpha}} d\mu \right)^{\frac{s-1-\alpha}{sk_\ell}} \leq C \left(\frac{\prod_{i=1}^\ell k_i!}{\prod_{i=1}^{\ell-1} k_{i+1}} \right)^{\frac{s-1-\alpha}{sk_\ell}} \|g_{t_1}^{t_2}\|_{L^\Phi(\Omega)}$$

whenever $k_i \in \mathbb{N}$, $i = 1, \dots, \ell$. From Lemma 2.1 and the weak form of (4.13) we obtain

$$\begin{aligned} t[\mu(\{v_{t_1}^{t_2} \geq t\})]^{\frac{s-1-\alpha}{sk_\ell}} &\leq C \inf_{c \in \mathbb{R}} \frac{t}{2} \left[\mu \left(\left\{ |v_{t_1}^{t_2} - c| \geq \frac{t}{2} \right\} \right) \right]^{\frac{s-1-\alpha}{sk_\ell}} \\ &\leq C (\mu(\Omega))^{\frac{s-1-\alpha}{sk_\ell}} \left(\frac{\prod_{i=1}^\ell k_i!}{\prod_{i=1}^{\ell-1} k_{i+1}} \right)^{\frac{s-1-\alpha}{sk_\ell}} \|g_{t_1}^{t_2}\|_{L^\Phi(\Omega)} \end{aligned}$$

for $k_i \in \mathbb{N}$, $i = 1, \dots, \ell$ and every $t > 0$. Since $(k!)^\frac{1}{k} \sim k^\frac{k}{k}$ if $k \leq l$, from above and from Lemma 3.3 we see that

$$(4.14) \quad t \left(\frac{\mu(\{v_{t_1}^{t_2} \geq t\})}{E\mu(\Omega)} \right)^{\frac{s-1-\alpha}{sa_\ell}} \leq C \left(\frac{\prod_{i=1}^\ell a_i^{\frac{\alpha_i}{a_\ell}}}{\prod_{i=1}^{\ell-1} a_i^{\frac{\alpha_{i+1}}{a_\ell}}} \right)^{\frac{s-1-\alpha}{s}} \|g_{t_1}^{t_2}\|_{L^\Phi(\Omega)}$$

for $a_i \in [1, \infty)$, $a_i \leq a_\ell$, $i = 1, \dots, \ell$ and $t > 0$.

STEP 2.

Our next step is to prove

$$(4.15) \quad \frac{2^i}{\log_{[\ell]}^{\frac{s-1-\alpha}{s}} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})} \right)} \leq C \|g_{2^i}^{2^{i+1}}\|_{L^\Phi(\Omega)} \quad \text{whenever } i \in \mathbb{N}.$$

Let us define $b = \frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})}$. We set

$$a_i = \frac{\log(b)}{\log_{[i+1]}(b)} \quad \text{for } i = 1, \dots, \ell - 1 \quad \text{and} \quad a_\ell = \log(b).$$

Hence as $t^{\frac{1}{\log(t)}} = e$, $(\frac{1}{t})^{\frac{1}{\log(t)}} = e^{-1}$, $b \geq E$ and $\lim_{t \rightarrow \infty} (\frac{1}{t})^{\frac{1}{t}} = 1$, we obtain

$$\begin{aligned}
 \frac{\prod_{i=1}^{\ell} a_i^{\frac{\alpha_i}{\alpha_{\ell}}}}{\prod_{i=1}^{\ell-1} a_i^{\frac{\alpha_{i+1}}{\alpha_{\ell}}}} &= \frac{\left(\prod_{i=1}^{\ell-1} \left(\frac{\log(b)}{\log_{[i+1]}(b)}\right)^{\frac{1}{\log_{[i+1]}(b)}}\right) \log(b)}{\left(\prod_{i=1}^{\ell-2} \left(\frac{\log(b)}{\log_{[i+1]}(b)}\right)^{\frac{1}{\log_{[i+2]}(b)}}\right) \frac{\log(b)}{\log_{[\ell]}(b)}} \\
 (4.16) \qquad &= \frac{\log_{[\ell]}(b) \log^{\frac{1}{\log_{[2]}(b)}}(b) \left(\prod_{i=1}^{\ell-1} \left(\frac{1}{\log_{[i+1]}(b)}\right)^{\frac{1}{\log_{[i+1]}(b)}}\right)}{\prod_{i=1}^{\ell-2} \left(\frac{1}{\log_{[i+1]}(b)}\right)^{\frac{1}{\log_{[i+2]}(b)}}} \\
 &\sim \log_{[\ell]}(b).
 \end{aligned}$$

Next we observe that $(\frac{1}{b})^{\frac{s-1-\alpha}{s \log(b)}} = e^{-\frac{s-1-\alpha}{s}} = C$ and $\{v_{2^i}^{2^{i+1}} \geq 2^i\} = \{v \geq 2^{i+1}\}$. Hence from (4.14) with $t = 2^i$, $t_1 = 2^i$, $t_2 = 2^{i+1}$ and (4.16) we obtain (4.15).

STEP 3.

Set $S_i = \{v \geq 2^i\}$,

$$G = \left\{ i \in \mathbb{N}_0 : \log_{[\ell]} \left(\frac{E\mu(\Omega)}{\mu(S_{i+1})} \right) < K 4^{\frac{s}{s-1-\alpha}} \log_{[\ell]} \left(\frac{E\mu(\Omega)}{\mu(S_i)} \right) \right\}$$

and $B = \mathbb{N}_0 \setminus G$, where $K \geq 1$ is large enough so that $0 \in G$. Notice that G and B are well-defined, because $v \notin L^\infty(\Omega)$.

Lemma 2.1 implies

$$\mu(\{v \geq 2^{i+1}\}) = \mu(\{v_{2^i}^{2^{i+1}} \geq 2^i\}) \leq 2 \inf_{c \in \mathbb{R}} \mu(\{|v_{2^i}^{2^{i+1}} - c| \geq 2^{i-1}\}).$$

Hence we can use (1.9) and the truncation property for $t_1 = 2^i$ and $t_2 = 2^{i+1}$ to obtain

$$\mu(\{v \geq 2^{i+1}\}) \exp_{[\ell]} \left(\left(\frac{2^{i-1}}{C \|g_{2^i}^{2^{i+1}}\|_{L^\Phi(\Omega)}} \right)^{\frac{s}{s-1-\alpha}} \right) \leq C_2.$$

Further we observe that

$$\{g_{2^i}^{2^{i+1}} \neq 0\} = \{g_{\chi_{2^i < v \leq 2^{i+1}}} \neq 0\} \subset \{2^i < v\} \subset \{2^i \leq v\} = S_i.$$

Thus for $i \in G$ we have

$$\begin{aligned}
 \frac{1}{\|g_{2^i}^{2^{i+1}}\|_{L^\Phi(\Omega)}} &\leq C \log_{[\ell]}^{\frac{s-1-\alpha}{s}} \left(E + \frac{C}{\mu(S_{i+1})} \right) \\
 &\leq C \log_{[\ell]}^{\frac{s-1-\alpha}{s}} \left(E + \frac{C}{\mu(S_i)} \right) \\
 &\leq C \log_{[\ell]}^{\frac{s-1-\alpha}{s}} \left(E + \frac{C}{\mu(\{g_{2^i}^{2^{i+1}} \neq 0\})} \right).
 \end{aligned}$$

This verifies assumption (4.1) and therefore Lemma 4.1 and (4.15) give us

$$(4.17) \quad \sum_{i \in G} \frac{2^{si}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})} \right)} \leq C \sum_{i \in G} \|g_{2^i}^{2^{i+1}}\|_{L^\Phi(\Omega)}^s < \infty.$$

Next, let us suitably decompose B . For each $i \in G$ we define

$$B_i = \{j \in B : j > i \text{ and } \{i + 1, i + 2, \dots, j\} \subset B\}.$$

From the definition of B , simple induction and (4.17) we have

$$(4.18) \quad \begin{aligned} \sum_{j \in B} \frac{2^{sj}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{j+1}\})} \right)} &= \sum_{i \in G} \sum_{j \in B_i} \frac{2^{sj}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(S_{j+1})} \right)} \\ &\leq C \sum_{i \in G} \sum_{j=i+1}^{\infty} \frac{2^{sj}}{4^{s(j-i)} \log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(S_{i+1})} \right)} \\ &\leq C \sum_{i \in G} \frac{2^{si}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})} \right)} \sum_{j=i+1}^{\infty} \frac{1}{2^{s(j-i)}} < \infty. \end{aligned}$$

From (4.17) and (4.18) we obtain

$$(4.19) \quad \sum_{i=0}^{\infty} \frac{2^{si}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})} \right)} < \infty.$$

STEP 4.

We here estimate (4.15) to the power s and sum over $i \in \mathbb{N}$ and we infer from (4.19)

$$(4.20) \quad \int_2^{\infty} \frac{t^{s-1}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq t\})} \right)} dt \leq C \sum_{i=0}^{\infty} \frac{2^{si}}{\log_{[\ell]}^{s-1-\alpha} \left(\frac{E\mu(\Omega)}{\mu(\{v \geq 2^{i+1}\})} \right)} < \infty.$$

From (4.20) for $v = v_+$ and $v = v_-$, respectively, we obtain (4.11). Since (4.11) implies (4.12), we are done. □

Acknowledgment. The work is a part of the research project MSM 0021620839 financed by MŠMT. The author would like to thank Stanislav Hencl for fruitful discussions.

REFERENCES

- [1] Brézis H., Wainger S., *A note on limiting case of Sobolev embeddings and convolution inequalities*, Comm. Partial Differential Equations **5** (1980), no. 7, 773–789.
- [2] Černý R., Mašková S., *A sharp form of an embedding into multiple exponential spaces*, Czechoslovak Math. J. **60** (2010), no. 3, 751–782.
- [3] Cianchi A., *A sharp embedding theorem for Orlicz-Sobolev spaces*, Indiana Univ. Math. J. **45** (1996), 39–65.

- [4] Cianchi A., *Optimal Orlicz-Sobolev embeddings*, Rev. Mat. Iberoamericana **20** (2004), 427–474.
- [5] Edmunds D.E., Gurka P., Opic B., *Double exponential integrability of convolution operators in generalized Lorentz-Zygmund spaces*, Indiana Univ. Math. J. **44** (1995), 19–43.
- [6] Edmunds D.E., Gurka P., Opic B., *Double exponential integrability, Bessel potentials and embedding theorems*, Studia Math. **115** (1995), 151–181.
- [7] Edmunds D.E., Gurka P., Opic B., *Sharpness of embeddings in logarithmic Bessel-potential spaces*, Proc. Roy. Soc. Edinburgh Sect. A **126** (1996), 995–1009.
- [8] Edmunds D.E., Kerman R., Pick L., *Optimal Sobolev imbeddings involving rearrangement-invariant quasinorms*, J. Funct. Anal. **170** (2000), no. 2, 307–355.
- [9] Fusco N., Lions P.L., Sbordone C., *Sobolev imbedding theorems in borderline cases*, Proc. Amer. Math. Soc. **124** (1996), 561–565.
- [10] Hajlasz P., Koskela P., *Sobolev met Poincaré*, Memoirs of the Amer. Math. Soc **145** (2000), 101pp.
- [11] Hansson K., *Imbeddings theorems of Sobolev type in potential theory*, Math. Scand. **49** (1979), 77–102.
- [12] Hedberg L.I., *On certain convolution inequalities*, Proc. Amer. Math. Soc. **36** (1972), 505–512.
- [13] Hempel J.A., Morris G.R., Trudinger N.S., *On the sharpness of a limiting case of the Sobolev imbedding theorem*, Bull. Austral. Math. Soc. **3** (1970), 369–373.
- [14] Hencl S., *A sharp form of an embedding into exponential and double exponential spaces*, J. Funct. Anal. **204** (2003), no. 1, 196–227.
- [15] Hencl S., *Sharp generalized Trudinger inequalities via truncation*, J. Math. Anal. Appl. **326** (2006), no. 1, 336–348.
- [16] Koskela P., Onninen J., *Sharp inequalities via truncation*, J. Math. Anal. Appl. **278** (2003), 324–334.
- [17] Maz'ya V., *Sobolev Spaces*, Springer, Berlin, 1975.
- [18] Maz'ya V., *A theorem on multidimensional Schrödinger operator* (Russian), Izv. Akad. Nauk **28** (1964), 1145–1172.
- [19] Malý J., Pick L., *An elementary proof of sharp Sobolev embeddings*, Proc. Amer. Math. Soc. **130** (2002), no. 2, 555–563.
- [20] O'Neil R., *Convolution operators and $L_{(p,q)}$ spaces*, Duke Math. J. **30** (1963), 129–142.
- [21] Peetre J., *Espaces d'interpolation et théorème de Soboleff*, Ann. Inst. Fourier **16** (1966), 279–317.
- [22] Pohozaev S.I., *On the imbedding Sobolev theorem for $pl = n$* , Doklady Conference, Section Math. Moscow Power Inst. (1965), 158–170.
- [23] Rao M.M., Ren Z.D., *Theory of Orlicz Spaces*, Monographs and Textbooks in Pure and Applied Mathematics, 146, Marcel Dekker, New York, 1991.
- [24] Strichartz R.S., *A note on Trudinger's extension of Sobolev's inequality*, Indiana Univ. Math. J. **21** (1972), 841–842.
- [25] Trudinger N.S., *On imbeddings into Orlicz spaces and some applications*, J. Math. Mech. **17** (1967), 473–484.
- [26] Yudovič V.I., *Some estimates connected with integral operators and with solutions of elliptic equations*, Soviet Math. Doklady **2** (1961), 746–749.

CHARLES UNIVERSITY, FACULTY OF MATHEMATICS AND PHYSICS, DEPARTMENT OF MATHEMATICAL ANALYSIS SOKOLOVSKÁ 83, 186 75 PRAHA 8, CZECH REPUBLIC

E-mail: rcerny@karlin.mff.cuni.cz

(Received May 4, 2010, revised October 11, 2010)