

D. Basile

More on κ -Ohio completeness

Commentationes Mathematicae Universitatis Carolinae, Vol. 52 (2011), No. 4, 551--559

Persistent URL: <http://dml.cz/dmlcz/141744>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2011

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

More on κ -Ohio completeness

D. BASILE

Abstract. We study closed subspaces of κ -Ohio complete spaces and, for κ uncountable cardinal, we prove a characterization for them. We then investigate the behaviour of products of κ -Ohio complete spaces. We prove that, if the cardinal κ^+ is endowed with either the order or the discrete topology, the space $(\kappa^+)^{\kappa^+}$ is not κ -Ohio complete. As a consequence, we show that, if κ is less than the first weakly inaccessible cardinal, then neither the space ω^{κ^+} , nor the space \mathbb{R}^{κ^+} is κ -Ohio complete.

Keywords: κ -Ohio complete, compactification, subspace, product

Classification: 54D35, 54B05, 54B10

1. Introduction

All spaces under discussion are Tychonoff. For all undefined notions we refer to [6].

The property of κ -Ohio completeness was introduced in [5] and it is a natural generalization of Ohio completeness, which was introduced by Arhangel'skii in [1] to study remainders in compactifications of topological spaces.

Recall that a topological space X is *κ -Ohio complete* if for every compactification γX of X there exists a G_κ -subset S of γX such that $X \subseteq S$ and for every $y \in S \setminus X$, there is a G_κ -subset of γX that contains y and misses X .

In [5] particular attention was given to sum theorems for κ -Ohio complete spaces. The aim of this paper is focusing on the behaviour that closed subspaces of κ -Ohio complete spaces and products of κ -Ohio complete spaces have. Indeed it is still an open question whether the κ -Ohio completeness property is closed-hereditary or finitely multiplicative.

The paper is divided in two parts. In the first we investigate the behaviour of closed subspaces. Our main result is a characterization of closed subspaces of κ -Ohio complete spaces, for κ uncountable cardinal. In the second part we study products of κ -Ohio complete spaces. We prove that, if the cardinal κ^+ is endowed with either the order or the discrete topology, the space $(\kappa^+)^{\kappa^+}$ is not κ -Ohio complete. From this results it follows that, for a large class of cardinals κ , neither the space ω^{κ^+} nor the space \mathbb{R}^{κ^+} is κ -Ohio complete. For more information see [2].

2. Preliminaries

Following the notation of [4] and [5] we say that a compactification γX of a space X is κ -good for X if there exists a G_κ -subset S of γX such that $X \subseteq S$ and for every $y \in S \setminus X$, there is a G_κ -subset of γX that contains y and misses X . We denote with the symbol $\kappa\mathcal{O}(X)$ the collection of all κ -good compactifications of X . Similarly, we say that a G_κ -subset of a compactification γX of X is a G_κ -good subset for X if it contains X , and if every point of $S \setminus X$ can be separated from X by a G_κ -subset of γX . If $\kappa = \omega$ we omit the symbol ω .

Observe that any space is κ -Ohio complete, for some large enough κ . Recall that the Čech-number of a space X , denoted by $\check{C}(X)$, is the smallest cardinality of a collection \mathcal{U} of open subsets of γX such that $X = \bigcap \mathcal{U}$, where γX is any compactification of X . Therefore, if X is any space, it follows that X is $\check{C}(X)$ -Ohio complete. On the other hand, for every infinite cardinal κ , there exist spaces which are not κ -Ohio complete, as it is shown in the next example (see also [4, Example 5.2]).

Example 2.1. Consider the cardinal κ^+ endowed with the discrete topology and its one point-compactification $\kappa^+ \cup \{\infty\}$. The example is the subspace X of the product $Z = (\kappa^+ \cup \{\infty\}) \times (\kappa^+ \cup \{\infty\})$ where $X = (\kappa^+ \times \kappa^+) \cup \{(\infty, \infty)\}$.

If G is a G_κ -subset of Z that contains the point (∞, ∞) , then $G \cap (Z \setminus X)$ is non-empty, so X is not a G_κ -subset of Z . Similarly, $Z \setminus X$ contains no non-empty G_κ -subset of Z ; this clearly implies that X is not κ -Ohio complete.

It is worth noting that, for a large class of cardinals κ , the space X we have just constructed has a good compactification, even if it is not κ -Ohio complete. Indeed, assume that κ is a non-measurable cardinal number. Then, the cardinal κ^+ is non-measurable as well, and in this case it is well-known that the discrete space of cardinality κ^+ is realcompact (see [6, Exercise 3.11.D(a)]). It follows that, under this hypothesis, the space X is realcompact (see [6, Exercise 3.11.A]), therefore its Čech-Stone compactification βX is good by [6, Theorem 3.11.10].

This means that, for a fixed cardinal κ , if the Čech-Stone compactification of a space X is κ -good for X , the space X need not be κ -Ohio complete. On the other hand, if a space X has a κ -good compactification γX , then the Čech-Stone compactification βX of X is always κ -good for X , as it is shown in the next proposition.

Proposition 2.2. *Let X be a space and let $\gamma X \in \kappa\mathcal{O}(X)$. Then $\{\delta X : \delta X \in \mathcal{C}(X) \text{ and } \delta X \geq \gamma X\} \subseteq \kappa\mathcal{O}(X)$.*

For the simple proof see [3, Proposition 4.3].

3. A characterization of closed subspaces of κ -Ohio complete spaces

In [3] we asked whether closed subspaces of Ohio complete spaces are again Ohio complete. Unfortunately we do not know the answer, as we do not know whether closed subspaces of κ -Ohio complete spaces are again κ -Ohio complete. However, there are some positive results; we will prove them in this section.

Proposition 2.2 asserts that if a space X has a κ -good compactification γX , then every compactification greater than or equal to γX (with respect to the order \leq) is κ -good for X . However, if a space is a closed subspace of a κ -Ohio complete space, then a sort of complementary property holds, as we are going to show. The formulation of the result is new, but it has actually been proved in [3]. We include the proof for completeness sake.

Lemma 3.1. *Let Y be a closed subspace of X . Fix a compactification αX of X and let $\gamma Y = \overline{Y}^{\alpha X}$. Then, for every compactification δY of Y such that $\delta Y \leq \gamma Y$, there exists some compactification ϱX of X such that $\delta Y = \overline{Y}^{\varrho X}$ and $\varrho X \leq \alpha X$.*

PROOF: Fix a compactification δY of Y such that $\delta Y \leq \gamma Y$. Hence, there exists a continuous map $f : \gamma Y \rightarrow \delta Y$ such that $f(y) = y$, for every $y \in Y$. Consider the adjunction space $Z = \alpha X \cup_f \delta Y$. Clearly Z is a compact Hausdorff space, since it is the image of the compact space $\alpha X \oplus \delta Y$ under a closed continuous function, that is, the natural quotient mapping π . Observe that π is closed since f is closed (see for instance [6, p. 94]).

First we shall prove that X , considered as a subspace of Z , has the original topology, by showing that $\pi|X : X \rightarrow \pi(X)$ is a homeomorphism. To verify that $\pi|X$ is one-to-one, pick two different points $x, y \in X$. Observe that, since Y is closed in X , we have $(\gamma Y \setminus Y) \cap X = \emptyset$. There are three different cases to consider. If $x, y \in X \setminus Y$ we have $x, y \in \alpha X \setminus \gamma Y$ and then, by construction, the equivalence classes of x and y are $\{x\}$ and $\{y\}$ respectively. If $x \in X \setminus Y$ and $y \in Y$, the equivalence classes of x and y are $\{x\}$ and $\{y\} \cup f^{-1}(y)$, respectively. Finally, if $x, y \in Y$, the equivalence classes of x and y are $\{x\} \cup f^{-1}(x)$ and $\{y\} \cup f^{-1}(y)$, respectively. In all cases we have $\pi(x) \neq \pi(y)$. This proves that $\pi|X$ is one-to-one.

We will now prove that $\pi|X$ is closed. As we observed before π is closed. Let D be a closed subspace of X , then we may find a closed subset C of $\alpha X \oplus \delta Y$, such that $D = C \cap X$. It follows that $\pi(D) = \pi(C \cap X) = \pi(C) \cap X$ is a closed subset of X . This shows that $\pi|X$ is a homeomorphism.

In a similar way we can prove that δY as a subspace of Z has the original topology. It follows that $\overline{Y}^Z = \delta Y$.

Since the space Z is clearly a compactification of X such that $Z \leq \alpha X$, we are done. □

Given a space X we say that a compactification γX of X is *very κ -good* if $\{\delta X : \delta X \in \mathcal{C}(X) \text{ and } \delta X \leq \gamma X\} \subseteq \kappa\mathcal{O}(X)$. In particular, if γX is a very κ -good compactification for X , then every compactification δX of X such that $\delta X \leq \gamma X$, is very κ -good for X .

Theorem 3.2. *Let Y be a closed subspace of a space X . Assume that X has a very κ -good compactification αX . Then $\gamma Y = \overline{Y}$ (closure in αX) is a very κ -good compactification for Y .*

PROOF: Fix a compactification δY of Y such that $\delta Y \leq \gamma Y$. By Lemma 3.1, there exists a compactification ϱX of X such that $\delta Y = \overline{Y}^{\varrho X}$ and $\varrho X \leq \alpha X$.

Since αX is a very κ -good compactification for X , the compactification ρX is κ -good for X . Let S be a G_κ -subset of ρX that is κ -good for X . Then the set $S \cap \delta Y$ is G_κ -good for Y . This completes the proof. \square

An application of Theorem 3.2 is the following result, which shows that κ -Ohio completeness is hereditary with respect to closed and C^* -embedded subspaces (see also [3]).

Corollary 3.3. *Let Y be a closed C^* -embedded subspace of a κ -Ohio complete space X . Then Y is κ -Ohio complete.*

PROOF: Closures are taken in βX . It follows from Theorem 3.2 that \overline{Y} is a very κ -good compactification for Y . But $\overline{Y} = \beta Y$, by [6, Corollary 3.6.7]. This proves that Y is κ -Ohio complete. \square

Corollary 3.4. *Let Y be a closed subspace of a κ -Ohio complete normal space X . Then Y is κ -Ohio complete.*

If $A \subseteq X$, a continuous function $f : X \rightarrow A$ is called a *retraction* of X onto A , if $f(x) = x$ for all $x \in A$. In this case A is called a *retract* of X .

Corollary 3.5. (1) *Every clopen subspace of a κ -Ohio complete space is κ -Ohio complete.*
 (2) *Every retract of a κ -Ohio complete space is κ -Ohio complete.*

PROOF: This follows from the fact that clopen subspaces and retracts are closed and C^* -embedded subspaces. \square

Unfortunately this does not answer to the following:

Question 3.6. *Is κ -Ohio completeness a closed-hereditary property?*

Theorem 3.2 implies in particular that a closed subspace of a κ -Ohio complete space has some very κ -good compactification. It is pretty natural to ask whether the converse is true, that is, whether, given a space having a very κ -good compactification, it can be embedded as a closed subspace in some κ -Ohio complete space.

The following theorem shows that, if κ is an uncountable cardinal number, the answer is yes.

Theorem 3.7. *Let κ be an uncountable cardinal number. The following statements are equivalent.*

- (1) *Y is a closed subspace of a κ -Ohio complete space X .*
- (2) *There exists a very κ -good compactification γY of Y .*

PROOF: (1) \Rightarrow (2) follows from Theorem 3.2.

(2) \Rightarrow (1). Fix a very κ -good compactification γY of Y . Consider the ordinal space $\omega_1 + 1$ and let Z be the space $(\omega_1 + 1) \times \gamma Y$, and let X be the subspace of Z given by

$$(\omega_1 \times \gamma Y) \cup \{\omega_1\} \times Y.$$

Then Y is clearly a closed subspace of X , so to prove the theorem it suffices to show that X is κ -Ohio complete.

First observe that $\beta X = Z$. Indeed, note that $\beta(\omega_1 \times \gamma Y) = (\omega_1 + 1) \times \gamma Y = Z$. This can be found in [6, Problem 3.12.20(c)]. Since $\omega_1 \times \gamma Y \subseteq X \subseteq Z$, it follows that $\beta X = Z$ by [6, Corollary 3.6.9].

To show that X is κ -Ohio complete, fix a compactification αX of X . Then $\alpha X \leq \beta X = Z$. So we may fix a continuous function $f : Z \rightarrow \alpha X$ such that $f \upharpoonright X$ is the identity on X . We let g be the restriction of f to the set $\{\omega_1\} \times \gamma Y$. Note that since the remainder $\beta X \setminus X$ is contained in the domain of g , it follows that the remainder $\alpha X \setminus X$ is contained in the range of g . So the range of g is given by

$$W = (\omega_1 \times Y) \cup (\alpha X \setminus X).$$

Clearly, the function g witnesses the fact that $W \leq \gamma Y$. By assumption it follows that W is a κ -good compactification for $\{\omega_1\} \times Y$, so we may fix a G_κ -subset S of αX such that every point in $(W \cap S) \setminus (\{\omega_1\} \times Y)$ can be separated from $\{\omega_1\} \times Y$ by a G_κ -subset of W .

Now let $S' = (\omega_1 \times \gamma Y) \cup S$. Since $\omega_1 \times \gamma Y$ is locally compact, it is an open subset of αX and therefore S' is a G_κ -subset of αX . We claim that S' is a G_κ -good subset for X .

So pick an arbitrary point $p \in S' \setminus X$. Then $p \in S \setminus (\{\omega_1\} \times Y)$. By the choice of S , there is a G_κ -subset T of W such that $p \in T$ and $T \cap (\{\omega_1\} \times Y) = \emptyset$. Now note that $\omega_1 \times \gamma Y$ is the union of ω_1 -many compact subspaces and therefore $\alpha X \setminus (\omega_1 \times \gamma Y) = W$ is a G_{ω_1} -subset and hence a G_κ -subset of αX . But then T is also a G_κ -subset of αX . Since T is disjoint from X , this set separates the point p from X . This completes the proof. \square

Question 3.8. *Does the equivalence of Theorem 3.7 also hold for $\kappa = \omega$?*

4. Products of κ -Ohio complete spaces

As we said in the introduction we do not know whether κ -Ohio completeness is finitely multiplicative. Actually, we do not know if even the product of a κ -Ohio complete space with a compact space is again κ -Ohio complete. However, there is some relation between these questions and Question 3.6, as the next theorem shows (see also [3, Theorem 3.4]):

Theorem 4.1. *Let κ be an infinite cardinal number. Consider the following statements.*

- (1) *Preimages of κ -Ohio complete spaces under perfect mappings are κ -Ohio complete.*
- (2) *The product of a κ -Ohio complete space and a compact space is always κ -Ohio complete.*
- (3) *Every closed subspace of a κ -Ohio complete space is κ -Ohio complete.*

Then (1) \Leftrightarrow (2) \Rightarrow (3).

PROOF: To prove that (1) \Rightarrow (2), let X be a κ -Ohio complete space and K be a compact space. Then $\pi_X : X \times K \rightarrow X$ is a perfect mapping, so the hypothesis implies that $X \times K$ is κ -Ohio complete.

For (2) \Rightarrow (3), let Y be a closed subspace of a κ -Ohio complete space X . Consider the product $Z = X \times \beta Y$ and its subspace $\Delta(Y)$. By [6, Theorem 3.6.1], Y is C^* -embedded in βY . From this fact it easily follows that $\Delta(Y)$ is a C^* -embedded copy of Y in Z . Since $\Delta(Y)$ is also closed in Z , by Corollary 3.3 it follows that if Z is κ -Ohio complete then so is Y .

We finally prove that (2) \Rightarrow (1). Since (2) \Rightarrow (3), it follows from [6, Theorem 3.7.26] that (1) holds. □

Therefore, if Question 4.2 below has a positive answer, then Question 3.6 has a positive answer as well.

Question 4.2. *Is the product of a κ -Ohio complete space with a compact space again κ -Ohio complete?*

On the other hand, it is straightforward to see that if a product space is κ -Ohio complete, then each of its factors is κ -Ohio complete as well.

Proposition 4.3. *Let $X = \prod_{\alpha < \tau} X_\alpha$ be a κ -Ohio complete space. Then, for every $\alpha < \tau$, the space X_α is κ -Ohio complete.*

PROOF: Note that every X_α is a retract of X . Now it suffices to apply Corollary 3.5(2). □

The following results show that the product of κ -many κ -Ohio complete spaces has many κ -good compactifications.

Lemma 4.4. *Let $\{X_\alpha : \alpha < \kappa\}$ be a family of spaces. For every $\alpha < \kappa$, let S_α be a G_κ -subset of X_α . Then $\prod_{\alpha < \kappa} S_\alpha$ is a G_κ -subset of $X = \prod_{\alpha < \kappa} X_\alpha$.*

Proposition 4.5. *Let $\{X_\alpha : \alpha < \kappa\}$ be a family of spaces. For every $\alpha < \kappa$, let $\gamma_\alpha X_\alpha \in \kappa\mathcal{O}(X_\alpha)$. Then $\prod_{\alpha < \kappa} \gamma_\alpha X_\alpha \in \kappa\mathcal{O}(\prod_{\alpha < \kappa} X_\alpha)$.*

PROOF: Since $\gamma_\alpha X_\alpha \in \kappa\mathcal{O}(X_\alpha)$, for every $\alpha < \kappa$ there exists a G_κ -subset S_α of $\gamma_\alpha X_\alpha$ which is κ -good with respect to X_α . By Lemma 4.4, the set $\prod_{\alpha < \kappa} S_\alpha$ is a G_κ -subset of $\prod_{\alpha < \kappa} \gamma_\alpha X_\alpha$ that clearly contains $\prod_{\alpha < \kappa} X_\alpha$. We will show that $\prod_{\alpha < \kappa} S_\alpha$ is κ -good with respect to $\prod_{\alpha < \kappa} X_\alpha$.

So, pick a point $p = (p_\alpha)_{\alpha < \kappa} \in \prod_{\alpha < \kappa} S_\alpha \setminus \prod_{\alpha < \kappa} X_\alpha$. Then, for some $\beta < \kappa$, we have $p_\beta \in S_\beta \setminus X_\beta$. Therefore, there exists a G_κ -subset T_β of $\gamma_\beta X_\beta$ containing p_β and missing X_β . The set $Z = \pi_\beta^{-1}(T_\beta)$ is a G_κ -subset of $\prod_{\alpha < \kappa} \gamma_\alpha X_\alpha$ that contains p and misses $\prod_{\alpha < \kappa} X_\alpha$. This proves the proposition. □

The proof of the preceding proposition is based on the fact that the intersection of κ -many G_κ -subsets is again a G_κ -subset. Since this property may fail for larger intersections, we might expect that Proposition 4.5 does not generalize to products with κ^+ -many factors. The next proposition shows that in fact this is the case.

Proposition 4.6. *Let Y be the cardinal κ^+ endowed with either the discrete or the order topology, and consider its one-point compactification $\omega Y = Y \cup \{\infty\}$. Then $(\omega Y)^{\kappa^+}$ is not a κ -good compactification for Y^{κ^+} .*

PROOF: Observe that the point ∞ is not a G_κ -subset of ωY . Hence, Y^{κ^+} is G_κ -dense in $(\omega Y)^{\kappa^+}$. But its remainder $(\omega Y)^{\kappa^+} \setminus Y^{\kappa^+}$ is G_κ -dense in $(\omega Y)^{\kappa^+}$ as well. So $(\omega Y)^{\kappa^+}$ cannot be a κ -good compactification for Y^{κ^+} . \square

Corollary 4.7. *If the cardinal κ^+ is endowed with either the order or the discrete topology, the space $(\kappa^+)^{\kappa^+}$ is not κ -Ohio complete.*

An application of this result is that the limit of an inverse system of κ -Ohio complete spaces need not be κ -Ohio complete.

Proposition 4.8. *The limit of an inverse system of κ -Ohio complete spaces need not be κ -Ohio complete.*

PROOF: If $\alpha < \kappa^+$, then it follows from [9, Proposition 1.10] that $\check{C}((\kappa^+)^\alpha) \leq |\alpha| \leq \kappa$. So it is clear that $(\kappa^+)^\alpha$ is κ -Ohio complete. Now observe that $(\kappa^+)^{\kappa^+}$ can be seen as the inverse limit of the system $\{(\kappa^+)^\alpha, \pi_\beta^\alpha, \kappa^+\}$, where $\pi_\beta^\alpha : (\kappa^+)^\alpha \rightarrow (\kappa^+)^\beta$ is the usual projection. \square

Remark 4.9. Let us remark that the behaviour of the space $(\kappa^+)^{\kappa^+}$ can be different if we consider κ^+ endowed with the discrete or with the order topology. Indeed, if κ^+ has the discrete topology, then, for a large class of cardinals (namely all non-measurable cardinals κ , see [6, Exercise 3.11.D(a)]), the space $(\kappa^+)^{\kappa^+}$ is realcompact and then it has a κ -good compactification.

If we now consider κ^+ with the order topology and we assume that $\kappa = \omega$, then the space $\omega_1^{\omega_1}$ is pseudocompact (see, for example [6, Exercise 3.12.21.(e)]). By a well-known result of Glicksberg ([7]), we have $\beta(\omega_1^{\omega_1}) = (\beta\omega_1)^{\omega_1}$. Since $\beta\omega_1 = \omega_1 + 1$, Proposition 4.6 implies that $\beta(\omega_1^{\omega_1})$ is not a good compactification for $\omega_1^{\omega_1}$. Therefore, it follows by Proposition 2.2, that $\omega_1^{\omega_1}$ does not have any good compactification.

A natural question is then whether the space ω^{κ^+} is or is not κ -Ohio complete. Observe that the argument used in Proposition 4.6 cannot be applied to ω^{κ^+} : every product compactification of ω^{κ^+} is indeed even good. This is a consequence of the next proposition. We will however answer our question in Corollary 4.16 below.

Recall that the compact covering number of a space X , denoted by $\text{kcov}(X)$, is the smallest cardinality of a collection \mathcal{K} of compact subsets of X such that $X = \bigcup \mathcal{K}$. It is well-known and easy to show that for a space X and for any compactification γX of X , we have $\text{kcov}(\gamma X \setminus X) = \check{C}(X)$.

Proposition 4.10. *Let $X = \prod_{\alpha < \kappa} X_\alpha$, where $\text{kcov}(X_\alpha) \leq \lambda$ for every $\alpha < \kappa$, and let $\gamma_\alpha X_\alpha \in \mathcal{C}(X_\alpha)$, for every $\alpha < \kappa$. Then $\prod_{\alpha < \kappa} \gamma_\alpha X_\alpha \in \lambda\mathcal{O}(X)$.*

PROOF: Let $Z = \prod_{\alpha < \kappa} \gamma_\alpha X_\alpha$. We will show that Z itself is the good G_λ -subset we are looking for. Note that, since $\text{kcov}(X_\alpha) \leq \lambda$, the remainder $\gamma_\alpha X_\alpha \setminus X_\alpha$ is a G_λ -subset of $\gamma_\alpha X_\alpha$, for every $\alpha < \kappa$.

Now, fix a point $x = (x_\alpha)_{\alpha < \kappa} \in Z \setminus X$. So, there exists some $\alpha < \kappa$ such that $x_\alpha \in \gamma_\alpha X_\alpha \setminus X_\alpha$. The set $W = \pi_\alpha^{-1}(\gamma_\alpha X_\alpha \setminus X_\alpha)$, is a G_λ -subset of Z that misses X . This completes the proof. \square

Corollary 4.11. *If a product space has σ -compact factors, then any compactification of its product is good.*

This raises the question whether spaces like ω^{κ^+} or \mathbb{R}^{κ^+} are κ -Ohio complete or not. Furthermore it turns out that finding a non κ -good compactification for such spaces is not trivial.

Nevertheless, using Proposition 4.6, which is a very simple but very useful result, we will be able to prove that, if κ is less than the first weakly inaccessible cardinal, neither ω^{κ^+} nor \mathbb{R}^{κ^+} is κ -Ohio complete.

Theorem 4.12. *If X contains a closed copy of the space κ^+ , endowed either with the discrete or the order topology, then X^{κ^+} is not κ -Ohio complete.*

PROOF: Let us prove the theorem assuming that X contains a closed copy of the discrete space of cardinality κ^+ . The other case is analogous. Since X contains a closed copy of κ^+ , the space X^{κ^+} contains a closed copy of $(\kappa^+)^{\kappa^+}$. Assume, striving for a contradiction, that X^{κ^+} is κ -Ohio complete and let $Z = (\gamma X)^{\kappa^+}$, where γX is any compactification of X . Closures are taken in Z .

Our hypothesis, combined with Theorem 3.2, imply that $\overline{(\kappa^+)^{\kappa^+}}$ is a very κ -good compactification for $(\kappa^+)^{\kappa^+}$. Since $\overline{(\kappa^+)^{\kappa^+}} \geq (\omega\kappa^+)^{\kappa^+}$, the latter product is a κ -good compactification for $(\kappa^+)^{\kappa^+}$, which is a contradiction with Proposition 4.6. \square

From the proof of Theorem 4.12 we get the following:

Corollary 4.13. *If X contains a closed copy of the space κ^+ , endowed either with the discrete or the order topology, then no compactification of X^{κ^+} of the form $(\gamma X)^{\kappa^+}$ can be very κ -good for X^{κ^+} .*

Recall that an uncountable cardinal is called weakly inaccessible if it is a regular limit cardinal. We denote by θ the first weakly inaccessible cardinal.

Corollary 4.14. *Assume that $\kappa < \theta$. If X^{κ^+} is κ -Ohio complete, then X is countably compact.*

PROOF: Observe at first that if $\kappa < \theta$, then $\kappa^+ < \theta$. If X were not countably compact, then X^{κ^+} would contain a closed copy of ω^{κ^+} . Since $\kappa^+ < \theta$, the power ω^{κ^+} contains a closed copy of the discrete space κ^+ , by [8]. Then X^{κ^+} would contain a closed copy of κ^+ , which is a contradiction with Theorem 4.12. \square

Question 4.15. *Can we improve Corollary 4.14 substituting ‘countably compact’ by ‘compact’?*

In [3] we showed that the answer is yes for $\kappa = \omega$.

Corollary 4.16. *If $\kappa < \theta$, then neither ω^{κ^+} nor \mathbb{R}^{κ^+} is κ -Ohio complete.*

Corollary 4.17. *If $\kappa < \theta$, then no compactification of ω^{κ^+} (resp. \mathbb{R}^{κ^+}) of the form Z^{κ^+} is very κ -good for ω^{κ^+} (resp. \mathbb{R}^{κ^+}).*

Question 4.18. *Let $\kappa < \theta$. Does exist some very κ -good compactification for ω^{κ^+} (resp. \mathbb{R}^{κ^+})?*

By Theorem 3.7 this question is equivalent to the question whether, if κ^+ is strictly less than the first weakly inaccessible cardinal, the space ω^{κ^+} (resp. \mathbb{R}^{κ^+}) can be embedded as a closed subspace in some κ -Ohio complete space. Moreover, let us point out that if Question 4.15 has a positive answer, then Question 4.18 has a negative answer.

Actually, to answer in the negative to Question 4.18 it would be enough to show that the space $(\kappa^+)^{\kappa^+}$, where κ^+ is endowed with the discrete topology does not have any very κ -good compactification. Unfortunately we do not know the answer to this.

REFERENCES

- [1] Arhangel'skii A.V., *Remainders in compactifications and generalized metrizable properties*, Topology Appl. **150** (2005), 79–90.
- [2] Basile D., *κ -Ohio completeness and related problems*, Doctoral Thesis, Vrije Universiteit, Amsterdam, 2009.
- [3] Basile D., van Mill J., *Ohio completeness and products*, Topology Appl. **155** (2008), no. 4, 180–189.
- [4] Basile D., van Mill J., Ridderbos G.J., *Sum theorems for Ohio completeness*, Colloq. Math. **113** (2008), 91–104.
- [5] Basile D., van Mill J., Ridderbos G.J., *κ -Ohio completeness*, J. Math. Soc. Japan **61** (2009), no. 4, 1293–1301.
- [6] Engelking R., *General Topology*, second ed., Heldermann, Berlin, 1989.
- [7] Glicksberg I., *Stone-Čech compactifications of products*, Trans. Amer. Math. Soc. **90** (1959), 369–382.
- [8] Mycielski J., *α -incompactness of N^α* , Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys. **12** (1964), 437–438.
- [9] Okunev O., Tamariz-Mascarúa A., *On the Čech number of $C_p(X)$* , Topology Appl. **137** (2004), no. 1–3, 237–249; IV Iberoamerican Conference on Topology and its Applications.

UNIVERSITÀ DEGLI STUDI DI CATANIA, DIPARTIMENTO DI MATEMATICA E INFORMATICA, VIALE ANDREA DORIA 6, 95125 CATANIA, ITALY

E-mail: basile@dmi.unict.it

(Received May 25, 2011)