

Applications of Mathematics

Yong-Fu Yang

Global classical solutions to a kind of mixed initial-boundary value problem for inhomogeneous quasilinear hyperbolic systems

Applications of Mathematics, Vol. 57 (2012), No. 3, 231–261

Persistent URL: <http://dml.cz/dmlcz/142852>

Terms of use:

© Institute of Mathematics AS CR, 2012

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

GLOBAL CLASSICAL SOLUTIONS TO A KIND OF MIXED
INITIAL-BOUNDARY VALUE PROBLEM FOR INHOMOGENEOUS
QUASILINEAR HYPERBOLIC SYSTEMS

YONG-FU YANG, Nanjing

(Received November 30, 2009)

Abstract. In this paper, the mixed initial-boundary value problem for inhomogeneous quasilinear strictly hyperbolic systems with nonlinear boundary conditions in the first quadrant $\{(t, x) : t \geq 0, x \geq 0\}$ is investigated. Under the assumption that the right-hand side satisfies a matching condition and the system is strictly hyperbolic and weakly linearly degenerate, we obtain the global existence and uniqueness of a C^1 solution and its L^1 stability with certain small initial and boundary data.

Keywords: quasilinear hyperbolic system, mixed initial-boundary value problem, global classical solution, weak linear degeneracy, matching condition

MSC 2010: 35L50

1. INTRODUCTION AND MAIN RESULTS

Consider the first order strictly quasilinear hyperbolic system

$$(1.1) \quad \frac{\partial u}{\partial t} + A(u) \frac{\partial u}{\partial x} = B(u),$$

where $u = (u_1, \dots, u_n)^\top$ is the unknown vector function of (t, x) , $A(u)$ is an $n \times n$ matrix with suitably smooth elements $a_{ij}(u)$ ($i, j = 1, \dots, n$), and $B(u) = (B_1(u), \dots, B_n(u))^\top$ is a vector function of u with suitably smooth elements.

The author's research was in part supported by National Natural Science Foundation of China under Grant No. 10926162, Fundamental Research Funds for the Central Universities under Grant No. 2009B01314, and Natural Science Foundation of Hohai university under Grant No. 2009428011.

By the definition of strict hyperbolicity, for any given u on the domain under consideration, $A(u)$ has n distinct real eigenvalues $\lambda_1(u), \dots, \lambda_n(u)$. We furthermore suppose that

$$(1.2) \quad \lambda_1(0) < \dots < \lambda_m(0) < 0 < \lambda_{m+1}(0) < \dots < \lambda_n(0).$$

Let $l_i(u) = (l_{i1}(u), \dots, l_{in}(u))$ and $r_i(u) = (r_{i1}(u), \dots, r_{in}(u))^\top$ be respectively a left and right eigenvector corresponding to $\lambda_i(u)$ ($i = 1, \dots, n$):

$$(1.3) \quad l_i(u)A(u) = \lambda_i(u)l_i(u) \quad \text{and} \quad A(u)r_i(u) = \lambda_i(u)r_i(u).$$

We have

$$(1.4) \quad \det|l_{ij}(u)| \neq 0 \quad (\text{equivalently, } \det|r_{ij}(u)| \neq 0).$$

Without loss of generality, we suppose that

$$(1.5) \quad l_i(u)r_j(u) \equiv \delta_{ij} \quad (i, j = 1, \dots, n),$$

$$(1.6) \quad r_i(u)^\top r_i(u) \equiv 1 \quad (i = 1, \dots, n),$$

where δ_{ij} stands for Kronecker's symbol.

All $\lambda_i(u)$, $l_{ij}(u)$ and $r_{ij}(u)$ ($i, j = 1, \dots, n$) are supposed to have the same regularity as $a_{ij}(u)$, ($i, j = 1, \dots, n$).

Under the assumption that

$$(1.7) \quad B(u) \equiv 0,$$

for the Cauchy problem of system (1.1) with the initial data

$$(1.8) \quad t = 0: u = \varphi(x), \quad x \in \mathbb{R},$$

where $\varphi(x)$ is a C^1 vector value function with bounded C^1 norm, many results concerning global existence and blow-up of classical solutions have been obtained (see [1], [5]–[6], and [16]). In particular, by means of the concept of weak linear degeneracy, for small initial data with certain decaying properties, the global existence and the blow-up phenomenon of a C^1 solution to the Cauchy problem (1.1) and (1.8) have been completely studied (see [9] and [14]–[15]). In virtue of two basic L^1 estimates, Zhou [20] furthermore relaxed the restrictions on the initial data and then showed that the Cauchy problem (1.1) and (1.8) for the weakly linearly degenerate and strictly hyperbolic system admits a unique C^1 solution which also satisfies L^1 stability.

In order to consider the effect of nonlinear boundary conditions on the global regularity of a classical solution of system (1.1), Li & Wang [12] investigated the mixed initial-boundary value problem for system (1.1) on a half-unbounded domain. The result obtained by Li & Wang indicates that the interaction of nonlinear boundary conditions with nonlinear hyperbolic waves does not cause any negative effect on the global regularity of the C^1 solution, provided that the C^1 norms of initial and boundary data both decaying at infinity are small enough. Recently, Zhou & Yang [21] also relaxed the restrictions on the initial and boundary data and then established results on the global C^1 solution for linearly degenerate and weakly linearly degenerate system, respectively.

For inhomogeneous quasilinear hyperbolic system (1.1), under the assumptions that $B(u)$ satisfies the so-called matching condition and the system is strictly hyperbolic and weakly linearly degenerate, Kong [7] and Li [8] established the corresponding results for the Cauchy problem (1.1) and (1.8). Recently, Wu [19] extended the results established by Zhou [20] to the inhomogeneous case. Chen [3] also obtained the corresponding inhomogeneous result in [12].

In this paper, we are going to re-prove the global existence result with less restrictions on the initial and boundary data. In particular, the supreme norms of the derivatives of the initial and boundary data are not assumed to be small. We shall also obtain global L^1 stability results in this situation.

Moreover, for the mixed initial-boundary value problem on a bounded domain $\{(t, x): t \geq 0, 0 \leq x \leq L\}$, the results on the global regularity can be found in [4], [9]–[11], and [18].

On the domain

$$(1.9) \quad D \stackrel{\text{def}}{=} \{(t, x): t \geq 0, x \geq 0\}$$

we consider the mixed initial-boundary value problem for system (1.1) with the initial data

$$(1.10) \quad t = 0: u = \varphi(x), \quad x \geq 0$$

and the boundary condition

$$(1.11) \quad x = 0: v_s = f_s(\alpha(t), v_1, \dots, v_m) + h_s(t) \quad (s = m + 1, \dots, n),$$

where

$$(1.12) \quad v_i = l_i(u)u \quad (i = 1 \dots, n),$$

$h_s(t)$ ($s = m + 1, \dots, n$) are given C^1 functions of t and

$$\alpha(t) \stackrel{\text{def}}{=} (\alpha_1(t), \dots, \alpha_k(t)).$$

Let

$$h(t) \stackrel{\text{def}}{=} (h_{m+1}(t), \dots, h_n(t)),$$

and let us define

$$|u| \stackrel{\text{def}}{=} \left(\sum_{k=1}^n u_k^2 \right)^{1/2}$$

for any vector value function $u = (u_1, \dots, u_n)^\top$. Without loss of generality, we suppose that

$$(1.13) \quad f_s(\alpha(t), 0, \dots, 0) \equiv 0 \quad (s = m + 1, \dots, n).$$

We also remark that in a neighborhood of $u = 0$, the boundary condition (1.11) takes the same form independently of the choice of left eigenvectors.

To state our results precisely, we shall first recall the concept of weak linear degeneracy (see [9], [14]) and matching condition (see [7], [8], [19]) as follows.

Definition 1.1. The i th characteristic $\lambda_i(u)$ is weakly linearly degenerate if along the i th characteristic trajectory $u = u^{(i)}(s)$ passing through $u = 0$, defined by

$$(1.14) \quad \frac{du}{ds} = r_i(u),$$

$$(1.15) \quad s = 0: u = 0,$$

we have

$$(1.16) \quad \nabla \lambda_i(u) r_i(u) \equiv 0, \quad \forall |u| \text{ small},$$

namely,

$$(1.17) \quad \lambda_i(u^{(i)}(s)) \equiv \lambda_i(0), \quad \forall |s| \text{ small}.$$

If all characteristics $\lambda_i(u)$ ($i = 1, \dots, n$) are weakly linearly degenerate, then the system (1.1) is said to be weakly linearly degenerate.

Definition 1.2. $B(u)$ satisfies the matching condition if along all characteristic trajectories passing through $u = 0$ we have $B(u) \equiv 0$, $\forall |u|$ small, i.e.,

$$(1.18) \quad B(u^{(i)}(s)) \equiv 0, \quad \forall |s| \text{ small} \quad (i = 1, \dots, n).$$

In this case, it is easy to see that

$$(1.19) \quad B(0) = 0, \quad \nabla B(0) = 0.$$

Our main results can be summarized as follows:

Theorem 1.1. *Suppose that in a neighborhood of $u = 0$ we have $A(u) \in C^2$, system (1.1) is strictly hyperbolic and weakly linearly degenerate, and $B(u) \in C^2$ satisfies the matching condition. Suppose furthermore that φ , α , f_s , and $h_s \in C^1$ ($s = m + 1, \dots, n$). Suppose finally that the conditions of C^1 compatibility are satisfied at the point $(0, 0)$ and assumptions (1.2) and (1.13) hold. Let*

$$(1.20) \quad M = \max \left\{ \sup_{x \geq 0} |\varphi'(x)|, \sup_{t \geq 0} |\alpha'(t)|, \sup_{t \geq 0} |h'(t)| \right\}.$$

Then there exists a positive constant ε independent of M such that the mixed initial-boundary value problem (1.1) and (1.10)–(1.11) admits a unique global C^1 solution $u = u(t, x)$ on the domain D provided that

$$(1.21) \quad \int_0^\infty |\varphi'(x)| dx \leq \varepsilon, \quad \int_0^\infty |\varphi(x)| dx \leq \frac{\varepsilon}{M+1}$$

and

$$(1.22) \quad \int_0^\infty (|h'(t)| + |\alpha'(t)|) dt \leq \varepsilon, \quad \int_0^\infty (|h(t)| + |\alpha(t)|) dt \leq \frac{\varepsilon}{M+1}.$$

Theorem 1.2. *Under the assumptions of Theorem 1.1, suppose furthermore that $B(u) \in C^3$. If $u^{(1)}, u^{(2)}$ are two solutions given by Theorem 1.1 with initial data $\varphi^{(1)}, \varphi^{(2)}$, boundary conditions $\alpha^{(1)}, h^{(1)}$ and $\alpha^{(2)}, h^{(2)}$, respectively, then we have*

$$(1.23) \quad \int_0^\infty |u^{(1)}(t, x) - u^{(2)}(t, x)| dx \\ \leq C \left(\int_0^\infty |\varphi^{(1)}(x) - \varphi^{(2)}(x)| dx + \int_0^\infty |h^{(1)}(t) - h^{(2)}(t)| dt \right. \\ \left. + \int_0^\infty |\alpha^{(1)}(t) - \alpha^{(2)}(t)| dt \right), \quad \forall t \geq 0,$$

where C is a positive constant independent of M and t .

Under the assumption that (1.7) holds, for the weak solution to the Cauchy problem (1.1) and (1.8) for general quasilinear hyperbolic systems, Bressan et al. [2] and Liu & Yang [17] both obtained global L^1 stability with respect to time t .

This paper is organized as follows: In Section 2, we recall and generalize John's formula on the decomposition of waves. Section 3 is devoted to two basic lemmas concerning the L^1 estimate. In Sections 4 and 5, we prove Theorem 1.1 and Theorem 1.2, respectively.

2. PRELIMINARIES

Suppose that $A(u) \in C^2$. By Lemma 2.5 in [14] (see also [9]), there exists an invertible C^3 transformation $u = u(\tilde{u})$ ($u(0) = 0$) such that in the \tilde{u} -space, for each $i = 1, \dots, n$, the i th characteristic trajectory passing through $\tilde{u} = 0$ coincides with the \tilde{u}_i -axis at least for $|\tilde{u}_i|$ small, namely

$$(2.1) \quad \tilde{r}_i(\tilde{u}_i e_i) \equiv e_i, \quad \forall |\tilde{u}_i| \text{ small} \quad (i = 1, \dots, n),$$

where

$$(2.2) \quad e_i = (0, \dots, 0, \overset{(i)}{1}, 0, \dots, 0)^\top \in \mathbb{R}^n \quad (i = 1, \dots, n).$$

Such a transformation is called a normalized transformation and the corresponding unknown variables $\tilde{u} = (\tilde{u}_1, \dots, \tilde{u}_n)^\top$ are called normalized variables or normalized coordinates.

Let v_i ($i = 1, \dots, n$) be defined by (1.12) and

$$(2.3) \quad w_i = l_i(u)u_x \quad (i = 1, \dots, n),$$

$$(2.4) \quad \beta_i = l_i(u)B(u) \quad (i = 1, \dots, n).$$

By (1.5), we have

$$(2.5) \quad u = \sum_{k=1}^n v_k r_k(u)$$

and

$$(2.6) \quad u_x = \sum_{k=1}^n w_k r_k(u).$$

Let

$$(2.7) \quad \frac{d}{d_i t} = \frac{\partial}{\partial t} + \lambda_i(u) \frac{\partial}{\partial x}$$

be the directional derivative with respect to time t along the i th characteristic. We have (see [19])

$$(2.8) \quad \frac{du}{d_i t} = \sum_{\substack{k=1 \\ k \neq i}}^n (\lambda_i(u) - \lambda_k(u)) w_k r_k(u) + \sum_{h=1}^n \beta_h r_h(u).$$

It is easy to see, in normalized coordinates, that

$$(2.9) \quad \frac{du_i}{dt} = \sum_{j,k=1}^n \varrho_{ijk}(u) u_j w_k + \sum_{j=1}^n \left(\sum_{h=1}^n \bar{\varrho}_{ijh}(u) \beta_h(u) \right) u_j + r_{ii}(u) \beta_i,$$

where

$$(2.10) \quad \varrho_{ijk} = \begin{cases} (\lambda_i(u) - \lambda_k(u)) \int_0^1 \frac{\partial r_{ki}(su_1, \dots, su_{k-1}, u_k, su_{k+1}, \dots, su_n)}{\partial u_j} ds, & k \neq i, j \neq k, \\ 0 & \text{otherwise,} \end{cases}$$

$$(2.11) \quad \bar{\varrho}_{ijh} = \begin{cases} \int_0^1 \frac{\partial r_{hi}(su_1, \dots, su_{h-1}, u_h, su_{h+1}, \dots, su_n)}{\partial u_j} ds, & h \neq i, j \neq h, \\ 0 & \text{otherwise.} \end{cases}$$

From (2.6) and (2.9), we get

$$(2.12) \quad \frac{\partial u_i}{\partial t} + \frac{\partial(\lambda(u)u_i)}{\partial x} = \sum_{j,k=1}^n F_{ijk}(u) u_j w_k + \sum_{j=1}^n \left(\sum_{h=1}^n \bar{\varrho}_{ijh}(u) \beta_h(u) \right) u_j + r_{ii}(u) \beta_i,$$

or equivalently,

$$(2.13) \quad d[u_i(dx - \lambda_i(u) dt)] \\ = \left[\sum_{j,k=1}^n F_{ijk}(u) u_j w_k + \sum_{j=1}^n \left(\sum_{h=1}^n \bar{\varrho}_{ijh}(u) \beta_h(u) \right) u_j + r_{ii}(u) \beta_i \right] dt \wedge dx,$$

where

$$(2.14) \quad F_{ijk}(u) = \varrho_{ijk}(u) + \nabla \lambda_i(u) r_k(u) \delta_{ij}.$$

By (2.10), it obviously follows that

$$(2.15) \quad F_{ijj}(u) \equiv 0, \quad \forall j \neq i.$$

Moreover, when system (1.1) is weakly linearly degenerate, then in normalized coordinates we have

$$(2.16) \quad F_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j|, \quad \forall i, j \in \{1, \dots, n\}.$$

On the other hand, we have

$$(2.17) \quad \frac{dw_i}{dt} = \sum_{j,k=1}^n \gamma_{ijk}(u)w_jw_k + \sum_{j=1}^n \left(\sum_{h=1}^n B_{ijh}(u)\beta_h(u) + \nu_{ij}(u) \right) w_j,$$

where

$$(2.18) \quad \gamma_{ijk}(u) = \frac{1}{2}\{(\lambda_j(u) - \lambda_k(u))l_i(u)\nabla r_k(u)r_j(u) - \nabla\lambda_k(u)r_j(u)\delta_{ik} + (j|k)\},$$

$$(2.19) \quad B_{ijh}(u) = -l_i(u)\nabla r_j(u)r_h(u),$$

and

$$(2.20) \quad \nu_{ij}(u) = l_i(u)\nabla B(u)r_j(u),$$

with $(j|k)$ standing for all terms obtained by changing j and k in the previous terms. Hence, we have

$$(2.21) \quad \gamma_{ijj}(u) \equiv 0, \quad \forall j \neq i.$$

When system (1.1) is weakly linearly degenerate, in normalized coordinates we get

$$(2.22) \quad \gamma_{iii}(u_i e_i) \equiv 0, \quad \forall |u_i| \quad (i = 1, \dots, n)$$

and

$$(2.23) \quad B_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j| \quad (i, j = 1, \dots, n).$$

Similarly to (2.13), we obtain

$$(2.24) \quad \frac{\partial w_i}{\partial t} + \frac{\partial(\lambda_i(u)w_i)}{\partial x} = \sum_{j,k=1}^n \Gamma_{ijk}(u)w_jw_k + \sum_{j=1}^n \left(\sum_{h=1}^n B_{ijh}(u)\beta_h(u) + \nu_{ij}(u) \right) w_j,$$

or equivalently,

$$(2.25) \quad d[w_i(dx - \lambda_i(u) dt)] \\ = \left[\sum_{j,k=1}^n \Gamma_{ijk}(u)w_jw_k + \sum_{j=1}^n \left(\sum_{h=1}^n B_{ijh}(u)\beta_h(u) + \nu_{ij}(u) \right) w_j \right] dt \wedge dx,$$

where

$$(2.26) \quad \Gamma_{ijk}(u) = \frac{1}{2}(\lambda_j(u) - \lambda_k(u))l_i(u)[\nabla r_k(u)r_j(u) - \nabla r_j(u)r_k(u)].$$

Thus, we have

$$(2.27) \quad \Gamma_{ijj}(u) \equiv 0, \quad \forall i, j \in \{1, \dots, n\}.$$

As for (2.9) and (2.17), we have

Lemma 2.1 ([19]). *Suppose that $A(u) \in C^2$ in a neighborhood of $u = 0$ and system (1.1) is strictly hyperbolic. Suppose furthermore that $B(u) \in C^2$ satisfies the matching condition. Then, in normalized coordinates, we have*

$$(2.28) \quad \begin{aligned} & \sum_{j=1}^n \left(\sum_{h=1}^n \bar{\varrho}_{ijh}(u) \beta_h(u) \right) u_j + r_{ii}(u) \beta_i(u) \\ &= \sum_{j,k=1}^n P_{ijk}(u) u_j u_k, \quad \forall |u| \text{ small} \quad (i = 1, \dots, n), \end{aligned}$$

where $P_{ijk}(u)$ are C^1 functions with respect to their arguments in a neighborhood of $u = 0$ and

$$(2.29) \quad P_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j| \text{ small} \quad (i, j = 1, \dots, n).$$

In addition, we get

$$(2.30) \quad \begin{aligned} & \sum_{j=1}^n \left(\sum_{h=1}^n B_{ijh}(u) \beta_h(u) + \nu_{ij}(u) \right) w_j \\ &= \sum_{j,k=1}^n Q_{ijk}(u) u_k w_j, \quad \forall |u| \text{ small} \quad (i = 1, \dots, n), \end{aligned}$$

where $Q_{ijk}(u)$ are C^1 functions with respect to their arguments in a neighborhood of $u = 0$ and

$$(2.31) \quad Q_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j| \text{ small} \quad (i, j = 1, \dots, n).$$

By Lemma 2.1, (2.9) and (2.12) can be rewritten as

$$(2.32) \quad \frac{du_i}{dt} = \sum_{j,k=1}^n \varrho_{ijk}(u) u_j w_k + \sum_{j,k=1}^n P_{ijk}(u) u_j u_k,$$

$$(2.33) \quad \frac{\partial u_i}{\partial t} + \frac{\partial(\lambda_i(u) u_i)}{\partial x} = \sum_{j,k=1}^n F_{ijk}(u) u_j w_k + \sum_{j,k=1}^n P_{ijk}(u) u_j u_k.$$

By Lemma 2.1, (2.17) and (2.24) can also be rewritten as

$$(2.34) \quad \frac{dw_i}{dt} = \sum_{j,k=1}^n \gamma_{ijk}(u) w_j w_k + \sum_{j,k=1}^n Q_{ijk}(u) u_k w_j,$$

$$(2.35) \quad \frac{\partial w_i}{\partial t} + \frac{\partial(\lambda_i(u) w_i)}{\partial x} = \sum_{j,k=1}^n \Gamma_{ijk}(u) w_j w_k + \sum_{j,k=1}^n Q_{ijk}(u) u_k w_j.$$

We next present a formula on the decomposition of waves for the difference of two solutions $u^{(1)}$, $u^{(2)}$ to system (1.1) (see [19], [20]). Let

$$(2.36) \quad w_i^{(1)} = l_i(u^{(1)})u_x^{(1)}, \quad w_i^{(2)} = l_i(u^{(2)})u_x^{(2)} \quad (i = 1, \dots, n).$$

By (1.5), we have

$$(2.37) \quad u_x^{(1)} = \sum_{i=1}^n w_i^{(1)} r_i(u^{(1)}), \quad u_x^{(2)} = \sum_{i=1}^n w_i^{(2)} r_i(u^{(2)}).$$

Let

$$(2.38) \quad u^{(0)} = u^{(1)} - u^{(2)},$$

$$(2.39) \quad \xi_i^{(1)} = l_i(u^{(1)})u^{(0)}, \quad \xi_i^{(2)} = l_i(u^{(2)})u^{(0)},$$

then we get

$$(2.40) \quad u^{(0)} = \sum_{j=1}^n \xi_j^{(1)} r_j(u^{(1)}) = \sum_{j=1}^n \xi_j^{(2)} r_j(u^{(2)}).$$

Thus,

$$(2.41) \quad \begin{aligned} & \xi_{it}^{(1)} + (\lambda_i(u^{(1)})\xi_i^{(1)})_x \\ &= \sum_{j,k=1}^n \Theta_{ijk}(u^{(1)})\xi_j^{(1)} w_k^{(1)} - (\lambda_i(u^{(1)}) - \lambda_i(u^{(2)}))l_i(u^{(1)})r_i(u^{(2)})w_i^{(2)} \\ & \quad + \sum_{j \neq i} (\lambda_i(u^{(1)}) - \lambda_j(u^{(2)}))l_i(u^{(1)})(r_j(u^{(1)}) - r_j(u^{(2)}))w_j^{(2)} \\ & \quad - \sum_{j=1}^n l_i(u^{(1)})\nabla r_j(u^{(1)})B(u^{(1)})\xi_j^{(1)} + l_i(u^{(1)})(B(u^{(1)}) - B(u^{(2)})) \end{aligned}$$

and

$$(2.42) \quad \begin{aligned} & \xi_{it}^{(2)} + (\lambda_i(u^{(2)})\xi_i^{(2)})_x \\ &= \sum_{j,k=1}^n \Theta_{ijk}(u^{(2)})\xi_j^{(2)} w_k^{(2)} + (\lambda_i(u^{(1)}) - \lambda_i(u^{(2)}))l_i(u^{(2)})r_i(u^{(1)})w_i^{(1)} \\ & \quad - \sum_{j \neq i} (\lambda_i(u^{(2)}) - \lambda_j(u^{(1)}))l_i(u^{(2)})(r_j(u^{(1)}) - r_j(u^{(2)}))w_j^{(1)} \\ & \quad - \sum_{j=1}^n l_i(u^{(2)})\nabla r_j(u^{(2)})B(u^{(2)})\xi_j^{(2)} - l_i(u^{(2)})(B(u^{(1)}) - B(u^{(2)})), \end{aligned}$$

where

$$(2.43) \quad \Theta_{ijk}(u) = (\lambda_k(u) - \lambda_i(u))l_i(u)\nabla r_j(u)r_k(u) + \nabla\lambda_i(u)r_k(u)\delta_{ij}.$$

We have, in normalized coordinates, that

$$(2.44) \quad \Theta_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j| \text{ small}, \quad \forall j \neq i.$$

When system (1.1) is weakly linearly degenerate, in normalized coordinates we get

$$(2.45) \quad \Theta_{ijj}(u_j e_j) \equiv 0, \quad \forall |u_j| \text{ small} \quad (i, j = 1, \dots, n).$$

3. TWO BASIC L^1 ESTIMATES

In this section, we present two basic L^1 estimates (see [21]).

Lemma 3.1. *Let $\varphi = \varphi(t, x) \in C^1$ satisfy*

$$(3.1) \quad \varphi_t + (\lambda(t, x)\varphi)_x = F, \quad 0 \leq t \leq T, \quad x \geq 0,$$

$$(3.2) \quad t = 0: \varphi = \varphi_0, \quad x \geq 0,$$

where $\lambda \in C^1$. Then for any $t \in [0, T]$,

(i) if $\lambda \geq 0$, we have

$$(3.3) \quad \|\varphi(t, \cdot)\|_{L^1(\mathbb{R}^+)} \leq \|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^t \|F(s, \cdot)\|_{L^1(\mathbb{R}^+)} ds + \int_0^t \lambda(s, 0)|\varphi(s, 0)| ds,$$

(ii) if $\lambda \leq 0$, we have

$$(3.4) \quad \begin{aligned} \|\varphi(t, \cdot)\|_{L^1(\mathbb{R}^+)} + \int_0^t (-\lambda(s, 0))|\varphi(s, 0)| ds \\ \leq \|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^t \|F(s, \cdot)\|_{L^1(\mathbb{R}^+)} ds, \end{aligned}$$

where $\|\varphi(t, \cdot)\|_{L^1(\mathbb{R}^+)} = \int_0^\infty |\varphi(t, x)| dx$.

Lemma 3.2. Let $\varphi = \varphi(t, x)$ and $\psi = \psi(t, x)$ be C^1 functions satisfying

$$(3.5) \quad \varphi_t + (\lambda(t, x)\varphi)_x = F, \quad 0 \leq t \leq T, \quad x \geq 0,$$

$$(3.6) \quad t = 0: \varphi = \varphi_0(x), \quad x \geq 0$$

and

$$(3.7) \quad \psi_t + (\mu(t, x)\psi)_x = G, \quad 0 \leq t \leq T, \quad x \geq 0,$$

$$(3.8) \quad t = 0: \psi = \psi_0(x), \quad x \geq 0,$$

where $\lambda, \mu \in C^1$ such that there exists a positive constant δ_0 independent of T verifying

$$(3.9) \quad \mu(t, x) - \lambda(t, x) \geq \delta_0, \quad 0 \leq t \leq T, \quad x \geq 0.$$

Then, for any $t \in [0, T]$,

(i) if $\lambda \geq 0$ and $\mu > 0$, we have

$$(3.10) \quad \delta_0 \int_0^T \int_0^\infty |\varphi(t, x)| \cdot |\psi(t, x)| \, dx \, dt \\ \leq 4 \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt + \int_0^T \lambda(t, 0) |\varphi(t, 0)| \, dt \right) \\ \times \left(\|\psi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt + \int_0^T \mu(t, 0) |\psi(t, 0)| \, dt \right),$$

(ii) if $\lambda \leq 0$ and $\mu \geq 0$, we have

$$(3.11) \quad \delta_0 \int_0^T \int_0^\infty |\varphi(t, x)| \cdot |\psi(t, x)| \, dx \, dt \\ \leq 4 \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right) \\ \times \left(\|\psi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt + \int_0^T \mu(t, 0) |\psi(t, 0)| \, dt \right),$$

(iii) if $\lambda < 0$ and $\mu \leq 0$, we have

$$(3.12) \quad \delta_0 \int_0^T \int_0^\infty |\varphi(t, x)| \cdot |\psi(t, x)| \, dx \, dt \\ \leq 4 \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right) \\ \times \left(\|\psi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right).$$

4. PROOF OF THEOREM 1.1

By the existence and uniqueness of the local C^1 solution to mixed initial-boundary value problem (1.1) and (1.10)–(1.11), in order to prove Theorem 1.1, it suffices to establish uniform a priori estimates for the C^0 norms of u and $\partial u/\partial x$ on the existence domain of the C^1 solution $u = u(t, x)$ (see [13]).

By (1.2), there exist positive constants δ and δ_0 small enough such that

$$(4.1) \quad \lambda_{j+1}(u) - \lambda_j(\bar{u}) \geq \delta_0, \quad \forall |u|, |\bar{u}| \leq \delta \quad (j = 1, \dots, n-1).$$

For the time being it is supposed that on the existence domain of the C^1 solution $u = u(t, x)$ to the mixed initial-boundary value problem (1.1) and (1.10)–(1.11) we have

$$(4.2) \quad |u(t, x)| \leq \delta.$$

At the end of the proof of Lemma 4.2, we shall explain that this hypothesis is reasonable. Thus, in order to prove Theorem 1.1, we only need to establish uniform a priori estimates for the supreme norms of u and $w = (w_1, \dots, w_n)^\top$ defined by (2.3) on any given time interval $[0, T]$.

By the assumptions $\int_0^\infty |\varphi'(x)| dx < \infty$ and $\int_0^\infty |\varphi(x)| dx < \infty$, it is easy to see that

$$(4.3) \quad \lim_{x \rightarrow \infty} \varphi(x) = 0.$$

Due to finite propagation speed of waves, we have

$$(4.4) \quad \lim_{x \rightarrow \infty} u(t, x) = 0, \quad \forall t \in [0, T].$$

Thus, it follows that

$$(4.5) \quad u(t, x) = - \int_x^\infty u_y(t, y) dy.$$

Differentiating the boundary condition (1.11) with respect to time t and taking into account (1.2) and (4.2), for $\delta > 0$ small enough we have (see [12])

$$(4.6) \quad \begin{aligned} x = 0: w_s = & \sum_{r=1}^m f_{sr}(\alpha(t), u) w_r + \sum_{i=1}^k \bar{f}_{si}(\alpha(t), u) \alpha'_i(t) \\ & + \sum_{\bar{s}=m+1}^n \tilde{f}_{s\bar{s}}(\alpha(t), u) h'_{\bar{s}}(t) \quad (s = m+1, \dots, n), \end{aligned}$$

where f_{sr} , \bar{f}_{si} , and $\tilde{f}_{s\bar{s}}$ are continuous functions with respect to their arguments.

By Lemma 2.5 in [14], there exists a normalized transformation. Without loss of generality, we assume that $u = (u_1, \dots, u_n)^\top$ are already normalized variables. To consider the mixed initial-boundary value problem in normalized coordinates, we need the following lemma.

Lemma 4.1 ([12]). *The boundary condition (1.11) and the assumption (1.22) keep the same form under any given smooth invertible transformation $u = u(\tilde{u})$ ($u(0) = 0$).*

Moreover, in normalized coordinates, it is easy to see that

$$(4.7) \quad v_i = l_i(u)u = l_i(0)u + o(|u|) = u_i + o(|u|) \quad (i = 1, \dots, n).$$

Substituting (4.7) into (1.11), the Implicit Function Theorem implies, for sufficiently small $|u|$, that

$$(4.8) \quad x = 0: u_s = g_s(\alpha(t), h_{m+1}(t), \dots, h_n(t), u_1, \dots, u_m) \\ \stackrel{\text{def}}{=} \bar{g}_s(\bar{\alpha}(t), u_1, \dots, u_m) + \bar{h}_s(t) \quad (s = m + 1, \dots, n),$$

where $g_s \in C^1$ ($s = m + 1, \dots, n$),

$$(4.9) \quad \bar{\alpha}(t) = (\alpha(t), h_{m+1}(t), \dots, h_n(t)),$$

$$(4.10) \quad \bar{h}_s(t) = g_s(\bar{\alpha}(t), 0, \dots, 0) \quad (s = m + 1, \dots, n),$$

and then

$$(4.11) \quad \bar{g}_s(\bar{\alpha}(t), 0, \dots, 0) \equiv 0 \quad (s = m + 1, \dots, n).$$

It furthermore follows from (1.13) that

$$(4.12) \quad g_s(\alpha(t), 0, \dots, 0) \equiv 0 \quad (s = m + 1, \dots, n).$$

Let

$$(4.13) \quad W_b(T) = \max_{r=1, \dots, m} \int_0^T (-\lambda_r(u(t, 0))) |w_r(t, 0)| dt,$$

$$(4.14) \quad U_b(T) = \max_{r=1, \dots, m} \int_0^T (-\lambda_r(u(t, 0))) |v_r(t, 0)| dt,$$

$$(4.15) \quad U_\infty(T) = \sup_{0 \leq t \leq T} \sup_{x \geq 0} |u(t, x)|,$$

$$(4.16) \quad W_\infty(T) = \sup_{0 \leq t \leq T} \sup_{x \geq 0} |w(t, x)|,$$

$$(4.17) \quad W_1(T) = \sup_{0 \leq t \leq T} \int_0^\infty |w(t, x)| dx,$$

$$(4.18) \quad \tilde{U}_1(T) = \max_{i=1, \dots, n} \max_{j \neq i} \sup_{C_j} \int_{C_j} |u_i| dt,$$

$$(4.19) \quad \widetilde{W}_1(T) = \max_{i=1, \dots, n} \max_{j \neq i} \sup_{C_j} \int_{C_j} |w_i| dt,$$

where C_j stands for any given j th characteristic on the domain $D(T)$.

Hence, we conclude from (4.5) that

$$(4.20) \quad |u(t, x)| \leq \int_0^\infty |u_x(t, x)| dx \leq CW_1(T).$$

Here and hereafter, C will denote a generic positive constant independent of ε , M , and T ; the meaning of C may change from line to line.

Lemma 4.2. *There exists a positive constant C independent of ε , M , and T such that*

$$(4.21) \quad W_b(T), W_1(T), \widetilde{W}_1(T) \leq C\varepsilon,$$

$$(4.22) \quad U_b(T), \tilde{U}_1(T) \leq C \frac{\varepsilon}{M+1},$$

$$(4.23) \quad W_\infty(T) \leq CM.$$

Proof. We introduce

$$(4.24) \quad Q_U(T) = \sum_{j=1}^n \sum_{i \neq j} \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| dx dt,$$

$$(4.25) \quad Q_W(T) = \sum_{j=1}^n \sum_{i \neq j} \int_0^T \int_0^\infty |w_i(t, x)| \cdot |w_j(t, x)| dx dt,$$

$$(4.26) \quad Q_{UW}(T) = \sum_{j=1}^n \sum_{i \neq j} \int_0^T \int_0^\infty |u_i(t, x)| \cdot |w_j(t, x)| dx dt,$$

and

$$(4.27) \quad J_1 \stackrel{\text{def}}{=} \{1, \dots, m\}, \quad J_2 \stackrel{\text{def}}{=} \{m+1, \dots, n\}.$$

To begin with, we estimate $Q_U(T)$.

(i) For $i, j \in J_1$ and $i \neq j$, by (2.33) and Lemma 3.2 we have

$$(4.28) \quad \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| dx dt \\ \leq C \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \right)^2,$$

where $F = (F_1, \dots, F_n)^\top$ with

$$(4.29) \quad F_i = \sum_{j,k=1}^n F_{ijk}(u) u_j w_k + \sum_{j,k=1}^n P_{ijk}(u) u_j u_k.$$

By Hadamard's formula (see [19]), we get

$$(4.30) \quad |F_{ijj}(u)| \leq C \sum_{m \neq j} |u_m|, \quad |P_{ijj}(u)| \leq C \sum_{m \neq j} |u_m|, \quad \forall i, j \in \{1, \dots, n\};$$

as a consequence, it follows that

$$(4.31) \quad \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \leq C(Q_U(T) + Q_{UW}(T)).$$

Thus, for $i, j \in J_1$ and $i \neq j$, we conclude that

$$(4.32) \quad \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| dx dt \leq C(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T))^2.$$

(ii) For $i \in J_1, j \in J_2$, it also follows from (2.33) and Lemma 3.2 that

$$(4.33) \quad \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| dx dt \\ \leq C \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F_i(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \right) \\ \times \left(\|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F_j(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt + \int_0^T \lambda_j(u(t, 0)) |u_j(t, 0)| dt \right).$$

By (4.8)–(4.12), taking into account (4.2) and Lemma 3.1, we can obtain

$$\begin{aligned}
(4.34) \quad & \int_0^T \lambda_j(u(t, 0)) |u_j(t, 0)| \, dt \\
&= \int_0^T \lambda_j(u(t, 0)) |\bar{g}_j(\bar{\alpha}(t), u_1, \dots, u_m) + \bar{h}_j(t)| \, dt \\
&\leq \sum_{r=1}^m \int_0^T \lambda_j(u(t, 0)) \left| \frac{\partial \bar{g}_j}{\partial u_r} \right| \cdot |u_r(t, 0)| \, dt + \int_0^T \lambda_j(u(t, 0)) |\bar{h}_j(t)| \, dt \\
&\leq C(\|h\|_{L^1(\mathbb{R}^+)} + U_b(T)) \\
&\leq \left(\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + \int_0^T \|F(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right)
\end{aligned}$$

Due to (4.31), we conclude from (4.33)–(4.34) that

$$\begin{aligned}
(4.35) \quad & \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| \, dx \, dt \\
&\leq C(\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T))^2.
\end{aligned}$$

(iii) For $i \in J_2$, $j \in J_1$ or $i, j \in J_2$ and $i \neq j$, a similar procedure yields

$$\begin{aligned}
(4.36) \quad & \int_0^T \int_0^\infty |u_i(t, x)| \cdot |u_j(t, x)| \, dx \, dt \\
&\leq C(\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T))^2.
\end{aligned}$$

Combining the three cases above gives

$$(4.37) \quad Q_U(T) \leq C(\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T))^2.$$

Similarly to $Q_U(T)$, due to (2.35) and (4.6), it follows that

$$(4.38) \quad Q_W(T) \leq C \left(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right)^2$$

where $G = (G_1, \dots, G_n)^\top$ with

$$(4.39) \quad G_i = \sum_{j,k=1}^n \Gamma_{ijk}(u) w_j w_k + \sum_{j,k=1}^n Q_{ijk}(u) u_k w_j.$$

By (2.31) and Hadamard's formula, we have

$$(4.40) \quad |Q_{ijj}(u)| \leq C \sum_{m \neq j} |u_m|, \quad \forall i, j \in \{1, \dots, n\},$$

and then

$$(4.41) \quad \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \leq C(Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

Thus, it follows that

$$(4.42) \quad Q_W(T) \leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T))^2.$$

By virtue of (2.33) and (2.35), we similarly have

$$(4.43) \quad Q_{UW}(T) \leq C(\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T)) \times (\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

We now estimate $W_1(T)$.

(i) For $i \in J_1$, by (2.35) and Lemma 3.1 we have

$$(4.44) \quad \int_0^\infty |w_i(t, x)| dx \leq W_1(0) + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \leq W_1(0) + C(Q_W(t) + (1 + W_\infty(T))Q_{UW}(T)).$$

(ii) For $i \in J_2$, by (2.35) and 3.1 we also have

$$(4.45) \quad \int_0^\infty |w_i(t, x)| dx \leq W_1(0) + \int_0^T \|G(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt + \int_0^T \lambda_i(u(t, 0))|w_i(t, 0)| dt.$$

Using (4.6), we obtain

$$(4.46) \quad \int_0^T \lambda_i(u(t, 0))|w_i(t, 0)| dt = \int_0^T \lambda_i(u(t, 0)) \left| \sum_{r=1}^m f_{ir}(\alpha(t), u(t, 0))w_r(t, 0) + \sum_{l=1}^k \tilde{f}_{il}(\alpha(t), u(t, 0))\alpha'_l(t) + \sum_{\bar{s}=m+1}^n \tilde{f}_{i\bar{s}}(\alpha(t), u(t, 0))h'_{\bar{s}}(t) \right| dt$$

$$\begin{aligned}
&\leq \sum_{r=1}^m \int_0^T \frac{\lambda_i(u(t,0))}{-\lambda_r(u(t,0))} |f_{ir}(\alpha(t), u(t,0))| |(-\lambda_r(u(t,0))|w_r(t,0))| dt \\
&\quad + \int_0^T \lambda_i(u(t,0)) \left(\sum_{l=1}^k |\bar{f}_{il}(\alpha(t), u(t,0))| \cdot |\alpha'_l(t)| \right. \\
&\quad \quad \quad \left. + \sum_{\bar{s}=m+1}^n |\tilde{f}_{i\bar{s}}(\alpha(t), u(t,0))| \cdot |h'_{\bar{s}}(t)| \right) dt \\
&\leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)}) \\
&\quad + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T);
\end{aligned}$$

consequently, for $i \in J_2$ it follows that

$$(4.47) \quad \int_0^\infty |w_i(t, x)| dx \leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)}) + W_1(0) + Q_W(T) \\
+ (1 + W_\infty(T))Q_{UW}(T).$$

Thus, combining (4.44) and (4.47) yields

$$(4.48) \quad W_1(T) \leq (\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)}) + W_1(0) + Q_W(T) \\
+ (1 + W_\infty(T))Q_{UW}(T).$$

We next estimate $\widetilde{W}_1(T)$.

For this purpose, we need to estimate

$$\int_{C_j} |w_i| dt \quad (i \neq j).$$

For any given point $(t, x) \in D(T)$, denoted by A , we have:

(i) For $i \in J_1$, there are only two possibilities:

(a) The j th characteristic passing through the point A intersects the x -axis at a point $B(0, \alpha_j)$ and the i th characteristic passing through the point A intersects the x -axis at a point $P'(0, \alpha_i)$. One can rewrite (2.35) as

$$(4.49) \quad d(|w_i(t, x)|(dx - \lambda_i(u) dt)) = \text{sgn}(w_i)G_i dt \wedge dx.$$

Integrating(4.49) on the domain ABP , utilizing the Stokes formula and using (4.41), we have

$$(4.50) \quad \left| \int_{C_j} |w_i(t, x)|(\lambda_j(u) - \lambda_i(u)) dt \right| \\
\leq \left| \int_{\alpha_i}^{\alpha_j} |w_i(0, x)| dx \right| + \iint_{ABP} |G_i| dx dt \\
\leq W_1(0) + C(Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

By the definition of \widetilde{W}_1 , we have $j \neq i$ and then

$$(4.51) \quad |\lambda_j(u) - \lambda_i(u)| \geq \delta_0;$$

consequently, it follows that

$$(4.52) \quad \int_{C_j} |w_i(t, x)| dt \leq W_1(0) + C(Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

(b) The j th characteristic passing through the point A intersects the t -axis at a point $B(\beta_j, 0)$ and the i th characteristic passing through the point A intersects the x -axis at a point $P(0, \alpha_i)$. Denote the origin $(0, 0)$ by O . We integrate (4.49) on the domain $ABOP$, utilize the Stokes formula and apply Lemma 3.1 to obtain

$$(4.53) \quad \left| \int_{C_j} |w_i(t, x)| (\lambda_j(u) - \lambda_i(u)) dt \right| \\ \leq \int_0^{\alpha_i} |w_i(0, x)| dx + \iint_{ABOP} |G_i| dx dt + \int_0^{\beta_j} (-\lambda_i(u(t, 0)) |w_i(t, 0)|) dt \\ \leq C(W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

Hence, by (4.51)–(4.53), for $i \in J_1$ it follows that

$$(4.54) \quad \int_{C_j} |w_i(t, x)| dt \leq C(W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

(ii) For $i \in J_2$, there are four possibilities as follows:

(a') The j th characteristic passing through the point A intersects the x -axis at a point $B(0, \alpha_j)$ and the i th characteristic passing through the point A intersects the x -axis at a point $P(0, \alpha_i)$. Similarly to the case (a) in (i), we integrate (4.49) on the domain ABP and use the Stokes formula to obtain

$$(4.55) \quad \int_{C_j} |w_i(t, x)| dt \leq C(W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).$$

(b') The j th characteristic passing through the point A intersects the x -axis at a point $B(0, \alpha_j)$ and the i th characteristic passing through the point A intersects the t -axis at a point $P(\beta_i, 0)$. Integrating (4.49) on the domain $ABOP$, using the Stokes formula and taking into account boundary condition (4.6), (1.22), and Lemma 3.1,

we have

$$\begin{aligned}
(4.56) \quad & \left| \int_{C_j} |w_i(t, x)| (\lambda_j(u) - \lambda_i(u)) dt \right| \\
& \leq \int_0^{\alpha_j} |w_i(0, x)| dx + \iint_{ABOP} |G_i| dx dt + \int_0^{\beta_i} (\lambda_i(u(t, 0)) |w_i(t, 0)|) dt \\
& \leq C(W_1(0) + Q_W(T) + Q_{UW}(T)) \\
& \quad + \sum_{r=1}^m \int_0^T \lambda_i(u(t, 0)) |f_{ir}(\alpha(t), u(t, 0))| \cdot |w_r(t, 0)| dt \\
& \quad + \int_0^T \lambda_i(u(t, 0)) \left(\sum_{l=1}^k |\bar{f}_{il}(\alpha(t), u(t, 0))| \cdot |\alpha'_l(t)| \right. \\
& \qquad \qquad \qquad \left. + \sum_{\bar{s}=m+1}^n |\tilde{f}_{i\bar{s}}(\alpha(t), u(t, 0))| \cdot |h'_{\bar{s}}(t)| \right) dt \\
& \leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).
\end{aligned}$$

Thus, by (4.51), it follows that

$$\begin{aligned}
(4.57) \quad & \int_{C_j} |w_i(t, x)| dt \leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) \\
& \quad + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).
\end{aligned}$$

(c') The j th characteristic passing through the point A intersects the t -axis at a point $B(\beta_j, 0)$ and the i th characteristic passing through the point A intersects the x -axis at a point $P(0, \alpha_i)$. Similarly to case (b'), we integrate (4.49) on the domain $ABOP$, utilize the Stokes formula and due to the boundary condition (4.6), (1.22), and (4.51) we obtain

$$\begin{aligned}
(4.58) \quad & \int_{C_j} |w_i(t, x)| dt \\
& \leq C(\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)),
\end{aligned}$$

where we have used Lemma 3.1.

(d') The j th characteristic passing through the point A intersects the t -axis at a point $B(\beta_j, 0)$ and the i th characteristic passing through the point A intersects the t -axis at a point $P(\beta_i, 0)$. Similarly, integrating (4.49) on the domain ABP , utilizing

the Stokes formula and the boundary conditions, we have

$$\begin{aligned}
(4.59) \quad & \left| \int_{C_j} |w_i(t, x)| (\lambda_j(u) - \lambda_i(u)) dt \right| \\
& \leq \iint_{ABP} |G_i| dx dt + \left| \int_{\beta_i}^{\beta_j} (\lambda_i(u(t, 0)) |w_i(t, 0)|) dt \right| \\
& \leq C (\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) + (1 + W_\infty(T))Q_{UW}(T)).
\end{aligned}$$

Hence, for any $i, j \in \{1, \dots, n\}$ and $i \neq j$, it follows that

$$\begin{aligned}
(4.60) \quad & \int_{C_j} |w_i| dt \\
& \leq C (\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) \\
& \quad + (1 + W_\infty(T))Q_{UW}(T)),
\end{aligned}$$

so that

$$\begin{aligned}
(4.61) \quad & \widetilde{W}_1(T) \leq C (\|h'\|_{L^1(\mathbb{R}^+)} + \|\alpha'\|_{L^1(\mathbb{R}^+)} + W_1(0) + Q_W(T) \\
& \quad + (1 + W_\infty(T))Q_{UW}(T)).
\end{aligned}$$

Similarly,

$$(4.62) \quad \widetilde{U}_1(T) \leq C (\|h\|_{L^1(\mathbb{R}^+)} + \|\varphi_0\|_{L^1(\mathbb{R}^+)} + Q_U(T) + Q_{UW}(T)).$$

We finally estimate $W_\infty(T)$.

(i) For $i \in J_1$ and any given point $(t, x) \in D(T)$, the i th characteristic passing through the point (t, x) must intersect the x -axis at a point $(0, \alpha_i)$. By (2.34), we have

$$(4.63) \quad |w_i(t, x)| \leq W_\infty(0) + \left| \int_{C_i} \left(\sum_{j, k=1}^n \gamma_{ijk}(u) w_j w_k + \sum_{j, k=1}^n Q_{ijk}(u) u_k w_j \right) \right|.$$

By (2.21) and (2.22), it follows that

$$\begin{aligned}
(4.64) \quad & \left| \int_{C_i} \left(\sum_{j, k=1}^n \gamma_{ijk}(u) w_j w_k + \sum_{j, k=1}^n Q_{ijk}(u) u_k w_j \right) \right| \\
& \leq \int_{C_i} |\gamma_{iii}(u) w_i^2| + CW_\infty(T) \widetilde{W}_1(T) + \int_{C_i} |Q_{iii}(u) u_i w_i| + C\delta \widetilde{W}_1(T) \\
& \leq CW_\infty^2(T) \int_{C_i} |\gamma_{iii}(u)| + CW_\infty(T) \widetilde{W}_1(T) \\
& \quad + \delta W_\infty(T) \int_{C_i} |Q_{iii}(u)| + C\delta \widetilde{W}_1(T).
\end{aligned}$$

By (2.22), Hadamard's formula yields

$$(4.65) \quad |\gamma_{iii}(u)| = |\gamma_{iii}(u) - \gamma_{iii}(u_i e_i)| \leq C \sum_{j \neq i} |u_j|.$$

Then

$$(4.66) \quad \int_{C_i} |\gamma_{iii}(u)| \leq C \tilde{U}_1(T).$$

Due to (2.31), we also have

$$(4.67) \quad \int_{C_i} |Q_{iii}(u)| \leq C \tilde{U}_1(T).$$

Hence, for $i \in J_1$, it follows that

$$(4.68) \quad \|w_i(t, \cdot)\|_{C^0} \leq W_\infty(0) + CW_\infty^2(T) \tilde{U}_1(T) + CW_\infty(T) \widetilde{W}_1(T) \\ + C\delta W_\infty(T) \tilde{U}_1(T) + C\delta \widetilde{W}_1(T).$$

(ii) For $i \in J_2$ and any given point $(t, x) \in D(T)$, there are only two possibilities:

(a'') The i th characteristic passing through the point (t, x) intersects the x -axis at a point $(0, \alpha_i)$, and similarly to (i) above we have

$$(4.69) \quad \|w_i(t, \cdot)\|_{C^0} \leq W_\infty(0) + CW_\infty^2(T) \tilde{U}_1(T) + CW_\infty(T) \widetilde{W}_1(T) \\ + C\delta W_\infty(T) \tilde{U}_1(T) + C\delta \widetilde{W}_1(T).$$

(b'') The i th characteristic passing through the point (t, x) intersects the t -axis at a point $(\beta_i, 0)$. By (4.6), (4.68), and (1.20), it follows that

$$(4.70) \quad |w_i(t, x)| \leq |w_i(\beta_i, 0)| + \left| \int_{C_i} \left(\sum_{j,k=1}^n \gamma_{ijk}(u) w_j w_k + \sum_{j,k=1}^n Q_{ijk}(u) u_k w_j \right) \right| \\ \leq \left| \sum_{r=1}^m f_{ir}(\alpha(\beta_i), u(\beta_i, 0)) w_r(\beta_i, 0) + \sum_{l=1}^k \bar{f}_{il}(\alpha(\beta_i), u(\beta_i, 0)) \alpha_l'(\beta_i) \right. \\ \left. + \sum_{\bar{s}=m+1}^n \tilde{f}_{i\bar{s}}(\alpha(\beta_i), u(\beta_i, 0)) h_{\bar{s}}'(\beta_i) \right| + CW_\infty^2(T) \tilde{U}_1(T) \\ + CW_\infty(T) \widetilde{W}_1(T) + C\delta W_\infty(T) \tilde{U}_1(T) + C\delta \widetilde{W}_1(T) \\ \leq C(M + W_\infty(0) + W_\infty^2(T) \tilde{U}_1(T) + W_\infty(T) \widetilde{W}_1(T) \\ + \delta W_\infty(T) \tilde{U}_1(T) + \delta \widetilde{W}_1(T)).$$

By virtue of (4.68) and (4.70), it is easy to see that

$$(4.71) \quad W_\infty(T) \leq C(M + W_\infty(0) + W_\infty^2(T)\tilde{U}_1(T) + W_\infty(T)\widetilde{W}_1(T) + \delta W_\infty(T)\tilde{U}_1(T) + \delta\widetilde{W}_1(T)).$$

Combining (4.37), (4.42)–(4.43), (4.48), (4.61), (4.62), and (4.71), we can prove that

$$(4.72) \quad Q_U(T) \leq C \frac{\varepsilon^2}{(M+1)^2},$$

$$(4.73) \quad Q_W(T) \leq C\varepsilon^2,$$

$$(4.74) \quad Q_{UW}(T) \leq C \frac{\varepsilon^2}{M+1}$$

as well as the conclusion of Lemma 4.2. This completes the proof of Lemma 4.2. \square

By (4.20), it follows that

$$(4.75) \quad U_\infty(T) \leq CW_1(T) \leq C\varepsilon.$$

Taking ε sufficiently small, we get

$$(4.76) \quad U_\infty(T) \leq \frac{1}{2}\delta,$$

so the hypothesis (4.2) is reasonable.

Theorem 1.1 is a direct consequence of Lemma 4.2.

5. PROOF OF THEOREM 1.2

Proof. To estimate

$$\int_0^\infty |u^{(0)}(t, x)| dx,$$

it suffices to estimate

$$(5.1) \quad \int_0^\infty |\xi_i^{(1)}(t, x)| dx, \quad \forall i \in \{1, \dots, n\}$$

on a fixed time interval $[0, T]$.

By the boundary condition (1.11), we have

$$(5.2) \quad x = 0: v_s^{(1)} = f_s(\alpha^{(1)}(t), v_1^{(1)}, \dots, v_m^{(1)}) + h_s^{(1)}(t) \quad (s = m+1, \dots, n)$$

and

$$(5.3) \quad x = 0: v_s^{(2)} = f_s(\alpha^{(2)}(t), v_1^{(2)}, \dots, v_m^{(2)}) + h_s^{(2)}(t) \quad (s = m + 1, \dots, n),$$

where $v_i^{(l)} = l_i(u^{(l)})u^{(l)}$ ($i = 1, \dots, n; l = 1, 2$). By (2.38) and (2.40), it follows that

$$(5.4) \quad \begin{aligned} v_i^{(1)} - v_i^{(2)} &= l_i(u^{(1)})u^{(1)} - l_i(u^{(2)})u^{(2)} \\ &= \xi_i^{(1)} + \sum_{k=1}^n \xi_k^{(1)} r_k^\top(u^{(1)}) \frac{\partial l_i}{\partial u} u^{(2)} \quad (i = 1, \dots, n). \end{aligned}$$

Then, subtracting (5.2) from (5.3) yields, at $x = 0$, that

$$(5.5) \quad \begin{aligned} \xi_s^{(1)} + \sum_{k=1}^n \xi_k^{(1)} r_k^\top(u^{(1)}) \frac{\partial l_s}{\partial u} u^{(2)} \\ = \left(\frac{\partial f_s}{\partial \alpha} \right) (\alpha^{(0)}(t))^\top + \sum_{r=1}^m \frac{\partial f_s}{\partial v_r} \left[\xi_r^{(1)} + \sum_{k=1}^n \xi_k^{(1)} r_k^\top(u^{(1)}) \frac{\partial l_r}{\partial u} u^{(2)} \right] + h_s^{(0)}(t) \end{aligned} \quad (s = m + 1, \dots, n),$$

where $\alpha^{(0)}(t) = \alpha^{(1)}(t) - \alpha^{(2)}(t)$, $h_s^{(0)}(t) = h_s^{(1)}(t) - h_s^{(2)}(t)$. When ε is small enough, we conclude from Theorem 1.1 that

$$(5.6) \quad x = 0: \xi_s^{(1)} = \sum_{r=1}^m g_{sr}^{(1)} \xi_r^{(1)} + \sum_{i=1}^k \bar{g}_{si}^{(1)} \alpha_i^{(0)}(t) + \sum_{\bar{s}} \tilde{g}_{s\bar{s}}^{(1)} h_{\bar{s}}^{(0)}(t) \quad (s = m + 1, \dots, n),$$

where $g_{sr}^{(1)}$, $\bar{g}_{si}^{(1)}$, and $\tilde{g}_{s\bar{s}}^{(1)}$ are continuous functions with respect to $(\alpha^{(1)}(t), \alpha^{(2)}(t), u^{(1)}, u^{(2)})$.

Similarly,

$$(5.7) \quad x = 0: \xi_s^{(2)} = \sum_{r=1}^m g_{sr}^{(2)} \xi_r^{(2)} + \sum_{i=1}^k \bar{g}_{si}^{(2)} \alpha_i^{(0)}(t) + \sum_{\bar{s}} \tilde{g}_{s\bar{s}}^{(2)} h_{\bar{s}}^{(0)}(t) \quad (s = m + 1, \dots, n),$$

where $g_{sr}^{(2)}$, $\bar{g}_{si}^{(2)}$, and $\tilde{g}_{s\bar{s}}^{(2)}$ are continuous functions with respect to $(\alpha^{(1)}(t), \alpha^{(2)}(t), u^{(1)}, u^{(2)})$.

(i) For $i \in J_1$, due (2.41), it follows from Lemma 3.1 that

$$(5.8) \quad \begin{aligned} \int_0^\infty |\xi_i^{(1)}(t, x)| dx + \int_0^T (-\lambda_i(u^{(1)}(t, 0)) |\xi_i^{(1)}(t, 0)|) dt \\ \leq \int_0^\infty |\xi_i^{(1)}(0, x)| dx + \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}_+)} dt, \end{aligned}$$

where

$$\begin{aligned}
 (5.9) \quad F_i^{(1)} &= \sum_{j,k=1}^n \Theta_{ijk}(u^{(1)}) \xi_j^{(1)} w_k^{(1)} - (\lambda_i(u^{(1)}) - \lambda_i(u^{(2)})) l_i(u^{(1)}) r_i(u^{(2)}) w_i^{(2)} \\
 &+ \sum_{j \neq i} (\lambda_i(u^{(1)}) - \lambda_j(u^{(2)})) l_i(u^{(1)}) (r_j(u^{(1)}) - r_j(u^{(2)})) w_j^{(2)} \\
 &- \sum_{j=1}^n l_i(u^{(1)}) \nabla r_j(u^{(1)}) \xi_j^{(1)} + l_i(u^{(1)}) (B(u^{(1)}) - B(u^{(2)})).
 \end{aligned}$$

By Hadamard's formula, some estimates are obtained in [19], [20] as follows:

$$(5.10) \quad \sum_{j \neq i} |u_j^{(0)}| \leq C \sum_{j \neq i} |\xi_j^{(2)}| + C \sum_{j \neq i} |u_j^{(2)}| \cdot |\xi_i^{(2)}|,$$

$$(5.11) \quad |\lambda_i(u^{(1)}) - \lambda_i(u^{(2)})| \leq C \sum_{j \neq i} |\xi_j^{(2)}| + C \sum_{j \neq i} |u_j^{(2)}| \cdot |\xi_i^{(2)}|,$$

$$(5.12) \quad |r_j(u^{(1)}) - r_j(u^{(2)})| \leq C \sum_{k \neq j} |\xi_k^{(2)}| + C \sum_{k \neq j} |u_k^{(2)}| \cdot |\xi_j^{(2)}|,$$

$$(5.13) \quad |B(u^{(1)})| \leq C \sum_{k \neq j} |u_k^{(1)}|,$$

$$(5.14) \quad |\Theta_{ijj}(u^{(1)})| \leq C \sum_{k \neq j} |u_k^{(1)}|, \quad \forall i, j \in \{1, \dots, n\},$$

and

$$(5.15) \quad |B(u^{(1)}) - B(u^{(2)})| \leq C \sum_{j=1}^n \sum_{k \neq j} |u_k^{(2)}| \cdot |\xi_j^{(2)}| + C \sum_{j=1}^n \sum_{k \neq j} |u_k^{(1)}| \cdot |\xi_j^{(1)}|.$$

Combining the above facts, we obtain

$$\begin{aligned}
 (5.16) \quad &\int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \\
 &\leq C(Q_{\xi^{(1)}}(T) + (M+1)D_{\xi^{(1)}}(T) + Q_{\xi^{(2)}}(T) + (M+1)D_{\xi^{(2)}}(T)),
 \end{aligned}$$

where

$$(5.17) \quad Q_{\xi^{(1)}}(T) = \sum_{j=1}^n \sum_{i \neq j} \int_0^T \int_0^\infty |\xi_i^{(1)}(t, x)| \cdot |w_j^{(1)}(t, x)| dx dt,$$

$$(5.18) \quad Q_{\xi^{(2)}}(T) = \sum_{j=1}^n \sum_{i \neq j} \int_0^T \int_0^\infty |\xi_i^{(2)}(t, x)| \cdot |w_j^{(2)}(t, x)| dx dt,$$

$$(5.19) \quad D_{\xi^{(1)}}(T) = \sum_{j=1}^n \sum_{h \neq j} \int_0^T \int_0^\infty |u_h^{(1)}(t, x)| \cdot |\xi_j^{(1)}(t, x)| \, dx \, dt,$$

$$(5.20) \quad D_{\xi^{(2)}}(T) = \sum_{j=1}^n \sum_{h \neq j} \int_0^T \int_0^\infty |u_h^{(2)}(t, x)| \cdot |\xi_j^{(2)}(t, x)| \, dx \, dt.$$

We first estimate $Q_{\xi^{(1)}}(T)$.

(a) For $i, j \in J_1$ and $i \neq j$, by (2.35) and (2.41), it follows from Lemma 3.2 and Lemma 4.2 that

$$(5.21) \quad \begin{aligned} & \int_0^T \int_0^\infty |\xi_i^{(1)}(t, x)| \cdot |w_j^{(1)}(t, x)| \, dx \, dt \\ & \leq C \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right) \\ & \quad \times \left(W_1^{(1)}(0) + \sum_{i,j,k=1}^n \int_0^T \int_0^\infty |\Gamma_{ijk}(u^{(1)}) w_j^{(1)} w_k^{(1)} \right. \\ & \quad \quad \quad \left. + Q_{ijk}(u^{(1)}) u_k^{(1)} w_j^{(1)} \, dx \, dt \right) \\ & \leq C\varepsilon \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right), \end{aligned}$$

where $\varphi^{(0)} = \varphi^{(1)} - \varphi^{(2)}$.

(b) For $i \in J_1$ and $j \in J_2$, by virtue of (4.46) and Lemma 4.2, we conclude from Lemma 3.2 that

$$(5.22) \quad \begin{aligned} & \int_0^T \int_0^\infty |\xi_i^{(1)}(t, x)| \cdot |w_j^{(1)}(t, x)| \, dx \, dt \\ & \leq C \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right) \\ & \quad \times \left(W_1^{(1)}(0) + \int_0^T \lambda_j(u^{(1)}(t, 0)) |w_j^{(1)}(t, 0)| \, dt \right. \\ & \quad \quad \left. + \sum_{i,j,k=1}^n \int_0^T \int_0^\infty |\Gamma_{ijk}(u^{(1)}) w_j^{(1)} w_k^{(1)} + Q_{ijk}(u^{(1)}) u_k^{(1)} w_j^{(1)} \, dx \, dt \right) \\ & \leq C\varepsilon \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right). \end{aligned}$$

(c) For $i \in J_2$ and $j \in J_1$, due to the boundary condition (5.6), it follows from Lemma 3.2 that

$$\begin{aligned}
(5.23) \quad & \int_0^T \int_0^\infty |\xi_i^{(1)}(t, x)| \cdot |w_j^{(1)}(t, x)| \, dx \, dt \\
& \leq C \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right. \\
& \quad \left. + \int_0^T \lambda_i(u^{(1)}(t, 0)) |\xi_i^{(1)}(t, 0)| \, dt \right) \\
& \quad \times \left(W_1^{(1)}(0) + \sum_{i,j,k=1}^n \int_0^T \int_0^\infty |\Gamma_{ijk}(u^{(1)}) w_j^{(1)} w_k^{(1)} \right. \\
& \quad \left. + Q_{ijk}(u^{(1)}) u_k^{(1)} w_j^{(1)} \, dx \, dt \right) \\
& \leq C \varepsilon \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)} \right. \\
& \quad \left. + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right).
\end{aligned}$$

(d) For $i, j \in J_2$ and $i \neq j$, the similar procedure gives

$$\begin{aligned}
(5.24) \quad & \int_0^T \int_0^\infty |\xi_i^{(1)}(t, x)| \cdot |w_j^{(1)}(t, x)| \, dx \, dt \\
& \leq C \varepsilon \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)} \right. \\
& \quad \left. + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right).
\end{aligned}$$

Thus, we have

$$\begin{aligned}
(5.25) \quad & Q_{\xi^{(1)}}(T) \leq C \varepsilon \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)} \right. \\
& \quad \left. + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right).
\end{aligned}$$

Because of (2.23) and (2.41), we conclude from Lemma 3.2 that

$$\begin{aligned}
(5.26) \quad & D_{\xi^{(1)}}(T) \leq C \frac{\varepsilon}{M+1} \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)} \right. \\
& \quad \left. + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} \, dt \right).
\end{aligned}$$

Similar estimates hold true for $Q_{\xi^{(2)}}(T)$ and $D_{\xi^{(2)}}(T)$.

Consequently, it follows from (5.16) that

$$(5.27) \quad \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \\ \leq C\varepsilon(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)}).$$

Hence, for $i \in J_1$ we have

$$(5.28) \quad \int_0^\infty |\xi_i^{(1)}(t, x)| dx \leq C(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)}).$$

(ii) For $i \in J_2$, due (5.6), it follows from Lemma 3.1 that

$$(5.29) \quad \int_0^\infty |\xi_i^{(1)}(t, x)| dx \\ \leq \int_0^\infty |\xi_i^{(1)}(0, x)| dx + \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \\ + \int_0^T (\lambda_i(u^{(1)}(t, 0)) |\xi_i^{(1)}(t, 0)|) dt \\ \leq C \left(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \sum_{i=1}^n \int_0^T \|F_i^{(1)}(t, \cdot)\|_{L^1(\mathbb{R}^+)} dt \right. \\ \left. + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)} \right).$$

Thus, from (5.27) and (5.29), for $i \in J_2$, we conclude that

$$(5.30) \quad \int_0^\infty |\xi_i^{(1)}(t, x)| dx \leq C(\|\varphi^{(0)}\|_{L^1(\mathbb{R}^+)} + \|\alpha^{(0)}\|_{L^1(\mathbb{R}^+)} + \|h^{(0)}\|_{L^1(\mathbb{R}^+)}).$$

Therefore, by (5.28) and (5.30), we prove (5.1), which completes the proof of Theorem 1.2. \square

Acknowledgement. The author would like to thank Prof. Ta-Tsien Li and Prof. Yi Zhou for their valuable suggestions and sustained support, guidance and encouragement.

References

- [1] *A. Bressan*: Contractive metrics for nonlinear hyperbolic systems. *Indiana Univ. Math. J.* *37* (1988), 409–420.
- [2] *A. Bressan, T.-P. Liu, T. Yang*: L^1 stability estimates for $n \times n$ conservation laws. *Arch. Ration. Mech. Anal.* *149* (1999), 1–22.
- [3] *Y. L. Chen*: Global classical solution of the mixed initial-boundary value problem for a kind of the first order quasilinear hyperbolic system. *J. Fudan Univ. Nat. Sci.* *45* (2006), 625–631.
- [4] *J. M. Greenberg, T.-T. Li*: The effect of boundary damping for the quasilinear wave equation. *J. Differ. Equations* *52* (1984), 66–75.
- [5] *L. Hörmander*: The lifespan of classical solutions of nonlinear hyperbolic equations. *Lecture Notes Math.* 1256. Springer, Berlin, 1987, pp. 214–280.
- [6] *F. John*: Formation of singularities in one-dimensional nonlinear wave propagation. *Commun. Pure Appl. Math.* *27* (1974), 377–405.
- [7] *D. X. Kong*: Cauchy problem for first order quasilinear hyperbolic systems. *J. Fudan Univ. Nat. Sci.* *33* (1994), 705–708.
- [8] *S. M. Li*: Cauchy problem for general first order inhomogeneous quasilinear hyperbolic systems. *J. Partial Differ. Equations* *15* (2002), 46–68.
- [9] *T.-T. Li*: *Global Classical Solutions for Quasilinear Hyperbolic Systems*. Masson/John Wiley, Paris/Chichester, 1994.
- [10] *T.-T. Li, Y.-J. Peng*: The mixed initial-boundary value problem for reducible quasilinear hyperbolic systems with linearly-degenerate characteristics. *Nonlinear Anal., Theory Methods Appl.* *52* (2003), 573–583.
- [11] *T.-T. Li, Y.-J. Peng*: Global C^1 solution to the initial-boundary value problem for diagonal hyperbolic systems with linearly degenerate characteristics. *J. Partial Differ. Equations* *16* (2003), 8–17.
- [12] *T.-T. Li, L. B. Wang*: Global classical solutions to a kind of mixed initial-boundary value problem for quasilinear hyperbolic systems. *Discrete Contin. Dyn. Syst.* *12* (2005), 59–78.
- [13] *T.-T. Li, W.-C. Yu*: *Boundary Value Problems for Quasilinear Hyperbolic Systems*, Series V. Duke University, Mathematical Department, Durham, 1985.
- [14] *T.-T. Li, Y. Zhou, D.-X. Kong*: Weak linear degeneracy and global classical solutions for general quasilinear hyperbolic systems. *Commun. Partial Differ. Equations* *19* (1994), 1263–1317.
- [15] *T.-T. Li, Y. Zhou, D.-X. Kong*: Global classical solutions for general quasilinear hyperbolic systems with decay initial data. *Nonlinear Anal., Theory Methods Appl.* *28* (1997), 1299–1232.
- [16] *T.-P. Liu*: Development of singularities in the nonlinear waves for quasi-linear hyperbolic partial differential equations. *J. Differ. Equations* *33* (1979), 92–111.
- [17] *T.-P. Liu, T. Yang*: Well-posedness theory for hyperbolic conservation laws. *Commun. Pure Appl. Math.* *52* (1999), 1553–1586.
- [18] *T. H. Qin*: Global smooth solutions of dissipative boundary value problems for first order quasilinear hyperbolic systems. *Chin. Ann. Math., Ser. B* *6* (1985), 289–298.
- [19] *P.-X. Wu*: Global classical solutions to the Cauchy problem for general first order inhomogeneous quasilinear hyperbolic systems. *Chin. Ann. Math., Ser. A* *27* (2006), 93–108. (In Chinese.)
- [20] *Y. Zhou*: Global classical solutions to quasilinear hyperbolic systems with weak linear degeneracy. *Chin. Ann. Math., Ser. B* *25* (2004), 37–56.

- [21] *Y. Zhou, Y.-F. Yang*: Global classical solutions of mixed initial-boundary value problem for quasilinear hyperbolic systems. *Nonlinear Anal., Theory Methods Appl.* *73* (2010), 1543–1561.

Author's address: *Y.-F. Yang*, Department of Mathematics, College of Sciences, Hohai University, Nanjing 210098, Jiangsu, P. R. China, e-mail: yongfuyang@sina.com.