

Ali Rezaei Aliabad; Mehdi Badie
Fixed-place ideals in commutative rings

Commentationes Mathematicae Universitatis Carolinae, Vol. 54 (2013), No. 1, 53--68

Persistent URL: <http://dml.cz/dmlcz/143152>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2013

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Fixed-place ideals in commutative rings

ALI REZAEI ALIABAD, MEHDI BADIE

Abstract. Let I be a semi-prime ideal. Then $P_o \in \text{Min}(I)$ is called irredundant with respect to I if $I \neq \bigcap_{P_o \neq P \in \text{Min}(I)} P$. If I is the intersection of all irredundant ideals with respect to I , it is called a fixed-place ideal. If there are no irredundant ideals with respect to I , it is called an anti fixed-place ideal. We show that each semi-prime ideal has a unique representation as an intersection of a fixed-place ideal and an anti fixed-place ideal. We say the point $p \in \beta X$ is a fixed-place point if $O^p(X)$ is a fixed-place ideal. In this situation the fixed-place rank of p , denoted by $\text{FP-rank}_X(p)$, is defined as the cardinal of the set of all irredundant prime ideals with respect to $O^p(X)$. Let p be a fixed-place point, it is shown that $\text{FP-rank}_X(p) = \eta$ if and only if there is a family $\{Y_\alpha\}_{\alpha \in A}$ of cozero sets of X such that: 1- $|A| = \eta$, 2- $p \in \text{cl}_{\beta X} Y_\alpha$ for each $\alpha \in A$, 3- $p \notin \text{cl}_{\beta X} (Y_\alpha \cap Y_\beta)$ if $\alpha \neq \beta$ and 4- η is the greatest cardinal with the above properties. In this case p is an F -point with respect to Y_α for any $\alpha \in A$.

Keywords: ring of continuous functions, fixed-place, anti fixed-place, irredundant, semi-prime, annihilator, affiliated prime, fixed-place rank, Zariski topology

Classification: Primary 13Axx; Secondary 54C40

1. Introduction

In this article, any ring R is commutative with unity. A semi-prime ideal means an ideal which is an intersection of prime ideals. For each ideal I of R and each element a of R , we denote the ideals $\{x \in R : ax \in I\}$ by $(I : a)$. When $I = \{0\}$ we write instead $\text{Ann}(a)$ and call this the annihilator of a . If $\text{Ann}(a)$ is maximal in the set of all annihilators of nonzero elements of R , then $\text{Ann}(a)$ is a prime ideal of R , and it is called an affiliated prime ideal. A prime ideal P is said to be a minimal prime ideal over an ideal I , if there are no prime ideals strictly contained in P that contain I . By $\text{Min}(I)$ we mean the set of all minimal prime ideals over I ; we use $\text{Min}(R)$ instead of $\text{Min}(\{0\})$. A ring is called reduced, if the ideal $\{0\}$ is a semi-prime ideal. $\text{Zd}(R)$ stands for the set of all zero divisors of R . A prime ideal P is called a Bourbaki (resp. Zariski-Samuel) associated prime divisor of an ideal I if $(I : x) = P$ (resp., $(I : x)$ is P -primary) for some $x \in R$. We denote the set of Bourbaki (Zariski-Samuel) associated prime divisors of an ideal I by $\text{B}(I)$ (resp., $[\text{Z-S}](I)$). It is clear that if I is a semi-prime ideal, then $\text{B}(I) = [\text{Z-S}](I)$. A representation $I = \bigcap_{P \in \mathcal{P}} P$ of I as an intersection of prime ideals is called irredundant if no $P \in \mathcal{P}$ may be omitted.

For every $S \subseteq R$, by $h(S)$ and $h^c(S)$ we mean the sets $\{P \in \text{Min}(R) : S \subseteq P\}$ and $\text{Min}(R) \setminus h(S)$, respectively. Recall that the Zariski topology on $\text{Min}(R)$ is the one generated by the closed base $\{h(a) : a \in R\}$.

We assume throughout the paper that any topological space X is Tychonoff, βX is the Stone-Ćech compactification of X and $C(X)$ is the ring of all real valued continuous functions on X . For any $f \in C(X)$, we denote $f^{-1}\{0\}$ and $X \setminus f^{-1}\{0\}$ by $Z(f)$ and $\text{Coz}(f)$, respectively. Supposing $S \subseteq C(X)$, we define $Z(S) = \{Z(f) : f \in S\}$ and we use $Z(X)$ instead of $Z(C(X))$. For any $\mathcal{B} \subseteq Z(X)$, we define $Z^{-1}(\mathcal{B}) = \{f \in C(X) : Z(f) \in \mathcal{B}\}$. Suppose that $A \subseteq \beta X$, then by $M^A(X)$ and $O^A(X)$ we mean the sets $\{f \in C(X) : A \subseteq \text{cl}_{\beta X} Z(f)\}$ and $\{f \in C(X) : A \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} Z(f)\}$, respectively. If $A = \{p\}$ for some $p \in \beta X$, then for brevity, we use the notations $M^p(X)$ and $O^p(X)$. We denote $Z(O^p(X))$ by $\mathcal{O}^p(X)$. An element $p \in \beta X$ is called an F -point with respect to X if $\mathcal{O}^p(X)$ is prime ideal. The set of all isolated points of a topological space X is denoted by $I(X)$. Clearly, $\overline{I(X)} = X$ if and only if X has a smallest dense subspace; exactly, this subspace is equal to $I(X)$. The reader is referred to [6] for other terms and notations.

In Section 2, we introduce irredundant families and irredundant prime ideals with respect to a semi-prime ideal. We use these notions to define central concepts of the article: fixed-place ideals, anti fixed-place ideals, and fixed-place rank for the fixed-place ideals. We give equivalence conditions for the above concepts. We prove that the intersection of two fixed-place ideals is fixed-place. In Section 3, we show that each semi-prime ideal has a unique intersection representation of a fixed-place ideal and an anti fixed-place ideal. In Section 4, we obtain some results from viewpoint of the Zariski topology. We show that I is a fixed-place (an anti fixed-place) ideal if and only if the set of isolated points of $\text{Min}(I)$ is a dense subspace ($\text{Min}(I)$ has no isolated point). We introduce fixed-place families in this section, and prove that $\mathcal{P} \subseteq \text{Spec}(R)$ is fixed-place if and only if \mathcal{P} is discrete in the Zariski topology. In the final section, we study fixed-place ideals in $C(X)$. We show that the zero ideal of $C(X)$ is fixed-place (anti fixed-place) if and only if $I(X)$ is dense in X (X has no isolated point). We introduce fixed-place points and fixed-place rank of these points and we generalize Proposition 3.1 in [8] and Theorem 3.1 in [10].

2. Fixed-place ideal

By Theorem 2.1 and Proposition 4.11 of [12], each irredundant prime ideal of a semi-prime ideal I is of the form $(I : x)$, for some $x \in R$, and if a semi-prime ideal I is equal to an irredundant intersection of the family $\{P_\alpha\}_{\alpha \in A}$ of prime ideals, then $\{P_\alpha\}_{\alpha \in A}$ is the set of all irredundant prime ideals of I . We can summarize these facts as follows.

Theorem 2.1. *If I is a semi-prime ideal, then the following statements are equivalent.*

- (a) There is a family \mathcal{P} of prime ideals such that $I = \bigcap \mathcal{P}$ is an irredundant intersection.
- (b) $I = \bigcap \mathcal{B}(I)$.
- (c) $I = \bigcap [Z-S](I)$.
- (d) $I = \bigcap \{(I : x) : x \in R \text{ and } (I : x) \text{ is a prime ideal}\}$.

In this situation, we have

$$\mathcal{P} = \mathcal{B}(I) = [Z-S](I) = \{(I : x) : x \in R \text{ and } (I : x) \text{ is a prime ideal}\}.$$

Definition 2.2. Suppose that I is a semi-prime ideal of a ring R and $\emptyset \neq \mathcal{P} \subseteq \text{Min}(I)$. We say \mathcal{P} is *irredundant* with respect to I if $I \neq \bigcap_{P \in \text{Min}(I) \setminus \mathcal{P}} P$. If $\mathcal{P} = \{P\}$, then we say that P is irredundant with respect to I . If I is equal to the intersection of irredundant prime ideals of I , then we call I a *fixed-place ideal*, exactly, by Theorem 2.1, we have $I = \bigcap \mathcal{B}(I)$. In this situation the *fixed-place rank* of I is denoted by $\text{FP-rank}(I)$, and it is defined by the cardinal of $\mathcal{B}(I)$. If $\mathcal{B}(I) = \emptyset$, i.e., I has no irredundant prime ideal, then we call I an *anti fixed-place ideal*.

The following proposition is an immediate consequence of Theorem 2.1.

Proposition 2.3. If I is a semi-prime ideal of a ring R and $\mathcal{A} \subseteq \mathcal{B}(I) \neq \emptyset$, then $J = \bigcap_{P \in \mathcal{A}} P$ is a fixed-place ideal and $\mathcal{B}(J) = \mathcal{A}$.

Proposition 2.4. Let I be a semi-prime ideal of a ring R and $\mathcal{P} \subseteq \text{Min}(I)$. If $I = \bigcap_{P \in \mathcal{P}} P$, then $\mathcal{B}(I) \subseteq \mathcal{P}$.

PROOF: If $P_0 \notin \mathcal{P}$, then $\mathcal{P} \subseteq \text{Min}(I) \setminus \{P_0\}$. Thus

$$I \subseteq \bigcap_{P_0 \neq P \in \text{Min}(I)} P \subseteq \bigcap_{P_0 \neq P \in \mathcal{P}} P = I \Rightarrow \bigcap_{P_0 \neq P \in \text{Min}(I)} P = I \Rightarrow P_0 \notin \mathcal{B}(I).$$

Hence $\mathcal{B}(I) \subseteq \mathcal{P}$. □

Theorem 2.5. Let I be a semi-prime ideal of a ring R , $\mathcal{P} \subseteq \text{Min}(I)$ and $\mathcal{Q} = \{P/I : P \in \mathcal{P}\}$. The family \mathcal{P} is irredundant with respect to I if and only if \mathcal{Q} is irredundant with respect to the zero ideal of the ring R/I .

PROOF: We know that

$$I(a) \in \bigcap_{P/I \in \text{Min}(R/I) \setminus \mathcal{Q}} \frac{P}{I} = \bigcap_{P \in \text{Min}(I) \setminus \mathcal{P}} \frac{P}{I} \Leftrightarrow a \in \bigcap_{P \in \text{Min}(I) \setminus \mathcal{P}} P.$$

Thus

$$\bigcap_{P/I \in \text{Min}(R/I) \setminus \mathcal{Q}} \frac{P}{I} = \{0\} \Leftrightarrow \bigcap_{P \in \text{Min}(I) \setminus \mathcal{P}} P = I.$$

Therefore, \mathcal{P} is irredundant with respect to ideal I if and only if \mathcal{Q} is irredundant with respect to the zero ideal of the ring R/I . □

By the above theorem, we can see that for studying the fixed-place ideals it is sufficient to focus on the zero ideal of the reduced rings. Thus, in the remainder of this section we assume that R is reduced. By this assumption, it is clear that if P is a minimal prime ideal of R , then P is irredundant with respect to the zero ideal if and only if P is an affiliated prime ideal.

Proposition 2.6. *Suppose that R is a ring and $\mathcal{P} \subseteq \text{Min}(R)$.*

- (a) *If $\mathcal{P} \cap \mathcal{B}(\{0\}) \neq \emptyset$, then \mathcal{P} is irredundant with respect to $\{0\}$.*
 (b) *If $J = \bigcap_{P \in \mathcal{B}(\{0\})} P$,*

$$\emptyset \neq \mathcal{S} = \{P \in \text{Min}(R) : P \notin \mathcal{B}(\{0\}) \text{ and } P \not\supseteq J\} \text{ and}$$

$$\emptyset \neq \mathcal{T} = \{P \in \text{Min}(R) : P \notin \mathcal{B}(\{0\}) \text{ and } P \supseteq J\}$$

then \mathcal{S} is irredundant with respect to the zero ideal, where $\mathcal{S} \cap \mathcal{B}(\{0\}) = \emptyset$. Also, \mathcal{T} is not irredundant with respect to the zero ideal.

PROOF: The proof is straightforward. □

In this part, we study the irredundant family with respect to the zero ideal and give some equivalent conditions.

Proposition 2.7. *Let $\mathcal{P} \subseteq \text{Min}(R)$. If \mathcal{P} is an irredundant family with respect to the zero ideal R , then there exists $0 \neq a \in R$ such that $\bigcap \mathcal{P} \subseteq \text{Ann}(a)$.*

PROOF: Since \mathcal{P} is an irredundant family with respect to the zero ideal of R , we have that

$$\bigcap_{P \in \text{Min}(R) \setminus \mathcal{P}} P \neq \{0\}.$$

Say $0 \neq a \in \bigcap_{P \in \text{Min}(R) \setminus \mathcal{P}} P$. For any $b \in \bigcap_{a \in \mathcal{P}} P$,

$$ab \in \bigcap_{P \in (\text{Min}(R) \setminus \mathcal{P}) \cup \mathcal{P}} P = \bigcap_{P \in \text{Min}(R)} P = \{0\}.$$

Therefore, $\bigcap_{P \in \mathcal{P}} P \subseteq \text{Ann}(a)$. □

Lemma 2.8. *For each element $a \in R$, we have $\text{Min}(\text{Ann}(a)) = h^c(a)$.*

PROOF: We claim that $a \notin P$ for each $P \in \text{Min}(\text{Ann}(a))$. Suppose, on the contrary, that $a \in P$ for some $P \in \text{Min}(\text{Ann}(a))$. Then

$$\exists b \notin P, ab \in \text{Ann}(a) \Rightarrow a^2b = 0 \Rightarrow ab = 0 \Rightarrow b \in \text{Ann}(a) \Rightarrow b \in P,$$

which is impossible. Now, we prove that P is a minimal prime ideal for each $P \in \text{Min}(\text{Ann}(a))$. To see this,

$$\forall x \in P \quad \exists y \notin P \quad xy \in \text{Ann}(a) \Rightarrow xy a = 0.$$

But $ya \notin P$, hence P is a minimal prime ideal and therefore $P \in h^c(a)$. Consequently, $\text{Min}(\text{Ann}(a)) \subseteq h^c(a)$. Now, we show that $h^c(a) \subseteq \text{Min}(\text{Ann}(a))$.

Suppose that $P \in h^c(a)$. It is sufficient to show that $\text{Ann}(a) \subseteq P$. But it is evident, because $a \text{Ann}(a) = \{0\} \subseteq P$. \square

Proposition 2.9. *Let a be a nonzero element of R and $\mathcal{P} \subseteq \text{Min}(\text{Ann}(a))$. The family \mathcal{P} is irredundant with respect to the zero ideal of R if and only if \mathcal{P} is irredundant with respect to $\text{Ann}(a)$.*

PROOF: \Rightarrow) Suppose that the assertion of the proposition is false, then

$$\text{Ann}(a) = \bigcap_{P \in \text{Min}(\text{Ann}(a)) \setminus \mathcal{P}} P.$$

Thus

$$\begin{aligned} \{0\} &= \bigcap_{P \in \text{Min}(R)} P = \left(\bigcap_{P \in \text{Min}(\text{Ann}(a))} P \right) \cap \left(\bigcap_{\substack{P \in \text{Min}(R) \\ P \notin \text{Min}(\text{Ann}(a))}} P \right) \\ &= \left(\bigcap_{P \in \text{Min}(\text{Ann}(a)) \setminus \mathcal{P}} P \right) \cap \left(\bigcap_{\substack{P \in \text{Min}(R) \\ P \notin \text{Min}(\text{Ann}(a))}} P \right) = \bigcap_{P \in \text{Min}(R) \setminus \mathcal{P}} P. \end{aligned}$$

Therefore, \mathcal{P} is not irredundant with respect to the zero ideal of R and this is a contradiction.

\Leftarrow) Let $\mathcal{P} \subseteq \text{Min}(\text{Ann}(a)) = h^c(a)$ be irredundant with respect to $\text{Ann}(a)$. We show that \mathcal{P} is irredundant with respect to the zero ideal. On the contrary

$$\{0\} = \bigcap_{P \in \text{Min}(R) \setminus \mathcal{P}} P = \left(\bigcap_{P \in h^c(a) \setminus \mathcal{P}} P \right) \cap \left(\bigcap_{P \in h(a)} P \right).$$

Since $\bigcap_{P \in h(a)} P \not\subseteq Q$ for each $Q \in h^c(a)$, it follows that

$$\forall Q \in h^c(a) \quad \bigcap_{P \in h^c(a) \setminus \mathcal{P}} P \subseteq Q.$$

Hence

$$\bigcap_{P \in h^c(a) \setminus \mathcal{P}} P \subseteq \bigcap_{P \in h^c(a)} P = \text{Ann}(a) \quad \Rightarrow \quad \bigcap_{P \in h^c(a) \setminus \mathcal{P}} P = \text{Ann}(a),$$

which is a contradiction. \square

An immediate consequence of the above proposition is that $\text{Min}(\text{Ann}(a))$ is irredundant with respect to the zero ideal of R .

Proposition 2.10. *If the zero ideal in a ring R is a fixed-place ideal, then $\text{Zd}(R) = \bigcup_{P \in \mathcal{B}(\{0\})} P$.*

PROOF: Clearly, $\bigcup_{P \in \mathcal{B}(\{0\})} P \subseteq \text{Zd}(R)$. We only need to show that $\text{Zd}(R) \subseteq \bigcup_{P \in \mathcal{B}(\{0\})} P$. On the contrary, suppose that there exists a zero divisor a which is not in $\bigcup_{P \in \mathcal{B}(\{0\})} P$, then a nonzero element b exists such that $ab = 0$. Since for

each $P \in \mathcal{B}(\{0\})$, $a \notin P$ and $ab = 0 \in P$, it follows that $0 \neq b \in \bigcap_{P \in \mathcal{B}(\{0\})} P$. This contradicts our assumption. \square

Now, we are going to show that the intersection of two fixed-place ideals is also a fixed-place ideal. To see this, we need the following lemma.

Lemma 2.11. *If I and J are two fixed-place ideals of R such that $I \not\subseteq P$ for each $P \in \mathcal{B}(J)$ and $J \not\subseteq P$ for each $P \in \mathcal{B}(I)$, then $I \cap J$ is a fixed-place ideal.*

PROOF: Clearly, we have $I \cap J = \bigcap \{P \in \text{Spec}(R) : P \in \mathcal{B}(I) \cup \mathcal{B}(J)\}$. Now, by Theorem 2.1, it is enough to prove that $\mathcal{B}(I \cap J) = \mathcal{B}(I) \cup \mathcal{B}(J)$. To prove this, suppose that $P_0 \in \mathcal{B}(I) \cup \mathcal{B}(J)$. Without loss of generality, we may assume that $P_0 \in \mathcal{B}(I)$. On the contrary, let $I \cap J = \bigcap_{\substack{P \in \mathcal{B}(I) \cup \mathcal{B}(J) \\ P \neq P_0}} P$. Thus

$$\begin{aligned} \bigcap_{\substack{P \in \mathcal{B}(I) \cup \mathcal{B}(J) \\ P \neq P_0}} P \subseteq P_0 &\Rightarrow J \cap \left(\bigcap_{\substack{P \in \mathcal{B}(I) \\ P \neq P_0}} P \right) \subseteq P_0 \\ \Rightarrow \bigcap_{\substack{P \in \mathcal{B}(I) \\ P \neq P_0}} P \subseteq P_0 &\Rightarrow \bigcap_{\substack{P \in \mathcal{B}(I) \\ P \neq P_0}} P = I, \end{aligned}$$

which is a contradiction. \square

Theorem 2.12. *If I and J are two fixed-place ideals of R , then $I \cap J$ is also a fixed-place ideal.*

PROOF: Let $I' = \bigcap \{P \in \mathcal{B}(I) : P \not\subseteq J\}$ and $J' = \bigcap \{P \in \mathcal{B}(J) : P \not\subseteq I'\}$. Then, it is clear that $I \cap J = I' \cap J = I' \cap J'$, $\mathcal{B}(J') = \{P \in \mathcal{B}(J) : P \not\subseteq I'\}$ and $\mathcal{B}(I') = \{P \in \mathcal{B}(I) : P \not\subseteq J\}$. Therefore

$$(1) \quad \forall P \in \mathcal{B}(I') \quad J' \not\subseteq P, \quad \forall P \in \mathcal{B}(J') \quad I' \not\subseteq P.$$

On the other side, $I \subseteq I'$ and $J \subseteq J'$. Therefore, I' and J' are two fixed-place ideals such that they satisfy the condition of Lemma 2.11 and consequently $I' \cap J' = I \cap J$ is a fixed-place ideal. \square

In the following, we show that if I is not a fixed-place ideal, then there is no smallest fixed-place ideal containing I .

Proposition 2.13. *Let I be a semi-prime ideal of a ring R . If I is not fixed-place, then the set of all fixed-place ideals containing I has no minimal element.*

PROOF: On the contrary, suppose that J is a minimal element of that set. Consequently, $J \cap P$ is a fixed-place ideal for each $P \in \text{Min}(I)$. Thus $J = J \cap P$, hence $J \subseteq P$. This implies that $J = \bigcap_{P \in \text{Min}(I)} P = I$. It follows that I is a fixed-place ideal, which is a contradiction. \square

3. Unique decomposition

Let \mathcal{P} be the set of all subsets of $\text{Spec}(R)$. Throughout this section \sim and \leq denote the relations on \mathcal{P} defined by

$$\mathcal{S} \leq \mathcal{T} \Leftrightarrow \forall P_1 \in \mathcal{S}, \exists P_2 \in \mathcal{T} \quad P_1 \subseteq P_2$$

and

$$\mathcal{S} \sim \mathcal{T} \Leftrightarrow \bigcap \mathcal{S} = \bigcap \mathcal{T}.$$

It is easy to check that \mathcal{P} with relation \leq is a complete lattice, in which $\text{Spec}(R)$ is the greatest element and \emptyset is the smallest element. Furthermore, the relation \sim is an equivalence relation. If we put $\mathfrak{P} = \{[\mathcal{S}] : \mathcal{S} \subseteq \text{Spec}(R)\}$ and $\mathfrak{I} = \{I : I \text{ is a semi-prime ideal of } R\}$, then the function $K : \mathfrak{P} \rightarrow \mathfrak{I}$ (resp. $H : \mathfrak{I} \rightarrow \mathfrak{P}$) is defined by $K([\mathcal{S}]) = \bigcap [\mathcal{S}]$ (resp. $H(I) = [\text{Min}(I)]$). Throughout this section we use the above notation.

Lemma 3.1. *Let $\{\mathcal{S}_\alpha\}_{\alpha \in A}$ and $\{\mathcal{T}_\alpha\}_{\alpha \in A}$ be two families of subsets of $\text{Spec}(R)$. If $\mathcal{S}_\alpha \sim \mathcal{T}_\alpha$ for each $\alpha \in A$, then $[\bigcup_{\alpha \in A} \mathcal{S}_\alpha] = [\bigcup_{\alpha \in A} \mathcal{T}_\alpha]$.*

PROOF: The proof is standard. \square

Definition 3.2. We call two families \mathcal{S} and \mathcal{T} of \mathcal{P} *essentially disjoint* if

$$\forall P \in \mathcal{S} \quad \bigcap \mathcal{T} \not\subseteq P, \quad \forall P \in \mathcal{T} \quad \bigcap \mathcal{S} \not\subseteq P.$$

In this way, we call two classes \mathcal{S} and \mathcal{T} essentially disjoint, if there are two essentially disjoint families \mathcal{S} and \mathcal{T} in \mathcal{S} and \mathcal{T} , respectively. Finally, we say two semi-prime ideals I and J are essentially disjoint if $[H(I)]$ and $[H(J)]$ are essentially disjoint.

Theorem 3.3. *If I is a semi-prime ideal of R , then we can write I as a unique intersection of a fixed-place ideal and an anti fixed-place ideal which are essentially disjoint.*

PROOF: Let $J = \bigcap_{P \in \mathcal{B}(I)} P$, $\mathcal{A} = \{P \in \text{Min}(I) : P \notin \mathcal{B}(I), P \not\supseteq J\}$, $K = \bigcap_{P \in \mathcal{A}} P$ and $\mathcal{B} = \{P \in \text{Min}(I) : P \notin \mathcal{B}(I), P \supseteq J\}$. Then

$$I = \bigcap_{P \in \text{Min}(I)} P = \left(\bigcap_{P \in \mathcal{B}(I)} P \right) \cap \left(\bigcap_{P \in \mathcal{B}} P \right) \cap \left(\bigcap_{P \in \mathcal{A}} P \right) = J \cap K.$$

By Corollary 2.3, J is fixed-place. We claim that $\mathcal{A} \subseteq \text{Min}(K)$, because for each $P_\circ \in \mathcal{A}$, $K \subseteq P \subseteq P_\circ$, we have $I \subseteq K \subseteq P \subseteq P_\circ$. Since $P_\circ \in \text{Min}(I)$, $P = P_\circ$ and therefore $P_\circ \in \text{Min}(K)$. By Proposition 2.4, $\mathcal{B}(K) \subseteq \mathcal{A}$. For each $P_\circ \in \mathcal{A}$,

$$\begin{aligned} P_\circ \supseteq I &= \bigcap_{P_\circ \neq P \in \text{Min}(I)} P = \left(\bigcap_{P \in \mathcal{B}(I)} P \right) \cap \left(\bigcap_{P \in \mathcal{B}} P \right) \cap \left(\bigcap_{P_\circ \neq P \in \mathcal{A}} P \right) \\ &= J \cap \left(\bigcap_{P_\circ \neq P \in \mathcal{A}} P \right). \end{aligned}$$

Since $J \not\subseteq P_o$, we can write

$$\bigcap_{P_o \neq P \in \mathcal{A}} P \subseteq P_o \quad \Rightarrow \quad K = \bigcap_{P \in \mathcal{A}} P = \left(\bigcap_{P_o \neq P \in \mathcal{A}} P \right) \cap P_o = \bigcap_{P_o \neq P \in \mathcal{A}} P.$$

This implies that $P_o \notin \mathcal{B}(K)$ and therefore $\mathcal{B}(K) = \emptyset$, thus K is an anti fixed-place ideal. Now, suppose that $I = J_1 \cap K_1$ where J_1 is a fixed-place ideal, K_1 is an anti fixed-place ideal and J_1 and K_1 are essentially disjoint. So there are some essentially disjoint families of prime ideals \mathcal{P} and \mathcal{Q} such that $J_1 = \bigcap \mathcal{P}$ and $K_1 = \bigcap \mathcal{Q}$. Since J_1 is fixed-place, we can assume that $\mathcal{P} = \mathcal{B}(J_1)$. Now, we prove that $J_1 = J$. To prove this, it is enough to show that $\mathcal{B}(J) = \mathcal{B}(J_1)$. If $P_o \in \mathcal{B}(J) = \mathcal{B}(I)$, then there exists $a \in R$ such that $P_o = (I : a)$, thus $P_o = (J_1 \cap K_1 : a) = (J_1 : a) \cap (K_1 : a)$. Since K_1 is anti fixed-place, by Theorem 2.1, we have $P_o \neq (K_1 : a)$. Hence $P_o = (J_1 : a)$ and this implies that $P_o \in \mathcal{B}(J_1)$. Therefore, $\mathcal{B}(J) \subseteq \mathcal{B}(J_1)$. Conversely, if $P_o \in \mathcal{B}(J_1)$, then $K_1 \not\subseteq P_o$ and so there exists $a \in K_1 \setminus P_o$. Hence, $(I : a) = (J_1 \cap K_1 : a) = (J_1 : a) = \bigcap_{a \notin P \in \mathcal{B}(J_1)} P$. Since $(J_1 : a)$ is fixed-place and $P_o \in \mathcal{B}(J_1 : a)$, Theorem 2.1 shows that there exists $b \in R$ such that $((J_1 : a) : b) = P_o$. Therefore, $(I : ab) = ((I : a) : b) = ((J_1 : a) : b) = P_o$. This implies that $P_o \in \mathcal{B}(I) = \mathcal{B}(J)$, thus $\mathcal{B}(J_1) \subseteq \mathcal{B}(J)$, and therefore $J_1 = J$. To complete the proof, we must show that $K = K_1$. Clearly, since $J \cap K = I = J_1 \cap K_1$, for each $Q \in \mathcal{Q}$, we have $J = J_1 \not\subseteq Q$ and so $K \subseteq Q$. It follows that $K \subseteq K_1$. By the same manner we can see $K_1 \subseteq K$. Thus, $K_1 = K$. \square

Proposition 3.4. *The zero ideal in a reduced ring R is fixed-place if and only if $\text{Ann}(a)$ is fixed-place, for each $a \in \text{Zd}(R)$.*

PROOF: \Rightarrow) Since the zero ideal is a fixed-place ideal, $\{0\} = \bigcap_{P \in \mathcal{B}(\{0\})} P$. Thus $\text{Ann}(a) = (0 : a) = \left(\bigcap_{P \in \mathcal{B}(\{0\})} P \right) \cap (0 : a) = \bigcap_{a \notin P \in \mathcal{B}(\{0\})} P$ for each $a \in \text{Zd}(R)$. By Corollary 2.3, it implies that $\text{Ann}(a)$ is a fixed-place ideal.

\Leftarrow) By Theorem 3.3, $\{0\} = J \cap K$ where J is a fixed-place ideal, K is an anti fixed-place ideal and J and K are essentially disjoint. To obtain contradiction, suppose that $\{0\}$ is not fixed-place, then $J \neq \{0\}$, so there exists $0 \neq a \in J$. It is clear that $a \in \text{Zd}(R)$. Consequently

$$(1) \quad \text{Ann}(a) = (0 : a) = (J \cap K : a) = (K : a).$$

Since $\text{Ann}(a)$ is a fixed-place ideal, it has an irredundant prime ideal P . Hence, by Theorem 2.1, there exists an element b in R such that $(\text{Ann}(a) : b) = P$. We conclude from (1) that $(K : ab) = ((K : a) : b) = (\text{Ann}(a) : b) = P$. Theorem 2.1 shows that $P \in \mathcal{B}(K)$, which contradicts our assumption. \square

The following corollary is an immediate consequence of Theorem 2.5 and Proposition 3.4. This corollary will be needed to prove the next proposition.

Corollary 3.5. *Suppose that J is a semi-prime ideal of a ring R . The ideal J is fixed-place if and only if $(J : a)$ is a fixed-place ideal for each $a \in R$ where $J \subset (J : a)$.*

Proposition 3.6. *The zero ideal in a reduced ring R is an anti fixed-place ideal if and only if $\text{Ann}(a)$ is an anti fixed-place ideal for each $a \in \text{Zd}(R)$.*

PROOF: \Rightarrow) On the contrary, if for some $a \in \text{Zd}(R)$, $\text{Ann}(a)$ is not anti fixed-place, then there exist b in R and a prime ideal P such that $P = (\text{Ann}(a) : b)$, by Theorem 2.1. In this way $P = (\text{Ann}(a) : b) = (0 : ab) = \text{Ann}(ab)$. So, by Theorem 2.1, $P \in \mathcal{B}(\{0\})$, which is impossible.

\Leftarrow) By Theorem 3.3, there is a fixed-place ideal J and an anti fixed-place ideal K such that $\{0\} = J \cap K$. Suppose that the zero ideal is not an anti fixed-place ideal, so $K \neq \{0\}$. Hence there exists $0 \neq a \in K$. It is clear that $a \in \text{Zd}(R)$ and

$$(2) \quad \text{Ann}(a) = (0 : a) = (J \cap K : a) = (J : a).$$

Since J is a fixed-place ideal, $(J : a)$ is also a fixed-place ideal, by Corollary 3.5. So, we can conclude from (2) that $\text{Ann}(a)$ is a fixed-place ideal, which is a contradiction. \square

4. Fixed-place ideal and Zariski topology

Throughout this section, for convenience, by Z we mean the set $\text{Min}(R)$.

Theorem 4.1. *Let R be a reduced ring and $\mathcal{P} \subseteq Z$. The family \mathcal{P} is irredundant with respect to the zero ideal of R if and only if $\text{int}_Z \mathcal{P} \neq \emptyset$.*

PROOF: \Rightarrow) Since \mathcal{P} is an irredundant family with respect to the zero ideal of R , we have

$$\bigcap_{P \in Z \setminus \mathcal{P}} P \neq \{0\}.$$

Hence, there is a nonzero element $a \in \bigcap_{P \in Z \setminus \mathcal{P}} P$. Obviously, $\emptyset \neq h^c(a) \subseteq \mathcal{P}$. Therefore, $\text{int}_Z \mathcal{P} \neq \emptyset$.

\Leftarrow) Since $\text{int}_Z \mathcal{P} \neq \emptyset$, there is an element $a \in R$ such that

$$\emptyset \neq h^c(a) \cap Z \subseteq \mathcal{P} \Rightarrow Z \setminus \mathcal{P} \subseteq h(a) \neq Z \Rightarrow 0 \neq a \in \bigcap_{P \in h(a)} P \subseteq \bigcap_{P \in Z \setminus \mathcal{P}} P.$$

Consequently, \mathcal{P} is irredundant with respect to the zero ideal. \square

An immediate conclusion of the above theorem is the following corollary.

Corollary 4.2. *Let P be a minimal prime ideal of a reduced ring R . The ideal P is an isolated point of Z if and only if $P \in \mathcal{B}(\{0\})$.*

Corollary 4.3. *The zero ideal of a reduced ring R is anti fixed-place if and only if Z has no isolated point.*

PROOF: It is evident, by Corollary 4.2. \square

Theorem 4.4. *The zero ideal of a reduced ring R is a fixed-place ideal if and only if $\overline{I(Z)} = Z$.*

PROOF: \Rightarrow) Clearly, by Corollary 4.2, $\mathcal{B}(\{0\}) = I(Z)$. On the other side,

$$\text{cl}_Z \mathcal{B}(\{0\}) = h\left(\bigcap_{P \in \mathcal{B}(\{0\})} P\right) = h(\{0\}) = Z.$$

Thus, $\overline{I(Z)} = Z$.

\Leftarrow) Using Corollary 4.2, since $\overline{I(Z)} = Z$, we have $Z = \text{cl}_Z \mathcal{B}(\{0\})$. So, we can write

$$\begin{aligned} Z &= \text{cl}_Z \mathcal{B}(\{0\}) = h\left(\bigcap_{P \in \mathcal{B}(\{0\})} P\right) \\ \Rightarrow \quad h\left(\bigcap_{P \in \mathcal{B}(\{0\})} P\right) &= Z \quad \Rightarrow \quad \{0\} = \bigcap_{P \in \mathcal{B}(\{0\})} P. \end{aligned}$$

Therefore, the zero ideal is fixed-place. \square

Definition 4.5. Let \mathcal{P} be a family of prime ideals of a ring R . We say \mathcal{P} is a *fixed-place family*, whenever $\bigcap_{P_o \neq P \in \mathcal{P}} P \not\subseteq P_o$ for each $P_o \in \mathcal{P}$. Obviously, if \mathcal{P} is a fixed-place family, by Theorem 2.1, the ideal $I = \bigcap \mathcal{P}$ is fixed-place and $\mathcal{B}(I) = \mathcal{P}$.

Theorem 4.6. *Let \mathcal{P} be a family of prime ideals of a ring R . The family \mathcal{P} is fixed-place if and only if \mathcal{P} is discrete as a subspace of $\text{Spec}(R)$ with Zariski topology.*

PROOF: \Rightarrow) For convenience, assume that $V(S)$ denotes the set of all prime ideals containing $S \subseteq R$. Suppose that $P_o \in \mathcal{P}$ and $I = \bigcap \mathcal{P}$. It is enough to show that P_o is isolated in $\text{Min}(I)$ with Zariski topology. Set $J = \bigcap_{P_o \neq P \in \mathcal{P}} P$. Since \mathcal{P} is a fixed-place family, we have $J \not\subseteq P_o$. Clearly, $V(J) \cap \text{Min}(I) = \text{Min}(I) \setminus \{P_o\}$ and hence P_o is isolated in $\text{Min}(I)$.

\Leftarrow) Consider $P_o \in \mathcal{P}$. Since P_o is isolated in \mathcal{P} , there exists an ideal K of R such that $V(K) \cap \mathcal{P} = \mathcal{P} \setminus \{P_o\}$. It follows that $K \not\subseteq P_o$, $K \subseteq \bigcap_{P_o \neq P \in \mathcal{P}} P$ and hence $\bigcap_{P_o \neq P \in \mathcal{P}} P \not\subseteq P_o$. Therefore, \mathcal{P} is a fixed-place family. \square

5. Fixed-place ideal and $C(X)$

This section is divided in two parts. First we study the zero ideal of $C(X)$ and give two equivalent topological conditions. In the second part, we introduce the fixed-place point and the fixed-place rank of this point. Throughout this section $I(X)$ denotes the set of all isolated points of a topological space X .

Lemma 5.1. *Let A be a subset of a topological space X . The subset A is dense in X if and only if $M^A(X) = O^A(X) = \{0\}$.*

PROOF: \Rightarrow) Suppose that $f \in M^A(X)$. Since $A \subseteq Z(f)$, A is dense in X and $Z(f)$ is closed, so $Z(f) = X$. Consequently, $M^A(X) = O^A(X) = \{0\}$.

\Leftarrow) If A is not dense in X , then there is a point p in X such that $p \notin \text{cl}_X A$. Thus, there is a function f in $C(X)$ such that $p \notin Z(f)$ and $A \subseteq \text{int}_X Z(f)$, hence $0 \neq f \in O^A(X)$. This clearly forces $M^A(X) \supseteq O^A(X) \neq 0$. \square

Theorem 5.2. *Let $I(X)$ be the set of all isolated points of X , then $\mathcal{B}(\{0\}) = \{O^p(X) : p \in I(X)\}$.*

PROOF: Define $A = \{p \in \beta X : \exists P \in \mathcal{B}(\{0\}) \ O^p(X) \subseteq P\}$. We first prove that A is a subset of each dense subset of βX . Assume that D is a dense subset of βX . Fix $\mathcal{P} = \{P \in \text{Min}(C(X)) : \exists p \in D \ O^p \subseteq P\}$. If P_o is a minimal prime ideal and $P_o \notin \mathcal{P}$, then

$$\bigcap_{P_o \neq P \in \text{Min}(C(X))} P \subseteq \bigcap_{P \in \mathcal{P}} P = \bigcap_{p \in D} \bigcap_{P \in \text{Min}(O^p(X))} P = \bigcap_{p \in D} O^p(X) = O^D(X).$$

By Lemma 5.1, $O^D(X) = \{0\}$, thus $\bigcap_{P_o \neq P \in \text{Min}(C(X))} P = \{0\}$. It shows that $P_o \notin \mathcal{B}(\{0\})$, and therefore $\mathcal{B}(\{0\}) \subseteq \mathcal{P}$. Now suppose $p \in A$, then there exists $P \in \mathcal{B}(\{0\})$ such that $O^p(X) \subseteq P$. Therefore $P \in \mathcal{P}$ and consequently there exists $p' \in D$ such that $O^{p'}(X) \subseteq P$. By [6, 2.11] and [6, 4I.4], every prime z -ideal contains a unique $O^x(X)$ for some $x \in \beta X$, hence $p = p' \in D$ and therefore $A \subseteq D$, which is desired. It is easily seen that the intersection of all dense subset of βX is equal to $I(X)$, thus $A \subseteq I(X)$, and therefore

$$(3) \quad \mathcal{B}(\{0\}) \subseteq \{O^p(X) : p \in I(X)\}.$$

Now, consider $p_o \in I(X)$. Then it is clear that $O^{p_o}(X)$ is prime and

$$\bigcap_{O^{p_o} \neq P \in \text{Min}(C(X))} P \supseteq \bigcap_{p_o \neq p \in X} O^p(X) = O^{X \setminus \{p_o\}}(X) \neq 0.$$

Hence, $O^{p_o}(X) \in \mathcal{B}(\{0\})$. Thus

$$(4) \quad \mathcal{B}(\{0\}) \supseteq \{O^p : p \in I(X)\}.$$

From (3) and (4), we obtain $\mathcal{B}(\{0\}) = \{O^p(X) : p \in I(X)\}$. \square

Corollary 5.3. *A space X has an isolated point if and only if the space $\text{Min}(C(X))$ has an isolated point.*

PROOF: Applying Corollary 4.2 and Theorem 5.2, it follows clearly. \square

Corollary 5.4. *The zero ideal of $C(X)$ is anti fixed-place if and only if X has no isolated point.*

PROOF: By Corollary 4.3 and Theorem 5.2, it is obvious. \square

Theorem 5.5. *The zero ideal in the ring $C(X)$ is fixed-place if and only if $\overline{I(X)} = X$. Then $\text{FP-rank}(\{0\}) = |I(X)|$.*

PROOF: \Rightarrow) Since the zero ideal is a fixed-place ideal, Theorem 5.2 shows that

$$\{0\} = \bigcap_{P \in \mathcal{B}(\{0\})} P = \bigcap_{p \in I(X)} O^p(X) = O^{I(X)}(X).$$

We conclude from Lemma 5.1 that $I(X)$ is dense in X .

\Leftarrow) Since $\overline{I(X)} = X$, by Lemma 5.1, $O^{I(X)}(X) = \{0\}$. Now, using Theorem 5.2, we have

$$\bigcap_{P \in \mathcal{B}(\{0\})} P = \bigcap_{p \in I(X)} O_p = O^{I(X)} = \{0\}.$$

Therefore, the zero ideal of $C(X)$ is fixed-place. \square

Corollary 5.6. $I(X)$ is a dense subspace if and only if the set of isolated points of $\text{Min}(C(X))$ is also a dense subspace.

PROOF: It is evident, by Theorems 4.4 and 5.5. \square

Example 5.7. Let X be an almost discrete space with the only non-isolated point p . The zero ideal of $C(X)$ is a fixed-place ideal and $\text{FP-rank}(\{0\}) = |X|$, by Theorem 5.5. Thus, for every cardinal number α there is a fixed-place ideal of fixed-place rank α .

The notion of rank of a point of a topological space was first introduced and studied in [8, 1.7], further in [10] and [11], and generalized in [1]. One can find in [9, 4.1] a basis of this concept as *FMP*-point. The following definition is based on similar definition in [3, 4.3] with a few differences. Actually, the root of this generalized definition may be found in [1].

Definition 5.8. Let X be a topological space and $p \in \beta X$. We call p a *fixed-place point* with respect to X if $O^p(X)$ is a fixed-place ideal and the fixed-place rank of p with respect to X , denoted by $\text{FP-rank}_X(p)$, is defined to be $\text{FP-rank}(O^p(X))$.

In [3, Theorem 4.4], it is shown that there is a point of fixed-place rank η for any given cardinal η . It is easy to see that p is a fixed-place point with respect to X if and only if it is a fixed-place point with respect to βX , for each $p \in \beta X$. Furthermore, if p is a fixed-place point with respect to X , then $\text{FP-rank}_X(p) = \text{FP-rank}_{\beta X}(p)$.

The Proposition 3.1 of [8] states: "Let X be a compact space. A point $p \in X$ has rank $k (< \infty)$ if and only if there is a family of k pairwise disjoint cozero sets with p being in each of their closures, but no larger family of pairwise disjoint cozero sets with this feature." Also, in [10, Theorem 3.1] the same proposition is given for completely regular spaces. Of course, this proposition, with a few differences, was also shown in [2]. In the remainder of this section, we want to generalize this proposition for any $p \in \beta X$ and any cardinal number. To do this, we need some facts.

Suppose that Y is a subspace of X and $\mathbf{F}(X)$ and $\mathbf{F}(Y)$ are the set of all z -filters of X and Y , respectively. Let $\gamma : \mathbf{F}(Y) \rightarrow \mathbf{F}(X)$ be given by

$$\gamma(\mathcal{F}) = \{Z \in Z(X) : Z \cap Y \in \mathcal{F}\}.$$

Then $\psi = Z^{-1}\gamma Z$ is a map from all z -ideals of $C(Y)$ to the set of all z -ideals of $C(X)$. We consider $\Phi : \beta Y \rightarrow \beta X$ as extension of the identity map $Y \rightarrow X$. In the remainder of this section we use the above notation. See [4] for more information about the above maps.

Proposition 5.9. *Let Y be a subspace of X . The subspace*

$$Y' = \{p \in \beta X : \Phi^{-1}(p) \text{ is a singleton}\}$$

is the largest subspace of βX that can be considered homeomorphically as a subspace of βY by Φ .

PROOF: The proof is standard. □

The above proposition leads us to the following definition.

Definition 5.10. Suppose that Y is a subspace of X and $p \in \beta X$. We say p is an F -point with respect to Y if $\Phi^{-1}(p)$ is a singleton and this point is an F -point with respect to Y .

Proposition 5.11. *Let X be a compact space, $p \in X$ and $Y = \text{Coz}(f)$, for some $f \in C(X)$. If $p \in \text{cl}_X Y$, then ψ is a one-to-one correspondence map between $\bigcup_{\Phi(q)=p} \text{Min}(O^q(Y))$ and $\{P \in \text{Min}(O^p(X)) : f \notin P\}$.*

PROOF: It follows from [4, Theorem 4.1 and Theorem 4.2]. □

Lemma 5.12. *Let X be a topological space, $f \in C(X)$ and $p \in \beta X$. A prime ideal P is irredundant with respect to $O^p(X)$ if and only if there is a cozero set Y in X such that p is an F -point with respect to Y and $P = \psi(O^q(Y))$, in which $\Phi(q) = p$.*

PROOF: Suppose that $Y = \text{Coz}(f)$ for some $f \in C(X)$. By Proposition 5.11, there is the one-to-one correspondence ψ between $\bigcup_{\Phi(q)=p} \text{Min}(O^q(Y))$ and $\{P \in \text{Min}(O^p(X)) : f \notin P\}$. Thus

$$\begin{aligned} (O^p : f) &= \left(\bigcap_{P \in \text{Min}(O^p(X))} P : f \right) = \bigcap_{P \in \text{Min}(O^p(X))} (P : f) \\ (5) \quad &= \bigcap_{\substack{P \in \text{Min}(O^p(X)) \\ f \notin P}} P = \bigcap_{\Phi(q)=p} \left(\bigcap_{Q \in \text{Min}(O^q(Y))} \psi(Q) \right). \end{aligned}$$

Now, by Theorem 2.1, it follows that P is irredundant with respect to $O^p(X)$ if and only if there is $f \in C(X)$ such that $(O^p : f) = P$. By (5),

$$P = \bigcap_{\Phi(q)=p} \left(\bigcap_{P \in \text{Min}(O^q(Y))} \psi(Q) \right).$$

Since the $\psi(Q)$'s in the above equality are minimal prime, this is equivalent to saying that $\Phi^{-1}(p)$ is a singleton and p is an F -point with respect to Y and consequently, this is equivalent to saying that $P = \psi(O^q(Y))$ where $\{q\} = \Phi^{-1}\{p\}$. \square

Lemma 5.13. *Let $p \in \beta X$ and $Y_1 = \text{Coz}(f_1)$, $Y_2 = \text{Coz}(f_2)$ for some $f_1, f_2 \in C(X)$. Then $f_1 f_2 \in O^p(X)$ if and only if $p \notin \text{cl}_{\beta X}(Y_1 \cap Y_2)$.*

PROOF: First we note that for each $f \in C^*(X)$

$$\text{cl}_{\beta X} \text{Coz}(f^\beta) = \text{cl}_{\beta X}(\text{Coz}(f^\beta) \cap X) = \text{cl}_{\beta X} \text{Coz}(f).$$

Now, without loss of generality, we assume that $f_1, f_2 \in C^*(X)$. Thus

$$\begin{aligned} p \notin \text{cl}_{\beta X}(Y_1 \cap Y_2) &= \text{cl}_{\beta X}(\text{Coz}(f_1) \cap \text{Coz}(f_2)) \\ &= \text{cl}_{\beta X} \text{Coz}(f_1 f_2) = \text{cl}_{\beta X} \text{Coz}(f_1^\beta f_2^\beta) \\ \Leftrightarrow p \in \text{int}_{\beta X} Z(f_1^\beta f_2^\beta) &= \text{int}_{\beta X} \text{cl}_{\beta X} Z(f_1 f_2) \Leftrightarrow f_1 f_2 \in O^p(X). \end{aligned}$$

\square

Proposition 5.14. *Let X be a topological space, $p \in \beta X$ and $\{Y_\alpha\}_{\alpha \in A}$ be a family of cozero sets of X . If*

- (a) p is F -point with respect to Y_α for each $\alpha \in A$,
- (b) $p \notin \text{cl}_{\beta X}(Y_\alpha \cap Y_\beta)$, if $\alpha \neq \beta$,

then $|\mathcal{B}(O^p(X))| \geq |A|$.

PROOF: Let $Y_\alpha = \text{Coz}(f_\alpha)$ for each $\alpha \in A$. By hypothesis, for each $\alpha \in A$ there is a unique $q_\alpha \in \beta Y_\alpha$ such that $\Phi_\alpha(q_\alpha) = p$. If we put $P_\alpha = \psi_\alpha(O^{q_\alpha}(Y_\alpha))$ for every $\alpha \in A$, then by Proposition 5.11, $\{P_\alpha\}_{\alpha \in A}$ is a family of irredundant ideals with respect to $O^p(X)$. It is sufficient to show that if $\alpha \neq \beta$, then $P_\alpha \neq P_\beta$. By Lemma 5.13, since $p \notin \text{cl}_{\beta X}(Y_\alpha \cap Y_\beta)$, we have $f_\alpha f_\beta \in O^p(X) \subseteq P_\alpha$. It follows from Proposition 5.11 that $f_\alpha \notin P_\alpha$. Therefore, $f_\beta \in P_\alpha$ and hence $P_\alpha \neq P_\beta$. \square

Lemma 5.15. *Let X be a topological space and $p \in \beta X$ be a fixed-place point. If $\text{FP-rank}_X(p) = \eta$ and ζ is a cardinal such that $\zeta \leq \eta$, then there exist a family $\{Y_\alpha\}_{\alpha \in A}$ of distinct cozero sets such that*

- (a) $|A| = \zeta$;
- (b) p is an F -point with respect to Y_α for every $\alpha \in A$;
- (c) $p \notin \text{cl}_{\beta X}(Y_\alpha \cap Y_\beta)$, if $\alpha \neq \beta$.

PROOF: Since $\text{FP-rank}_X(p) = \eta$, Theorem 2.1 shows that there exists a family $\{P_\alpha\}_{\alpha \in B}$ of minimal prime ideals such that $|B| = \eta$, $O^p(X) = \bigcap_{\alpha \in B} P_\alpha$ and this intersection is irredundant. Suppose that $\{P_\alpha\}_{\alpha \in A}$ is a subfamily of $\{P_\alpha\}_{\alpha \in B}$ such that $|A| = \zeta$. By Lemma 5.12, for every $\alpha \in A$, there exists a cozero set $Y_\alpha = \text{Coz}(f_\alpha)$ such that p is an F -point with respect to Y_α , for each $\alpha \in A$. If $\alpha \neq \beta$, it is easy to check that either f_α or f_β must be in P_θ , for each $\theta \in B$. Thus $f_\alpha f_\beta \in P_\theta$, for each $\theta \in B$, and therefore $f_\alpha f_\beta \in O^p(X)$. It shows that $p \notin \text{int}_{\beta X}(Y_\alpha \cap Y_\beta)$, by Lemma 5.13. \square

The following is another version of Lemma 5.14.

Lemma 5.16. *Let $p \in \beta X$ be a fixed-place point and $\{Y_\alpha\}_{\alpha \in A}$ be a family of cozero sets of X . If*

- (a) $p \in \text{cl}_{\beta X} Y_\alpha$ for each $\alpha \in A$,
- (b) $p \notin \text{cl}_{\beta X} (Y_\alpha \cap Y_\beta)$ for $\alpha \neq \beta$,

then $\text{FP-rank}_X(p) \geq |A|$.

PROOF: For each $\alpha \in A$, set $Y_\alpha = \text{Coz}(f_\alpha)$ with $f_\alpha \in C^*(X)$. Define

$$\mathcal{Q}_\alpha = \{Q \in \text{Min}(O^q(Y_\alpha)) : \Phi_\alpha(q) = p\}.$$

We claim that $\psi_\alpha(\mathcal{Q}_\alpha) \cap \psi_\beta(\mathcal{Q}_\beta) = \emptyset$, for each $\alpha \neq \beta$. Let $\psi_\alpha(Q) \in \psi_\alpha(\mathcal{Q}_\alpha)$, then $f_\alpha \notin \psi_\alpha(Q)$. Since $p \notin \text{cl}_{\beta X} (Y_\alpha \cap Y_\beta)$, by Lemma 5.13, obviously $f_\alpha f_\beta \in O^p(X)$. Thus $f_\beta \in \psi_\alpha(Q)$, this implies that $\psi_\alpha(Q) \notin \psi_\beta(\mathcal{Q}_\beta)$. To complete the proof, it is sufficient to show that $\mathcal{B}(O^p(X)) \cap \psi_\alpha(\mathcal{Q}_\alpha) \neq \emptyset$ for each $\alpha \in A$. We know that $f_\alpha \notin \psi_\alpha(Q)$, for each $Q \in \mathcal{Q}_\alpha$ and $f_\alpha \in \bigcap_{P \in \text{Min}(O^p(X)) \setminus \psi_\alpha(\mathcal{Q}_\alpha)} P$. Thus

$$\begin{aligned} f_\alpha &\in \bigcap_{P \in \text{Min}(O^p(X)) \setminus \psi_\alpha(\mathcal{Q}_\alpha)} P \setminus O^p(X) \\ &\Rightarrow O^p(X) \neq \bigcap_{P \in \text{Min}(O^p(X)) \setminus \psi_\alpha(\mathcal{Q}_\alpha)} P. \end{aligned}$$

This implies that $\psi(\mathcal{Q}_\alpha)$ is irredundant with respect to $O^p(X)$. Since $O^p(X)$ is fixed-place, $\mathcal{B}(O^p(X)) \cap \psi_\alpha(\mathcal{Q}_\alpha) \neq \emptyset$. \square

Now, we are ready to state the main theorem of this section.

Theorem 5.17. *Let $p \in \beta X$ be a fixed-place point. Then $\text{FP-rank}_X(p) = \eta$ if and only if there exists a family $\{Y_\alpha\}_{\alpha \in A}$ of cozero sets of X such that*

- (a) $|A| = \eta$;
- (b) $p \in \text{cl}_{\beta X} Y_\alpha$, for each $\alpha \in A$;
- (c) $p \notin \text{cl}_{\beta X} (Y_\alpha \cap Y_\beta)$, if $\alpha \neq \beta$;
- (d) η is the greatest cardinal with above properties.

In this case, p is an F -point with respect to Y_α for each $\alpha \in A$.

PROOF: \Rightarrow) By Lemma 5.15, there is a family $\{Y_\alpha\}_{\alpha \in A}$ of cozero sets of X with properties (a)–(c), and by Lemma 5.16, it follows that η is the greatest cardinal with properties (a)–(c).

\Leftarrow) Lemma 5.16 shows that $\text{FP-rank}_X(p) \geq \eta$. Since η is the greatest cardinal with properties (a)–(c), by Lemma 5.15, $\text{FP-rank}_X(p) = \eta$.

According to Lemma 5.12, p is an F -point with respect to Y_α , for each $\alpha \in A$. \square

It is clear to see that if $f \in O^p(X)$, then there exists $g \notin M^p(X)$ such that $fg = 0$. Using this fact, we may obtain the following result.

Corollary 5.18. *A point $p \in \beta X$ has finite fixed-place rank n if and only if there is a collection of n pairwise disjoint cozero sets $\{Y_i\}_{i=1}^n$ such that p is in the closure of each Y_i , and there is no larger such collection. In this case, p is an F -point with respect to Y_i , for each $i = 1, \dots, n$.*

REFERENCES

- [1] Aliabad A.R., *z° -ideals in $C(X)$* , Ph.D. Thesis, Chamran University of Ahvaz, Iran.
- [2] Aliabad A.R., *Connections between $C(X)$ and $C(Y)$, where Y is a subspace of X* , Abstracts of International Conference on Applicable General Topology, August 12–18, 2001, Hacettepe University, Ankara, Turkey.
- [3] Aliabad A.R., *Pasting topological spaces at one point*, Czechoslovak Math. J. **56** (131) (2006), 1193–1206.
- [4] Aliabad A.R., Badie M., *Connection between $C(X)$ and $C(Y)$, where Y is subspace of X* , Bull. Iranian Math. Soc. **37** (2011), no. 4, 109–126.
- [5] Engelking R., *General Topology*, PWN-Polish Scientific Publishing, Warsaw, 1977.
- [6] Gillman L., Jerison M., *Rings of Continuous Functions*, Van Nostrand Reinhold, New York, 1960.
- [7] Goodearl K.R., Warfield R.B., Jr. *Introduction to Noncommutative Noetherian Rings*, Cambridge University Press, Cambridge, 1989.
- [8] Henriksen M., Larson S., Martinez J., Woods R.G., *Lattice-ordered algebras that are sub-direct product of valuation domains*, Trans. Amer. Math. Soc. **345** (1994), 195–221.
- [9] Henriksen M., Wilson R.G., *Almost discrete SV -space*, Topology Appl. **46** (1992), 89–97.
- [10] Larson S., *f -Rings in which every maximal ideal contains finitely many minimal prime ideals*, Comm. Algebra **25** (1997), no. 12, 3859–3888.
- [11] Larson S., *Constructing rings of continuous functions in which there are many maximal ideals with nontrivial rank*, Comm. Algebra **31** (2003), 2183–2206.
- [12] Underwood D.H., *On some Uniqueness questions in primary representations of ideals*, Kyoto Math. J. **35** (1969), 69–94.
- [13] Willard S., *General Topology*, Addison Wesley, Reading, Mass., 1970.

MATHEMATICS DEPARTMENT, CHAMRAN UNIVERSITY, AHVAZ, IRAN

E-mail: aliabady_r@scu.ac.ir

BASIC SCIENCE DEPARTMENT, JUNDI SHAPUR UNIVERSITY OF TECHNOLOGY, DEZFUL, IRAN

E-mail: badie@jsu.ac.ir

(Received July 13, 2012, revised October 20, 2012)