

Jiří Benedikt

Estimates of the principal eigenvalue of the p -Laplacian and the p -biharmonic operator

Mathematica Bohemica, Vol. 140 (2015), No. 2, 215–222

Persistent URL: <http://dml.cz/dmlcz/144327>

Terms of use:

© Institute of Mathematics AS CR, 2015

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ESTIMATES OF THE PRINCIPAL EIGENVALUE OF THE
 p -LAPLACIAN AND THE p -BIHARMONIC OPERATOR

JIŘÍ BENEDIKT, Plzeň

(Received October 14, 2013)

Abstract. We survey recent results concerning estimates of the principal eigenvalue of the Dirichlet p -Laplacian and the Navier p -biharmonic operator on a ball of radius R in \mathbb{R}^N and its asymptotics for p approaching 1 and ∞ .

Let p tend to ∞ . There is a critical radius R_C of the ball such that the principal eigenvalue goes to ∞ for $0 < R \leq R_C$ and to 0 for $R > R_C$. The critical radius is $R_C = 1$ for any $N \in \mathbb{N}$ for the p -Laplacian and $R_C = \sqrt{2N}$ in the case of the p -biharmonic operator.

When p approaches 1, the principal eigenvalue of the Dirichlet p -Laplacian is $NR^{-1} \times (1 - (p-1) \log R(p-1)) + o(p-1)$ while the asymptotics for the principal eigenvalue of the Navier p -biharmonic operator reads $2N/R^2 + O(-(p-1) \log(p-1))$.

Keywords: eigenvalue problem for p -Laplacian; eigenvalue problem for p -biharmonic operator; estimates of principal eigenvalue; asymptotic analysis

MSC 2010: 35J66, 35J92, 35P15, 35P30

1. p -LAPLACIAN

Let us consider the eigenvalue problem for the Dirichlet p -Laplacian

$$(1.1) \quad \begin{cases} -\operatorname{div}(|\nabla u|^{p-2} \nabla u) = \lambda |u|^{p-2} u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

where $p > 1$ and Ω is a bounded open subset of \mathbb{R}^N , $N \geq 1$. It is well-known that the principal eigenvalue of (1.1) is

$$(1.2) \quad \lambda_1(\Omega, p) \stackrel{\text{def}}{=} \min \left(\int_{\Omega} |\nabla u|^p \, dx / \int_{\Omega} |u|^p \, dx \right)$$

where the minimum is taken over all $u \in W_0^{1,p}(\Omega)$, $u \neq 0$.

The author was supported by the Grant Agency of the Czech Republic, Grant No. 13-00863S.

In the one dimensional case $N = 1$ the precise formula

$$(1.3) \quad \lambda_1((-R, R), p) = \frac{1}{R^p}(p-1) \left(\frac{\pi}{p \sin(\pi/p)} \right)^p, \quad p > 1$$

is known (see, e.g., [7], page 244). It implies

$$\lim_{p \rightarrow 1+} \lambda_1((-R, R), p) = \frac{1}{R}, \quad \lim_{p \rightarrow 1+} \frac{\lambda_1((-R, R), p) - 1/R}{p-1} = \infty,$$

and

$$\begin{aligned} 0 < R \leq 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1((-R, R), p) = \infty, \\ R > 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1((-R, R), p) = 0 \end{aligned}$$

(see Figure 1).

When $N \geq 2$, an explicit formula for $\lambda_1(\Omega, p)$ is not known even in the case when $\Omega = B_N(0, R)$, the open ball of radius $R > 0$ and centered at the origin. Using the Cheeger constant, Kawohl and Fridman [14], Remark 5, proved the lower estimate

$$(1.4) \quad \lambda_1(B_N(0, R), p) \geq \left(\frac{N}{Rp} \right)^p, \quad p > 1$$

which together with (1.2) implies (see [14], Corollary 6)

$$\lim_{p \rightarrow 1+} \lambda_1(B_N(0, R), p) = \frac{N}{R}.$$

A more precise asymptotics for $\lambda_1(B_N(0, R), p)$ as $p \rightarrow 1+$ follows from the estimates

$$(1.5) \quad \frac{N}{R} \left(\frac{p'}{R} \right)^{p-1} \leq \lambda_1(B_N(0, R), p) \leq \frac{N}{R} \left(\frac{p'}{R} \right)^{p-1} \frac{\Gamma(p+1+N/p')}{\Gamma(p+1)\Gamma(2+N/p')}, \quad p > 1$$

where Γ is the Gamma function and $p' \stackrel{\text{def}}{=} p/(p-1)$. The estimate from below was proved in ([8], (8.10) on page 332) and both the estimates from below and from above in [3]. The proof of the estimate from below is based on the Picone identity [1], the estimate from above follows from (1.2) by choosing an appropriate function u .

Moreover, it is proved in [3] that the estimates (1.5) yield the asymptotics

$$\lambda_1(B_N(0, R), p) = \frac{N}{R} (1 - (p-1) \log R(p-1)) + o(p-1) \quad \text{as } p \rightarrow 1+.$$

This follows from the fact that both the lower and the upper bound in (1.5) are subject to the same asymptotics.




Figure 1. Dependence of λ_1 on p —second-order case.

On the other hand, it follows from [12], Lemma 1.5, that

$$\begin{aligned} 0 < R < 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = \infty, \\ R > 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = 0. \end{aligned}$$

The critical case $R = R_C \stackrel{\text{def}}{=} 1$ is not covered. In [5] we proved the estimates

$$(1.6) \quad \frac{Np}{R^p} \leq \lambda_1(B_N(0, R), p) \leq \frac{(p+1)(p+2)\dots(p+N)}{N!R^p}, \quad p > 1$$

which imply that, similarly to the one dimension,

$$\begin{aligned} 0 < R \leq 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = \infty, \\ R > 1 &\Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = 0. \end{aligned}$$

The estimates (1.6) can also be generalized to domains Ω other than a ball. Since the variational characterization (1.2) implies that $\lambda_1(\Omega, p)$ is decreasing with respect to Ω (in the sense of the set inclusion), the upper estimate in (1.6) applies to any bounded open subset of \mathbb{R}^N that contains an inscribed ball of radius $R > 0$ as well. On the other hand, it follows from the Schwarz symmetrization (see [13]) that the lower estimate in (1.6) holds also for any Ω such that $|\Omega| = |B_N(0, R)|$. Moreover, it is proved in [5] that

$$\lambda_1(\Omega, p) \geq \frac{kp}{R^p}$$

for any $\Omega \subset B_k(0, R) \times \mathbb{R}^{N-k}$ where $B_k(0, R)$ is the open ball in \mathbb{R}^k of radius $R > 0$ and centered at the origin, $k \in \{1, 2, \dots, N\}$. In particular, for $k = 1$ and $R = 1$ it implies $\lim_{p \rightarrow \infty} \lambda_1(\Omega, p) = \infty$ for any Ω situated between two parallel hyperplanes of distance 2. However, if Ω cannot be squeezed between two parallel hyperplanes of distance 2 but the radius of the largest inscribed ball has the radius $R \leq 1$, the asymptotic behavior of $\lambda_1(\Omega, p)$ as $p \rightarrow \infty$ is an *open problem*. A concrete example of such Ω in the plane is the open equilateral triangle with the largest inscribed disc of the radius 1.

In Figure 1 we present estimates of the principal eigenvalue $\lambda_1(B_N(0, R), p)$ in different dimensions $N = 1, 2, 3, 4, 8, 9$. The solid curve for $N = 1$ depicts the exact value (1.3). For $N = 2, 3$ and 4 the thick dots represent approximate values of λ_1 for certain discrete values of p , which were evaluated in [6]. The dashed curves represent lower and upper estimates from (1.5), the dotted curves visualize those from (1.6). Finally, the dash-dotted curves illustrate the lower estimate (1.4). The shaded regions reflect all the above mentioned estimates for $\lambda_1(B_N(0, R), p)$.

The well-known continuous embedding $W_0^{1,p}(B_N(0, R)) \hookrightarrow L^p(B_N(0, R))$ and the Rellich-Kondrachov Theorem (e.g., [9], Theorem 1.2.28) imply the existence of the minimal constant $C = C(p, N, R) = \lambda_1^{-1/p}(B_N(0, R), p)$ such that for all $u \in W_0^{1,p}(B_N(0, R))$

$$\|u\|_p \leq C(p, N, R) \|u\|_{1,p}$$

where

$$\|u\|_p \stackrel{\text{def}}{=} \left(\int_{B_N(0, R)} |u|^p dx \right)^{1/p}$$

while

$$\|u\|_{1,p} \stackrel{\text{def}}{=} \left(\int_{B_N(0, R)} |\nabla u|^p dx \right)^{1/p}$$

is an equivalent (radially symmetric) norm on $W_0^{1,p}(B_N(0, R))$. It then follows from the estimates (1.5) and (1.6) that

$$\frac{R}{N^{1/p}(p')^{1/p'}} \left(\frac{\Gamma(p+1)\Gamma(2+N/p')}{\Gamma(p+1+N/p')} \right)^{1/p} \leq C(p, N, R) \leq \frac{R}{N^{1/p}(p')^{1/p'}}$$

and

$$R \left(\frac{N!}{(p+1)(p+2)\dots(p+N)} \right)^{1/p} \leq C(p, N, R) \leq \frac{R}{N^{1/p}p^{1/p}},$$

respectively. Consequently, for all $u \in W_0^{1,p}(B_N(0, R))$ we have

$$\|u\|_p \leq \frac{R}{N^{1/p} \max\{p^{1/p}, (p')^{1/p'}\}} \|u\|_{1,p}.$$

2. p -BIHARMONIC OPERATOR

We also study the Navier p -biharmonic (fourth-order) eigenvalue problem

$$(2.1) \quad \begin{cases} \Delta(|\Delta u|^{p-2}\Delta u) = \lambda|u|^{p-2}u & \text{in } B_N(0, R), \\ u = \Delta u = 0 & \text{on } \partial B_N(0, R) \end{cases}$$

where $p > 1$. The principal eigenvalue of (2.1) is

$$(2.2) \quad \lambda_1(B_N(0, R), p) \stackrel{\text{def}}{=} \min \frac{\int_{B_N(0, R)} |\Delta u|^p dx}{\int_{B_N(0, R)} |u|^p dx}$$

where the minimum is taken over all $u \in W^{2,p}(B_N(0, R)) \cap W_0^{1,p}(B_N(0, R))$, $u \neq 0$ (see [10]).

A precise formula for $\lambda_1(B_N(0, R), p)$ is not known even in one dimension. The estimates

$$(2.3) \quad \left(\frac{2N}{R^2}\right)^p \left(\frac{\sqrt{\pi}\Gamma(p')}{\Gamma(p'+1/2)} - \frac{1}{p'}\right)^{1-p} \\ \leq \lambda_1(B_N(0, R), p) \leq \left(\frac{2N}{R^2}\right)^p \left(\frac{2\Gamma(p'+1+N/2)}{N\Gamma(N/2)\Gamma(p'+1)}\right)^{p-1}, \quad p > 1$$

were proved in [2] using [4]. These estimates imply the asymptotics

$$\lambda_1(B_N(0, R), p) = \frac{2N}{R^2} + O(-(p-1)\log(p-1)) \quad \text{as } p \rightarrow 1+.$$

On the other hand, using the Picone identity for the p -biharmonic operator due to Jaroš [11] and the variational characterization (2.2), respectively, the lower and the upper estimate,

$$(2.4) \quad \left(\frac{2N}{R^2}\right)^p \frac{1}{\sqrt{\pi}\Gamma(p)/[\Gamma(p+1/2)] - 1/p} \\ \leq \lambda_1(B_N(0, R), p) \leq \left(\frac{2N}{R^2}\right)^p \frac{2\Gamma(p+1+N/2)}{N\Gamma(N/2)\Gamma(p+1)}$$

were proved in [4]. They yield that, similarly to the second-order case, there is a critical radius $R_C = \sqrt{2N}$ such that

$$0 < R \leq R_C \Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = \infty, \\ R > R_C \Rightarrow \lim_{p \rightarrow \infty} \lambda_1(B_N(0, R), p) = 0.$$

However, here the critical radius does depend on the dimension.

In Figure 2 we present estimates for the principal eigenvalue in different dimensions $N = 1, 2, 3,$ and 4 . The dashed curves represent lower and upper estimates from (2.3), the dotted curves visualize those from (2.4). The shaded regions reflect all the above mentioned estimates for λ_1 .

Again, the well-known continuous embedding $W^{2,p}(B_N(0, R)) \cap W_0^{1,p}(B_N(0, R)) \hookrightarrow L^p(B_N(0, R))$ and the Rellich-Kondrachov Theorem imply the existence of the minimal constant $C = C(p, N, R) = \lambda_1^{-1/p}(B_N(0, R), p)$ such that for all $u \in W^{2,p}(B_N(0, R)) \cap W_0^{1,p}(B_N(0, R))$

$$\|u\|_p \leq C(p, N, R) \|u\|_{2,p}$$

where

$$\|u\|_p \stackrel{\text{def}}{=} \left(\int_{B_N(0, R)} |u|^p dx \right)^{1/p}$$




Figure 2. Dependence of λ_1 on p —fourth-order case.

and

$$\|u\|_{2,p} \stackrel{\text{def}}{=} \left(\int_{B_N(0,R)} |\Delta u|^p dx \right)^{1/p}$$

is an equivalent (radially symmetric) norm on $W^{2,p}(B_N(0, R)) \cap W_0^{1,p}(B_N(0, R))$. It follows from the estimates (2.3) and (2.4) that

$$\frac{R^2}{2N} \left(\frac{N\Gamma(N/2)\Gamma(p'+1)}{2\Gamma(p'+1+N/2)} \right)^{1/p'} \leq C(p, N, R) \leq \frac{R^2}{2N} \left(\frac{\sqrt{\pi}\Gamma(p')}{\Gamma(p'+1/2)} - \frac{1}{p'} \right)^{1/p'}$$

and

$$\frac{R^2}{2N} \left(\frac{N\Gamma(N/2)\Gamma(p+1)}{2\Gamma(p+1+N/2)} \right)^{1/p} \leq C(p, N, R) \leq \frac{R^2}{2N} \left(\frac{\sqrt{\pi}\Gamma(p)}{\Gamma(p+1/2)} - \frac{1}{p} \right)^{1/p},$$

respectively. Consequently, for all $u \in W^{2,p}(B_N(0, R)) \cap W_0^{1,p}(B_N(0, R))$ we have

$$\|u\|_p \leq \frac{R^2}{2N} \min \left\{ \left(\frac{\sqrt{\pi}\Gamma(p)}{\Gamma(p+1/2)} - \frac{1}{p} \right)^{1/p}, \left(\frac{\sqrt{\pi}\Gamma(p')}{\Gamma(p'+1/2)} - \frac{1}{p'} \right)^{1/p'} \right\} \|u\|_{2,p}.$$

References

- [1] *W. Allegretto, Y. X. Huang*: A Picone's identity for the p -Laplacian and applications. *Nonlinear Anal., Theory Methods Appl.* *32* (1998), 819–830.
- [2] *J. Benedikt, P. Drábek*: Asymptotics for the principal eigenvalue of the p -biharmonic operator on the ball as p approaches 1. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* *95* (2014), 735–742.
- [3] *J. Benedikt, P. Drábek*: Asymptotics for the principal eigenvalue of the p -Laplacian on the ball as p approaches 1. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* *93* (2013), 23–29.
- [4] *J. Benedikt, P. Drábek*: Estimates of the principal eigenvalue of the p -biharmonic operator. *Nonlinear Anal., Theory Methods Appl., Ser. A, Theory Methods* *75* (2012), 5374–5379.
- [5] *J. Benedikt, P. Drábek*: Estimates of the principal eigenvalue of the p -Laplacian. *J. Math. Anal. Appl.* *393* (2012), 311–315.
- [6] *R. J. Biezuner, J. Brown, G. Ercole, E. M. Martins*: Computing the first eigenpair of the p -Laplacian via inverse iteration of sublinear supersolutions. *J. Sci. Comput.* *52* (2012), 180–201.
- [7] *R. J. Biezuner, G. Ercole, E. M. Martins*: Computing the first eigenvalue of the p -Laplacian via the inverse power method. *J. Funct. Anal.* *257* (2009), 243–270.
- [8] *H. Bueno, G. Ercole, A. Zumpano*: Positive solutions for the p -Laplacian and bounds for its first eigenvalue. *Adv. Nonlinear Stud.* *9* (2009), 313–338.
- [9] *P. Drábek, J. Milota*: *Methods of Nonlinear Analysis. Applications to Differential Equations.* Birkhäuser Advanced Texts: Basel Lehrbücher, Birkhäuser, Basel, 2007.
- [10] *P. Drábek, M. Ôtani*: Global bifurcation result for the p -biharmonic operator. *Electron. J. Differ. Equ. (electronic only)* *2001* (2001), No. 48, 19 pages.
- [11] *J. Jaroš*: Picone's identity for the p -biharmonic operator with applications. *Electron. J. Differ. Equ. (electronic only)* *2011* (2011), No. 122, 6 pages.
- [12] *P. Juutinen, P. Lindqvist, J. J. Manfredi*: The ∞ -eigenvalue problem. *Arch. Ration. Mech. Anal.* *148* (1999), 89–105.
- [13] *B. Kawohl*: *Rearrangements and Convexity of Level Sets in PDE.* Lecture Notes in Mathematics 1150, Springer, Berlin, 1985.
- [14] *B. Kawohl, V. Fridman*: Isoperimetric estimates for the first eigenvalue of the p -Laplace operator and the Cheeger constant. *Commentat. Math. Univ. Carol.* *44* (2003), 659–667.

Author's address: Jiří Benedikt, Department of Mathematics and New Technologies for the Information Society, Faculty of Applied Sciences, University of West Bohemia, Univerzita 22, 306 14 Plzeň, Czech Republic, e-mail: benedikt@kma.zcu.cz.