

Tai-Wei Chen; Chung-I Ho; Jyh-Haur Teh
 E_1 -degeneration and $d'd''$ -lemma

Commentationes Mathematicae Universitatis Carolinae, Vol. 57 (2016), No. 2, 155–162

Persistent URL: <http://dml.cz/dmlcz/145757>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2016

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

E_1 -degeneration and $d'd''$ -lemma

TAI-WEI CHEN, CHUNG-I HO, JYH-HAUR TEH*

Abstract. For a double complex (A, d', d'') , we show that if it satisfies the $d'd''$ -lemma and the spectral sequence $\{E_r^{p,q}\}$ induced by A does not degenerate at E_0 , then it degenerates at E_1 . We apply this result to prove the degeneration at E_1 of a Hodge-de Rham spectral sequence on compact bi-generalized Hermitian manifolds that satisfy a version of $d'd''$ -lemma.

Keywords: $\partial\bar{\partial}$ -lemma; Hodge-de Rham spectral sequence; E_1 -degeneration; bi-generalized Hermitian manifold

Classification: 55T05, 53C05

1. Introduction

Complex manifolds that satisfy the $\partial\bar{\partial}$ -lemma enjoy some nice properties such as they are formal manifolds ([DGMS]), their Bott-Chern cohomology, Aeppli cohomology and Dolbeault cohomology are all isomorphic. Compact Kähler manifolds are examples of such manifolds. The Hodge-de Rham spectral sequence $E_{*,*}^{*,*}$ of a complex manifold M is built from the double complex $(\Omega^{*,*}(M), \partial, \bar{\partial})$ of complex differential forms which relates the Dolbeault cohomology of M to the de Rham cohomology of M . It is well known that $E_1^{p,q}$ is isomorphic to $H^p(M, \Omega^q)$ and the spectral sequence $E_r^{*,*}$ converges to $H^*(M, \mathbb{C})$. The goal of this paper is to prove an algebraic version of the result that the $\partial\bar{\partial}$ -lemma implies the E_1 -degeneration of a Hodge-de Rham spectral sequence. The following is our main result.

Theorem 1.1. *If a double complex (A, d', d'') satisfies the $d'd''$ -lemma and the spectral sequence $\{E_r^{p,q}\}$ induced by A does not degenerate at E_0 , then it degenerates at E_1 .*

We define a spectral sequence that is analogous to the Hodge-de Rham spectral sequence of complex manifolds for bi-generalized Hermitian manifolds. Applying result above, we are able to show that for compact bi-generalized Hermitian manifolds that satisfy a version of $\partial\bar{\partial}$ -lemma, the sequence degenerates at E_1 .

2. Degeneration of a Hodge-de Rham spectral sequence

Definition 2.1. A spectral sequence is a sequence of differential bi-graded modules $\{(E_r^{*,*}, d_r)\}$ such that d_r is of degree $(r, 1 - r)$ and $E_{r+1}^{p,q}$ is isomorphic to $H^{p,q}(E_r^{*,*}, d_r)$.

Definition 2.2. A filtered differential graded module is an \mathbb{N} -graded module $A = \bigoplus_{k=0}^{\infty} A^k$, endowed with a filtration F and a linear map $d : A \rightarrow A$ satisfying

- (1) d is of degree 1: $d(A^k) \subset A^{k+1}$;
- (2) $d \circ d = 0$;
- (3) the filtered structure is descending:

$$A = F^0 A \supseteq F^1 A \supseteq \dots \supseteq F^k A \supseteq F^{k+1} A \supseteq \dots ;$$

- (4) the map d preserves the filtered structure: $d(F^k A) \subset F^k A$ for all k .

For $p, q, r \in \mathbb{Z}$, let

$$\begin{aligned} Z_r^{p,q} &= \left\{ \xi \in F^p A^{p+q} \mid d\xi \in F^{p+r} A^{p+q+1} \right\}, \quad Z_{\infty}^{p,q} = F^p A^{p+q} \cap \ker d \\ B_r^{p,q} &= F^p A^{p+q} \cap dF^{p-r} A^{p+q-1}, \quad B_{\infty}^{p,q} = F^p A^{p+q} \cap \text{Im}d \\ E_r^{p,q} &= \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}}, \quad E_{\infty}^{p,q} = \frac{F^p A^{p+q} \cap \ker d}{F^{p+1} A^{p+q} \cap \ker d + F^p A^{p+q} \cap \text{Im}d} \end{aligned}$$

with the convention $F^{-k} A^{p+q} = A^{p+q}$ and $A^{-k} = \{0\}$ for $k \geq 0$. Let $d_r : E_r^{p,q} \rightarrow E_r^{p+r,q-r+1}$ be the differential induced by $d : Z_r^{p,q} \rightarrow Z_r^{p+r,q-r+1}$.

Throughout this paper, we always assume that $A = \bigoplus_{p,q \geq 0} A^{p,q}$ is a double complex of vector spaces over some field with two maps $d'_{p,q} : A^{p,q} \rightarrow A^{p+1,q}$ and $d''_{p,q} : A^{p,q} \rightarrow A^{p,q+1}$ satisfying $d'_{p+1,q} d'_{p,q} = 0, d''_{p,q+1} d''_{p,q} = 0$ and $d'_{p,q+1} d''_{p,q} + d''_{p+1,q} d'_{p,q} = 0$ for all $p, q \geq 0$. To make notation cleaner, we allow p, q to be any integers by defining $A^{p,q} = 0$ for $p < 0$ or $q < 0$.

Let $A^k = \bigoplus_{p+q=k} A^{p,q}$. Define

$$F^p A^k = \bigoplus_{s=p}^k A^{s,k-s}.$$

For $p > k$, define $F^p A^k = \{0\}$. This gives a descending filtration on A^k .

Let $d = d' + d''$. The double complex (A, d', d'') then defines a filtered differential graded module (A, d, F) . Let $\{E_r^{p,q}\}$ be the corresponding spectral sequence. We are interested in the convergence of $E_r^{p,q}$.

Definition 2.3. Let $\{E_r^{p,q}\}$ be the spectral sequence associated to the double complex (A, d', d'') . If $d_s = 0$ for all $s \geq r$, then we say that $\{E_r^{p,q}\}$ or A degenerates at E_r .

The following simple lemmas will be used frequently.

Lemma 2.4. *If G' is a vector space and $H < G, H < H'$ are subspaces of G' , the natural map $\varphi : \frac{G}{H} \rightarrow \frac{G'}{H'}$ is injective if and only if $G \cap H' = H$, and is surjective if and only if $G' = G + H'$.*

Lemma 2.5. *Let $p, q, r \in \mathbb{Z}$. There are inclusions*

$$\begin{aligned} \dots \subset B_0^{p,q} \subset B_1^{p,q} \subset \dots \subset B_\infty^{p,q} \subset Z_\infty^{p,q} \subset \dots \subset Z_1^{p,q} \subset Z_0^{p,q} \subset \dots, \\ Z_{r-1}^{p+1,q-1} \subset Z_r^{p,q}, \quad B_{r+1}^{p+1,q-1} \subset Z_r^{p,q}, \quad d(Z_r^{p-r,q+r-1}) = B_r^{p,q}. \end{aligned}$$

Definition 2.6. Let $\alpha_{p,q,r} : E_{r+1}^{p,q} \rightarrow \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_r^{p,q}}$ be the map induced by the composition of inclusion and projection, and $\beta_{p,q,r} : E_r^{p,q} \rightarrow \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_r^{p,q}}$ be the map induced by the projection.

Proposition 2.7. *Let $r \in \mathbb{Z}$. Then*

- (1) $d_r = 0$ if and only if $\beta_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}$,
- (2) $d_r = 0$ implies that $\alpha_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}$.

PROOF: (1) We first note that the map $\beta_{p,q,r}$ is always surjective. By Lemma 2.4, $\beta_{p,q,r}$ is an isomorphism if and only if $Z_r^{p,q} \cap (Z_{r-1}^{p+1,q-1} + B_r^{p,q}) = Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$, or equivalently, $B_r^{p,q} \subseteq Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$. The map $d_r^{p-r,q+r-1} : E_r^{p-r,q+r-1} \rightarrow E_r^{p,q}$ is the zero map if and only if $\text{Im} d_r^{p-r,q+r-1} = \{0\}$. This is equivalent to $d(Z_r^{p-r,q+r-1}) = B_r^{p,q} \subseteq Z_{r-1}^{p+1,q-1} + B_{r-1}^{p,q}$, which is equivalent to $\beta_{p,q,r}$ being an isomorphism.

(2) We recall that the isomorphism $E_{r+1}^{p,q} \xrightarrow{\cong} H^{p,q}(E_r^{*,*}, d_r)$ (see [M, Proof of Theorem 2.6]) is induced from some canonical projections and inclusions. If $d_r = 0$, $H^{p,q}(E_r^{*,*}, d_r) \cong E_r^{p,q}$ and we have a commutative diagram

$$\begin{array}{ccc} E_{r+1}^{p,q} & \xrightarrow{\cong} & E_r^{p,q} \\ & \searrow \alpha_{p,q,r} & \swarrow \beta_{p,q,r} \\ & & \frac{Z_r^{p,q}}{Z_{r-1}^{p+1,q-1} + B_r^{p,q}} \end{array}$$

By (1), $\beta_{p,q,r}$ is an isomorphism and hence $\alpha_{p,q,r}$ is an isomorphism. □

Definition 2.8. Fix a pair of integers (p, q) . For nonzero

$$\xi = \sum_i \xi_i \in \bigoplus_{i \geq 0} A^{p+i,q-i}$$

where $\xi_i \in A^{p+i,q-i}$, let $i_0 = \min_i \{\xi_i \neq 0\}$. We call ξ_{i_0} the leading term of ξ and denote it as $\ell^{p,q}(\xi)$. We define $\ell^{p,q}(0) = 0$. For $r \geq 1, p, q \in \mathbb{Z}$, let $\mathcal{E}_r^{p,q}$ be the set of $\xi = \xi_0 + \xi_1 + \dots + \xi_{r-1}$ such that $\xi_i \in A^{p+i,q-i}, d\xi = d'\xi_{r-1} \notin \text{Im} d''$,

$\ell^{p,q}(\eta) \neq \xi_0$ for all d -closed η and let

$$\mathcal{E}_0^{p,q-1} := B_0^{p,q} - (Z_{-1}^{p+1,q-1} + B_{-1}^{p,q}).$$

Lemma 2.9. Fix $r_0 \geq 1$.

- (1) If the map $\alpha_{p,q,r}$ is an isomorphism for all $p, q \in \mathbb{Z}, r \geq r_0$, then $\mathcal{E}_r^{p,q} = \emptyset$ for all $p, q \in \mathbb{Z}, r \geq r_0$.
- (2) If the map α_{p,q,r_0} is not an isomorphism, then $\mathcal{E}_{r_0}^{p,q} \neq \emptyset$.

PROOF: Note that by Lemma 2.4, the surjectivity of $\alpha_{p,q,r}$ is equivalent to the condition

$$Z_r^{p,q} = Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1} + B_r^{p,q} = Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}.$$

(1) Suppose that $\alpha_{p,q,r}$ is an isomorphism for all $r \geq r_0$. Then $Z_i^{p,q} = Z_{i+1}^{p,q} + Z_{i-1}^{p+1,q-1}$ for all $i \geq r_0$. Assume that $\mathcal{E}_r^{p,q} \neq \emptyset$ for some $r \geq r_0, p, q \in \mathbb{Z}$. Let $\xi \in \mathcal{E}_r^{p,q}$. By definition, $Z_{q+2}^{p,q} = Z_{q+3}^{p,q} = \cdots = Z_{\infty}^{p,q}$. So we may take $j > r$ such that $Z_j^{p,q} = Z_{\infty}^{p,q}$. Note that $\xi \in Z_r^{p,q}$. Using the relation above, we may write $\xi = \eta_1 + \eta_2$ where $\eta_1 \in Z_j^{p,q}, \eta_2 \in Z_{j-2}^{p+1,q-1} + \cdots + Z_{r-1}^{p+1,q-1}$. Since $\ell^{p,q}(\xi) \neq 0$, by comparing the degrees of both sides of $\xi = \eta_1 + \eta_2$, we have $\ell^{p,q}(\xi) = \ell^{p,q}(\eta_1)$. But $d\eta_1 = 0$ which contradicts to the fact that $\ell^{p,q}(\xi)$ is not the leading term of any d -closed element.

(2) Fix $r \geq 1$. Suppose that $\alpha_{p,q,r}$ is not an isomorphism, then $Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1} \subsetneq Z_r^{p,q}$. Let

$$\xi = \xi_0 + \xi_1 + \cdots + \xi_k \in Z_r^{p,q} - (Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}) \text{ where } \xi_i \in A^{p+i,q-i}.$$

If $k > r - 1$, let $\xi' = \xi_r + \xi_{r+1} + \cdots + \xi_k \in F^{p+r}A^{p+q} \subset F^{p+1}A^{p+q}$. We have

$$d\xi' = d\xi_r + \cdots + d\xi_k \in F^{p+r}A^{p+q+1} = F^{(p+1)+(r-1)}A^{(p+1)+(q-1)+1}$$

which means that $\xi' \in Z_{r-1}^{p+1,q-1}$. Let $\xi'' = \xi - \xi'$. If $\xi'' \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$, then $\xi = \xi' + \xi'' \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$ which contradicts to our assumption. Therefore $\xi'' = \xi_0 + \cdots + \xi_{r-1} \in Z_r^{p,q} - (Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1})$. Hence we may assume $\xi = \xi_0 + \cdots + \xi_{r-1}$.

- (i) Since $\xi \in Z_r^{p,q}$, by definition, $d\xi \in F^{p+r}A^{p+q+1}$. But $d(\xi_0 + \cdots + \xi_{r-2}) + d''\xi_{r-1} \in A^{p,q+1} \oplus A^{p+1,q} \oplus \cdots \oplus A^{p+r-1,q-r+2}$. This forces $d(\xi_0 + \cdots + \xi_{r-2}) + d''\xi_{r-1} = 0$ and hence $d\xi = d'\xi_{r-1}$.
- (ii) If $d'\xi_{r-1} = d''\eta_r$ for some $\eta_r \in A^{p+r,q-r}$, then $d(\xi - \eta_r) = d'\xi_{r-1} - d'\eta_r - d''\eta_r = -d'\eta_r \in A^{p+r+1,q-r} \subset F^{p+(r+1)}A^{p+q+1}$. Hence $\xi - \eta_r \in Z_{r+1}^{p,q}$. Since $\eta_r \in F^pA^{p+q}$ and $d\eta_r \in A^{p+r,q-r+1} \oplus A^{p+r+1,q-r} \subset F^{(p+1)+(r-1)}A^{p+q+1}$, we have $\eta_r \in Z_{r-1}^{p+1,q-1}$. Therefore $\xi = (\xi - \eta_r) + \eta_r \in Z_{r+1}^{p,q} + Z_{r-1}^{p+1,q-1}$, which is a contradiction. Hence $d'\xi_{r-1} \notin \text{Im}d''$.
- (iii) If ξ_0 is the leading term of a d -closed form $\tau \in F^pA^{p+q}$, then $\xi - \tau \in F^{p+1}A^{p+q}$ and $d(\xi - \tau) = d\xi \in F^{p+r}A^{p+q+1} = F^{(p+1)+(r-1)}A^{p+q+1}$.

Hence $\xi - \tau \in Z_{r-1}^{p+1, q-1}$. Then $\xi = \tau + (\xi - \tau) \in Z_{\infty}^{p, q} + Z_{r-1}^{p+1, q-1} \subset Z_{r+1}^{p, q} + Z_{r-1}^{p+1, q-1}$, which is a contradiction.

Hence $\xi \in \mathcal{E}_r^{p, q}$. □

- Lemma 2.10.** (1) $\mathcal{E}_0^{p, q-1} = \emptyset$ if and only if $\beta_{p, q, 0}$ is an isomorphism.
 (2) For $r \geq 1$, if $\mathcal{E}_r^{p-r, q+r-1} = \emptyset$, then $\beta_{p, q, r}$ is an isomorphism.
 (3) For $r \geq 1$, if $\mathcal{E}_r^{p-r, q+r-1} \neq \emptyset$, then $\beta_{p, q, j}$ is not an isomorphism for $j = 1$ or r .

PROOF: We note that $\beta_{p, q, r}$ is an isomorphism if and only if $B_r^{p, q} \subset Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q}$.

(1) This follows from the definition.

(2) Assume that $\beta_{p, q, r}$ is not an isomorphism. Then there exists $\xi \in B_r^{p, q} - (Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q})$. So $\xi = d\eta$ for some $\eta \in F^{p-r}A^{p+q-1}$. Let

$$\eta = \eta_0 + \eta_1 + \cdots + \eta_k \text{ where } \eta_i \in A^{p-r+i, q+r-i-1}.$$

If $k \geq r$, let $\eta' = \eta_r + \cdots + \eta_k \in F^pA^{p+q-1} \subset F^{p-(r-1)}A^{p+q-1}$. Then $d\eta' \in F^pA^{p+q} \cap d(F^{p-(r-1)}A^{p+q-1}) = B_{r-1}^{p, q}$. If $d(\eta - \eta') \in Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q}$, then $\xi = d(\eta - \eta') + d\eta' \in Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q}$, which is a contradiction. So $d(\eta - \eta') \in B_r^{p, q} - (Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q})$. Hence we may assume $\xi = d\eta$ where $\eta = \eta_0 + \cdots + \eta_{r-1}$.

- (i) Comparing the degrees of ξ and $d\eta$, we see that $d\eta = d'\eta_{r-1}$.
- (ii) If $\eta_0 = 0$, then $\xi = d(\eta_1 + \cdots + \eta_{r-1}) \in F^pA^{p+q} \cap d(F^{p-(r-1)}A^{p+q-1}) = B_{r-1}^{p, q}$, which is a contradiction. So $\eta_0 \neq 0$.
- (iii) If η_0 is the leading term of a d -closed form η'' , $\eta - \eta'' \in F^{p-r+1}A^{p+q-1}$ and $\xi = d\eta = d(\eta - \eta'') \in d(F^{p-(r-1)}A^{p+q-1}) \cap F^pA^{p+q} = B_{r-1}^{p, q}$, which is a contradiction. Hence η_0 is not the leading term of any d -closed form.
- (iv) If $d'\eta_{r-1} \in \text{Im}d''$, $\xi = d\eta = d'\eta_{r-1} = -d''\eta_r$ for some $\eta_r \in A^{p, q-1}$, then $\xi = d'\eta_r - d\eta_r \in Z_{\infty}^{p+1, q-1} + B_0^{p, q} \subset Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q}$, which is a contradiction. Hence $d'\eta_{r-1} \notin \text{Im}d''$.

Therefore, $\eta \in \mathcal{E}_r^{p-r, q+r-1}$.

(3) Assume that $\mathcal{E}_r^{p-r, q+r-1} \neq \emptyset$. Let $\eta = \eta_0 + \cdots + \eta_{r-1} \in \mathcal{E}_r^{p-r, q+r-1}$ where $\eta_i \in A^{p-r+i, q+r-i-1}$. Since $d\eta \in B_r^{p, q}$, if $d\eta \notin Z_{r-1}^{p+1, q-1} + B_{r-1}^{p, q}$, $\beta_{p, q, r}$ is not an isomorphism. So we may assume $d\eta = d'\eta_{r-1} = \xi' + d\eta'$ where $\xi' \in Z_{r-1}^{p+1, q-1}$ and $d\eta' \in B_{r-1}^{p, q}$. Let $\eta' = \eta'_1 + \eta'_2 + \cdots + \eta'_i$, where $\eta'_i \in A^{p-r+i, q+r-1-i}$. The degree of $d'\eta_{r-1}$ is (p, q) , so by comparing degrees of both sides of $d'\eta_{r-1} = \xi' + d\eta'$, we get

$$d'\eta_{r-1} = d'\eta'_{r-1} + d''\eta'_r \text{ and } d''\eta'_{r-1} = 0.$$

If $d'\eta'_{r-1} \in \text{Im}d''$, then $d'\eta_{r-1} \in \text{Im}d''$ which contradicts to the fact that $\eta \in \mathcal{E}_r^{p-r, q+r-1}$. So $d'\eta'_{r-1} \notin \text{Im}d''$. Note that if η'_{r-1} is the leading term of a d -closed element τ , we may write $\tau = \eta'_{r-1} + \tau_r + \cdots + \tau_k$ for some $k > r-1$ and each $\tau_i \in A^{p-r+i, q+r-1-i}$. Then comparing the degrees of $d'\tau = -d''\tau$, we get $d'\eta_{r-1} = -d''\tau_r$ which contradicts to the fact that $d'\eta_{r-1} \notin \text{Im}d''$.

From the above verification, we see that $\eta'_{r-1} \in \mathcal{E}_1^{p-1,q}$. Assume that $d\eta'_{r-1} \in Z_0^{p+1,q-1} + B_0^{p,q}$. Write $d\eta'_{r-1} = \gamma + d\sigma$ where $\gamma = \gamma_1 + \gamma_2 + \dots \in Z_0^{p+1,q-1}$, $\gamma_i \in A^{p+i,q-i}$, $\sigma = \sigma_0 + \sigma_1 + \dots \in B_0^{p,q}$ and $\sigma_i \in A^{p+i,q-1-i}$. Since the degree of $d\eta'_{r-1}$ is (p, q) , comparing the degrees of both sides of $d\eta'_{r-1} = \gamma + d\sigma$, we get $d\eta'_{r-1} = d''\sigma_0$ which contradicts to the fact that $\eta'_{r-1} \in \mathcal{E}_1^{p-1,q}$. Therefore $d\eta'_{r-1} \notin Z_0^{p+1,q-1} + B_0^{p,q}$ and hence $\beta_{p,q,1}$ is not an isomorphism. \square

Theorem 2.11. *Suppose that $(A = \bigoplus_{p,q \geq 0} A^{p,q}, d', d'')$ is a double complex and $r \geq 1$. The spectral sequence $\{E_r^{p,q}\}$ induced by A degenerates at E_r but not at E_{r-1} if and only if the following conditions hold:*

- (1) $\mathcal{E}_k^{p,q} = \emptyset$ for all $p, q \in \mathbb{Z}, k \geq r$ and
- (2) $\mathcal{E}_{r-1}^{p,q} \neq \emptyset$ for some p, q .

PROOF: Suppose that $\{E_r^{p,q}\}$ degenerates at E_r but not at E_{r-1} for some $r \geq 1$. By Proposition 2.7(2), $\alpha_{p,q,i}$ is an isomorphism for all $p, q \in \mathbb{Z}, i \geq r$. Then by Lemma 2.9, $\mathcal{E}_i^{p,q} = \emptyset$ for all $p, q \in \mathbb{Z}, i \geq r$. Since $d_{r-1} \neq 0$, by Proposition 2.7(1), there are some $p, q \in \mathbb{Z}$ such that $\beta_{p,q,r-1}$ is not an isomorphism. Then by Lemma 2.10, $\mathcal{E}_{r-1}^{p-r+1,q+r-2} \neq \emptyset$.

Conversely, suppose that (1) and (2) hold. By Lemma 2.10, $\beta_{p,q,k}$ is an isomorphism for all $p, q \in \mathbb{Z}, k \geq r$. Then by Proposition 2.7, $d_k = 0$ for $k \geq r$. For the case $r = 1$, by definition, $\mathcal{E}_0^{p,q} \neq \emptyset$ implies that $\beta_{p,q+1,0}$ is not an isomorphism. And hence by Proposition 2.7, $d_0 \neq 0$. For the case $r \geq 2$, if $\beta_{p,q,r-1}$ is an isomorphism for all $p, q \in \mathbb{Z}$, by Proposition 2.7, $d_{r-1} = 0$. Then we have $d_k = 0$ for $k \geq r - 1$. By the proof above, $\mathcal{E}_k^{p,q} = \emptyset$ for $k \geq r - 1$. In particular, $\mathcal{E}_{r-1}^{p,q} = \emptyset$ for all $p, q \in \mathbb{Z}$ which contradicts to our assumption (2). Therefore there exist some p_0, q_0 such that $\beta_{p_0,q_0,r-1}$ is not an isomorphism. By Proposition 2.7, $d_{r-1} \neq 0$. \square

Definition 2.12. We say that a double complex (A, d', d'') satisfies the $d'd''$ -lemma at (p, q) if

$$\text{Im}d' \cap \ker d'' \cap A^{p,q} = \ker d' \cap \text{Im}d'' \cap A^{p,q} = \text{Im}d'd'' \cap A^{p,q}$$

and A satisfies the $d'd''$ -lemma if A satisfies the $d'd''$ -lemma at (p, q) for all (p, q) .

Now we can give a proof of the main result Theorem 1.1.

PROOF: Note that by definition, $d'd''$ -lemma implies that $\text{Im}d' \cap \ker d'' \cap A^{p,q} = \text{Im}d' \cap \text{Im}d'' \cap A^{p,q}$ for all p, q . Since $\{E_r^{p,q}\}$ does not degenerate at E_0 , $\beta_{p,q,0}$ is not an isomorphism for some p, q , hence by Lemma 2.10, $\mathcal{E}_0^{p,q-1} \neq \emptyset$. Assume that $\mathcal{E}_r^{p,q} \neq \emptyset$ for some $p, q \in \mathbb{Z}, r \geq 1$. Then there is $\alpha = \sum_{i=0}^{r-1} \alpha_i \in \mathcal{E}_r^{p,q}$ where $\alpha_i \in A^{p+i,q-i}$. From the condition $d\alpha = d'\alpha_{r-1}$, we have $d''\alpha_{r-1} = -d'\alpha_{r-2}$ and hence $d''d\alpha = -d'd''\alpha_{r-1} = 0$. So $d\alpha = d'\alpha_{r-1} \in (\text{Im}d' \cap \ker d'') \cap A^{p,q} = (\text{Im}d' \cap \text{Im}d'') \cap A^{p,q}$. But by the definition of $\mathcal{E}_r^{p,q}$, $d'\alpha_{r-1} \notin \text{Im}d''$ which leads to a contradiction. Therefore by Theorem 2.11, $\{E_r^{p,q}\}$ degenerates at E_1 . \square

In the following, we apply the main result to prove the E_1 -degeneration of a spectral sequence of bi-generalized Hermitian manifolds. We refer the reader to [G1], [C] for generalized complex geometry, and to [CHT] for bi-generalized complex manifolds. We give a brief recall here. A bi-generalized complex structure on a smooth manifold M is a pair $(\mathcal{J}_1, \mathcal{J}_2)$ where $\mathcal{J}_1, \mathcal{J}_2$ are commuting generalized complex structures on M . A bi-generalized complex manifold is a smooth manifold M with a bi-generalized complex structure. A bi-generalized Hermitian manifold $(M, \mathcal{J}_1, \mathcal{J}_2, \mathbb{G})$ is an oriented bi-generalized complex manifold $(M, \mathcal{J}_1, \mathcal{J}_2)$ with a generalized metric \mathbb{G} which commutes with \mathcal{J}_1 and \mathcal{J}_2 . We define

$$U^{p,q} := U_1^p \cap U_2^q$$

where $U_1^p, U_2^q \subset \Gamma(\Lambda^* TM \otimes \mathbb{C})$ are eigenspaces of $\mathcal{J}_1, \mathcal{J}_2$ associated to the eigenvalues ip and iq respectively and $TM = TM \oplus T^*M$ is the generalized tangent space. It can be shown that the exterior derivative d is an operator from $U^{p,q}$ to $U^{p+1,q+1} \oplus U^{p+1,q-1} \oplus U^{p-1,q+1} \oplus U^{p-1,q-1}$ and we write

$$\delta_+ : U^{p,q} \rightarrow U^{p+1,q+1}, \quad \delta_- : U^{p,q} \rightarrow U^{p+1,q-1}$$

for the projection of d into corresponding spaces.

Definition 2.13. On a bi-generalized Hermitian manifold M , there is a double complex $\{(A, d', d'')\}$ given by

$$A^{p,q} := U^{p+q,p-q}, \quad d' = \delta_+, \quad d'' = \delta_-.$$

We call the spectral sequence $\{E_*^{*,*}\}$ associated to this double complex the ∂_1 -Hodge-de Rham spectral sequence.

By Theorem 1.1, we have the following result.

Theorem 2.14. *Suppose that M is a compact bi-generalized Hermitian manifold which satisfies the $\delta_+\delta_-$ -lemma and has positive dimension. Then the ∂_1 -Hodge-de Rham spectral sequence degenerates at E_1 .*

Now we give a proof of the E_1 -degeneration of the ∂_1 -Hodge-de Rham spectral sequence.

PROOF: Since $\bigoplus_{p,q} U^{p,q} = \Omega^\bullet(M) \otimes \mathbb{C}$ (see [Ca07], p. 36) where $\Omega^\bullet(M)$ is the collection of smooth forms on M , some $U^{p,q}$ is not empty. The space $U^{p,q}$ is a $C^\infty(M, \mathbb{C})$ -module where $C^\infty(M, \mathbb{C})$ is the ring of complex-valued smooth functions on M , and M has positive dimension, therefore $U^{p,q}$ is an infinite dimensional complex vector space. If δ_- is a zero map, we have $H_{\delta_-}^{p,q}(M) = U^{p,q}$ for all p, q . But M is compact, this contradicts to the fact that $H_{\delta_-}^{p,q}(M)$ is finite dimensional ([CHT, Theorem 2.14, Corollary 3.11]). Hence δ_- is not the zero map and the spectral sequence does not degenerate at E_0 . Since we assume that M satisfies the $\delta_+\delta_-$ -lemma, by Theorem 1.1, the spectral sequence degenerates at E_1 . \square

Acknowledgment. The authors thank the referee for his/her extremely careful review which largely improves this paper.

REFERENCES

- [C] Cavalcanti G., *New aspects of the dd^c -lemma*, Oxford Univ. DPhil. thesis, arXiv:math/0501406v1[math.DG].
- [Ca07] Cavalcanti G., *Introduction to generalized complex geometry*,impa, 26-Coloquio Brasileiro de Matematica, 2007.
- [CHT] Chen T.W., Ho C.I., Teh J.H., *Aeppli and Bott-Chern cohomology for bigeneralized Hermitian manifolds and $d'd''$ -lemma*, J. Geom. Phys. **93** (2015), 40–51.
- [DGMS] Deligne P., Griffiths P., Morgan J., Sullivan D., *Real homotopy theory of Kähler manifolds*, Invent. Math. **29** (1975), no. 3, 245–274.
- [G1] Gualtieri M., *Generalized complex geometry*, Ann. of Math. **174** (2011), 75–123.
- [M] McCleary J., *A User's Guide to Spectral Sequences*, 2nd edition, Cambridge studies in advanced mathematics, 58, Cambridge University Press, Cambridge, 2001.

Tai-Wei Chen, Jyh-Haur Teh:

MATHEMATICS DEPARTMENT, NATIONAL TSING HUA UNIVERSITY, HSINCHU, TAIWAN

E-mail: d937203@oz.nthu.edu.tw

jyhaur@math.nthu.edu.tw

Chung-I Ho:

MATHEMATICS DEPARTMENT, NATIONAL TSING HUA UNIVERSITY, NATIONAL CENTER OF THEORETICAL SCIENCES, MATHEMATICAL DIVISION, HSINCHU, TAIWAN

E-mail: ciho@math.cts.nthu.edu.tw

(Received January 25, 2015, revised January 26, 2016)