

D. Guerrero Sánchez; Vladimir Vladimirovich Tkachuk  
Some applications of the point-open subbase game

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 58 (2017), No. 3, 383–395

Persistent URL: <http://dml.cz/dmlcz/146904>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2017

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## Some applications of the point-open subbase game

D. GUERRERO SÁNCHEZ<sup>1</sup>, V.V. TKACHUK<sup>2</sup>

*Abstract.* Given a subbase  $\mathcal{S}$  of a space  $X$ , the game  $PO(\mathcal{S}, X)$  is defined for two players  $P$  and  $O$  who respectively pick, at the  $n$ -th move, a point  $x_n \in X$  and a set  $U_n \in \mathcal{S}$  such that  $x_n \in U_n$ . The game stops after the moves  $\{x_n, U_n : n \in \omega\}$  have been made and the player  $P$  wins if  $\bigcup_{n \in \omega} U_n = X$ ; otherwise  $O$  is the winner. Since  $PO(\mathcal{S}, X)$  is an evident modification of the well-known point-open game  $PO(X)$ , the primary line of research is to describe the relationship between  $PO(X)$  and  $PO(\mathcal{S}, X)$  for a given subbase  $\mathcal{S}$ . It turns out that, for any subbase  $\mathcal{S}$ , the player  $P$  has a winning strategy in  $PO(\mathcal{S}, X)$  if and only if he has one in  $PO(X)$ . However, these games are not equivalent for the player  $O$ : there exists even a discrete space  $X$  with a subbase  $\mathcal{S}$  such that neither  $P$  nor  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$ . Given a compact space  $X$ , we show that the games  $PO(\mathcal{S}, X)$  and  $PO(X)$  are equivalent for any subbase  $\mathcal{S}$  of the space  $X$ .

*Keywords:* point-open game; subbase; winning strategy; players; discrete space; compact space; scattered space; measurable cardinal

*Classification:* Primary 54A25; Secondary 91A05, 54D30, 54D70

### 1. Introduction

The game we are going to study here is a slight variation of the well-known point-open game  $PO(X)$  that was defined and studied independently by Galvin [4] and Telgársky [8]. Given a topological space  $X$ , the game  $PO(X)$  is played on  $X$  as follows: the  $n$ -th move of the first player (from now on denoted by  $P$ ) consists in taking a point  $x_n \in X$ . The second player (called  $O$ ) answers by choosing an open set  $U_n \subset X$  with  $x_n \in U_n$ . The play is finished after  $\omega$ -many moves and  $P$  wins if  $\bigcup_{n \in \omega} U_n = X$ . If  $\bigcup_{n \in \omega} U_n \neq X$ , then  $O$  wins the play  $\{x_n, U_n : n \in \omega\}$ . The game  $PO(X)$  is said to be determined on a space  $X$  if one of the players has a winning strategy.

In the paper [7] Pawlikowski gave a complete description of spaces  $X$  of countable pseudocharacter in which the game  $PO(X)$  is undetermined: this happens if and only if  $X$  is uncountable and has the Rothberger property  $C''$ . In particular, the game  $PO(X)$  is undetermined on an uncountable set  $X \subset \mathbb{R}$  if and only if  $X$  is a  $C''$ -set. It follows from a result of Laver [6] that there exist models of ZFC

---

DOI 10.14712/1213-7243.2015.210

<sup>1</sup>Research supported by CONACyT grant MOD-ORD-28-2015 PCI-085-08-15 (Mexico).

<sup>2</sup>Research supported by CONACyT grant CB-2012-01-178103 (Mexico).

in which every  $C''$ -subset of  $\mathbb{R}$  is countable so it is consistent with ZFC that the game  $PO(X)$  is determined on every set  $X \subset \mathbb{R}$ .

Telgársky established in [8] that if  $X$  is a  $\sigma$ -Čech-complete or pseudocompact space then  $PO(X)$  is determined on  $X$ . Later in [9] he gave a ZFC example of a non-metrizable space  $X$  on which  $PO(X)$  is undetermined. Daniels and Gruenhage [2] as well as Baldwin [1] studied the point-open game of uncountable length.

In this paper we consider a variation  $PO(\mathcal{S}, X)$  of the game  $PO(X)$  where  $\mathcal{S}$  is a fixed subbase of the space  $X$ . The game  $PO(\mathcal{S}, X)$  is played exactly as  $PO(X)$  with the only difference that at every move Player  $O$  must pick an element of  $\mathcal{S}$ . Of course, the first question that must be answered about the game  $PO(\mathcal{S}, X)$  is how different it is from  $PO(X)$ . We will show that, for any subbase  $\mathcal{S}$ , Player  $P$  has a winning strategy in  $PO(\mathcal{S}, X)$  if and only if he has one in  $PO(X)$ . However, these games are not equivalent for Player  $O$ : there exists even a discrete space  $X$  with a subbase  $\mathcal{S}$  such that neither  $P$  nor  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$ . We also establish that a discrete space  $X$  of a measurable cardinality is determined: for any subbase  $\mathcal{S}$  in  $X$ , Player  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$ .

Given a compact space  $X$  and a subbase  $\mathcal{S}$  in  $X$ , we prove that Player  $O$  has a winning strategy in  $PO(\mathcal{S}, X)$  if and only if  $X$  is not scattered; since the same characterization holds for  $PO(X)$ , for any subbase  $\mathcal{S}$  of the space  $X$ , the games  $PO(\mathcal{S}, X)$  and  $PO(X)$  are equivalent for both players.

## 2. Notation and terminology

All spaces are assumed to be Tychonoff. Given a space  $X$ , the symbol  $\tau(X)$  denotes the topology of  $X$  and  $\tau^*(X) = \tau(X) \setminus \{\emptyset\}$ . If  $X$  is a space and  $A \subset X$ , then  $\tau(A, X) = \{U \in \tau(X) : A \subset U\}$ . As usual,  $\mathbb{R}$  is the set of reals; the set  $\omega \setminus \{0\}$  is denoted by  $\mathbb{N}$  and  $\mathbb{I} = [0, 1] \subset \mathbb{R}$ . The symbol  $\mathbb{D}$  stands for the two-point space  $\{0, 1\}$  with the discrete topology.

If  $\mathcal{P} = \{x_n, U_n : n \in \omega\}$  is a play in the point-open game on a space  $X$ , then  $\langle x_n, U_n : n \leq k \rangle$  is called an initial segment (or simply segment) of the play  $\mathcal{P}$  for any  $k \in \omega$ .

A strategy of Player  $P$  in the point-open game  $PO(X)$  on a space  $X$  is a function  $\sigma$  with values in  $X$  defined on the initial segments of  $PO(X)$  called  $\sigma$ -admissible; they are inductively defined as follows. The empty segment is  $\sigma$ -admissible; if  $n > 0$ , then a segment  $\langle x_0, U_0, \dots, x_n, U_n \rangle$  is  $\sigma$ -admissible if  $\langle x_0, U_0, \dots, x_{n-1}, U_{n-1} \rangle$  is  $\sigma$ -admissible and  $x_n = \sigma(x_0, U_0, \dots, x_{n-1}, U_{n-1})$ . The definition of a strategy  $s$  for Player  $O$  is analogous for  $s$ -admissible segments  $\langle x_0, U_0, \dots, x_{n-1}, U_{n-1}, x_n \rangle$ . A play  $\mathcal{P} = \{x_n, U_n : n \in \omega\}$  is called  $\sigma$ -admissible for a strategy  $\sigma$  of Player  $P$  if every initial segment of  $\mathcal{P}$  is  $\sigma$ -admissible; in this case we will also say that  $P$  applies the strategy  $\sigma$ . An  $s$ -admissible play for a strategy  $s$  of Player  $O$  is defined analogously. A strategy  $\sigma$  of Player  $P$  is *winning* on  $X$  if  $P$  wins in any  $\sigma$ -admissible play. Analogously, a strategy  $s$  of Player  $O$  is winning on  $X$  if  $O$  is the winner in any  $s$ -admissible play.

A game  $PO(X)$  or  $PO(\mathcal{S}, X)$  is *undetermined* on a space  $X$  if neither of the players  $P$  and  $O$  has a winning strategy in the respective game on  $X$ . If a game is considered on a space  $X$  and  $A$  is one of the players, then  $X$  is called *A-favorable* if  $A$  has a winning strategy on  $X$ . We say that a space  $X$  is *crowded* if it has no isolated points. The space  $X$  is *scattered* if every non-empty subspace of  $X$  has an isolated point. The rest of our notation is standard and the unexplained notions can be found in the book [3].

### 3. Point-open subbase game

The point-open subbase game requires the player  $O$  to pick larger sets than in the point-open game so it is formally easier to win for the player  $P$ . Our main purpose is to establish that the point-open subbase game is equivalent to the point-open game for the player  $P$  while it might not be equivalent for the player  $O$  even in a discrete space.

The following statement is evident.

#### 3.1 Proposition. For any space $X$ ,

- (a) if  $P$  has a winning strategy in the game  $PO(X)$ , then the same strategy is winning in the game  $PO(\mathcal{S}, X)$  for any subbase  $\mathcal{S}$  in the space  $X$ ;
- (b) if  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$  for some subbase  $\mathcal{S}$  in the space  $X$ , then the same strategy is winning in the game  $PO(X)$ .

Denote by  $FO(X)$  the game in which the first player (called  $F$ ) at the  $n$ -th move picks a finite set  $F_n \subset X$  and the second player (called  $O$ ) chooses an open set  $U_n \supset F_n$ . The play is finished after  $\omega$ -many moves and  $F$  wins if  $\bigcup_{n \in \omega} U_n = X$ ; otherwise  $O$  is the winner. The game  $FO(X)$  is equivalent to  $PO(X)$  for both players (see Corollary 4.3 and Corollary 4.4 of the paper [8]) so it can be used instead of  $PO(X)$  when it is convenient.

**3.2 Proposition.** Assume that  $X$  is a space and  $\mathcal{S}$  is a subbase in  $X$ . If Player  $P$  has a winning strategy in the game  $PO(\mathcal{S}, X)$ , then Player  $F$  has a winning strategy in the game  $FO(X)$ .

PROOF: Let  $\rho$  be a winning strategy of  $P$  in  $PO(\mathcal{S}, X)$ . For any finite set  $F \subset X$  and  $U \in \tau(F, X)$  fix a finite family  $\mathcal{A}(U, F) \subset \mathcal{S}$  such that for each  $x \in F$  there exists a subfamily  $\mathcal{B} \subset \mathcal{A}(U, F)$  with  $x \in \bigcap \mathcal{B} \subset U$ .

To construct a strategy  $\sigma$  for Player  $F$  in the game  $FO(X)$  take the point  $x_0 = \rho(\emptyset)$  and consider the set  $F_0 = \{x_0\}$ ; letting  $\sigma(\emptyset) = F_0$  we define the strategy for the first move of  $F$ . Given any  $U_0 \in \tau(F_0, X)$  define  $\sigma(F_0, U_0)$  to be the set  $F_1 = \{\rho(x_0, S) : S \in \mathcal{A}(U_0, F_0)\}$ .

Proceeding inductively, assume that  $n \in \mathbb{N}$  and the strategy  $\sigma$  has been defined for every move  $i \leq n$  in such a way that for any  $i < n$  and any  $\sigma$ -admissible initial segment  $\langle F_0, U_0, \dots, F_i, U_i \rangle$  we have the following property:

- (1) if a point  $x_j \in F_j$  and a set  $S_j \in \mathcal{A}(U_j, F_j)$  are chosen for every  $j \leq n$  in such a way that the segment  $\langle x_j, S_j : j \leq i \rangle$  is  $\rho$ -admissible, then  $\rho(x_0, S_0, \dots, x_i, S_i) \in F_{i+1}$ .

Given an arbitrary  $\sigma$ -admissible segment  $\langle F_0, U_0, \dots, F_{n-1}, U_{n-1}, F_n \rangle$  take any  $U_n \in \tau(F_n, X)$  and consider the family  $\mathcal{E} = \{I : I = \langle x_0, S_0, \dots, x_n, S_n \rangle$  is a  $\rho$ -admissible segment such that  $x_i \in F_i$  and  $S_i \in \mathcal{A}(U_i, F_i)$  for every  $i \leq n\}$ . It is clear that  $\mathcal{E}$  is finite so letting  $F_{n+1} = \sigma(F_0, U_0, \dots, F_n, U_n) = \{\rho(I) : I \in \mathcal{E}\}$  we define our strategy  $\sigma$  for the move  $n + 1$  and it is straightforward that the property (1) holds if we replace  $n$  with  $n + 1$ . Therefore the construction of our strategy  $\sigma$  is complete and the condition (1) is satisfied for any  $n \in \mathbb{N}$ .

To see that  $\sigma$  is winning, suppose that  $\{F_i, U_i : i \in \omega\}$  is a play in which  $F$  applies the strategy  $\sigma$  and there exists a point  $p \in X \setminus \bigcup_{n \in \omega} U_n$ . It follows from  $p \notin U_0$  and the definition of  $\mathcal{A}(U_0, F_0)$  that there exists  $S_0 \in \mathcal{A}(U_0, F_0)$  such that  $x_0 \in U_0$  and  $p \notin S_0$ . Proceeding by induction assume that, for some  $n \in \omega$ , we have a  $\rho$ -admissible initial segment  $\langle x_0, S_0, \dots, x_n, S_n \rangle$  such that  $x_i \in F_i \cap S_i$  while  $S_i \in \mathcal{A}(U_i, F_i)$  and  $p \notin S_i$  for every  $i \leq n$ . It follows from (1) that  $x_{n+1} = \rho(x_0, S_0, \dots, x_n, S_n) \in F_{n+1} \subset U_{n+1}$  so it follows from  $p \notin U_{n+1}$  that we can choose  $S_{n+1} \in \mathcal{A}(U_{n+1}, F_{n+1})$  such that  $x_{n+1} \in S_{n+1}$  and  $p \notin S_{n+1}$ .

Therefore our inductive procedure can be continued to construct a play  $\{x_i, S_i : i \in \omega\}$  in the game  $PO(\mathcal{S}, X)$  where  $P$  applies the strategy  $\rho$  and  $p \notin S_i$  for every  $i \in \omega$ . However, this implies that  $p \notin \bigcup_{i \in \omega} S_i$  which is a contradiction with the fact that  $\rho$  is a winning strategy. This shows that  $\bigcup_{n \in \omega} U_n = X$  and hence  $\sigma$  is also a winning strategy. □

**3.3 Theorem.** *If  $X$  is a space and  $\mathcal{S}$  is a subbase in  $X$ , then the games  $PO(X)$  and  $PO(\mathcal{S}, X)$  are equivalent for  $P$ , i.e., Player  $P$  has a winning strategy in the game  $PO(X)$  if and only if he has a winning strategy in the game  $PO(\mathcal{S}, X)$ .*

PROOF: Since the game  $FO(X)$  is equivalent to the game  $PO(X)$  for both players, the games  $PO(X)$  and  $PO(\mathcal{S}, X)$  are equivalent for Player  $P$  by Proposition 3.1(a) and Proposition 3.2. □

**3.4 Corollary.** *If  $PO(X)$  is undetermined on a space  $X$ , then so is  $PO(\mathcal{S}, X)$  for any subbase  $\mathcal{S}$  of the space  $X$ .*

PROOF: It suffices to observe that, for such a space  $X$ , Player  $P$  does not have a winning strategy by Theorem 3.3 and Player  $O$  has no winning strategy by Proposition 3.1(b). □

Recall that  $X$  is a  $P$ -space if every  $G_\delta$ -subset of  $X$  is open.

**3.5 Observations.** Telgársky constructed in [9] a Lindelöf  $P$ -space  $X$  on which  $PO(X)$  is undetermined. By Corollary 3.4, on the same space  $X$  the game  $PO(\mathcal{S}, X)$  is undetermined for any subbase  $\mathcal{S}$ .

In [7], a complete characterization was given by Pawlikowski for the game  $PO(X)$  to be undetermined on a space  $X$  of countable pseudocharacter. In particular, the game  $PO(M)$  is undetermined on a set  $M \subset \mathbb{R}$  if and only if  $|M| > \omega$  and  $M$  is a  $C''$ -set, i.e., for every sequence  $\{U_n : n \in \omega\}$  of open covers of  $M$ , there exists a sequence  $\{U_n : n \in \omega\} \subset \tau(X)$  such that  $U_n \in \mathcal{U}_n$  for each  $n \in \omega$  and  $\bigcup_{n \in \omega} U_n = M$ . Therefore the game  $PO(\mathcal{S}, M)$  is undetermined on a set

$M \subset \mathbb{R}$  for every subbase  $\mathcal{S}$  of  $M$  if  $M$  is a  $C'''$ -set. We will see later that the above implication cannot be reversed.

Telgársky proved in [8] that for every Lindelöf scattered space  $X$ , Player  $P$  has a winning strategy in the game  $PO(X)$ . He also established in [8] that a compact space  $X$  is scattered if and only if Player  $O$  has a winning strategy in  $PO(X)$ . As an immediate consequence, the game  $PO(X)$  is determined on the class of compact spaces. We will show that the same is true for the game  $PO(\mathcal{S}, X)$  whenever  $X$  is compact and  $\mathcal{S}$  is a subbase in  $X$ .

**3.6 Theorem.** *Assume that a space  $X$  has a pseudocompact crowded subspace. Then Player  $O$  has a winning strategy in  $PO(\mathcal{S}, X)$  for any subbase  $\mathcal{S}$  in the space  $X$ .*

PROOF: Let  $Y$  be a pseudocompact crowded subspace of  $X$ ; since  $\bar{Y}$  is also pseudocompact and crowded, we can consider that  $Y$  is closed in  $X$ . We will use the following trivial observation.

(2) If  $Z$  is a space and  $\mathcal{G}$  is a finite family of closed subsets of  $Z$  such that the interior of  $\bigcup \mathcal{G}$  is non-empty, then the interior of  $G$  is non-empty for some  $G \in \mathcal{G}$ .

The set  $Y$  is infinite being crowded, so for any point  $x \in X$  we can find a set  $U \in \tau^*(Y)$  such that  $x \notin \bar{U}$ . There exists a finite family  $\mathcal{F} \subset \mathcal{S}$  such that  $x \in \bigcap \mathcal{F} \subset X \setminus \bar{U}$ . It follows from  $\bar{U} \subset \bigcup \{X \setminus S : S \in \mathcal{F}\}$  that we can apply (2) to find a set  $V \in \tau^*(Y)$  such that  $\bar{V} \subset (X \setminus S) \cap U$ . This proves that

(3) for any point  $x \in X$ , if  $U$  is a non-empty open subset of  $Y$ , then we can find  $S \in \mathcal{S}$  and a non-empty open subset  $V$  of  $Y$  such that  $x \in S$  and  $\bar{V} \subset U \setminus S$ .

Now it is easy to construct a winning strategy  $\sigma$  for Player  $O$ . If  $P$  chooses a point  $x_0 \in X$ , we can apply (3) to find a set  $S_0 \in \mathcal{S}$  and  $U_0 \in \tau^*(Y)$  such that  $x_0 \in S_0 \subset X \setminus \bar{U}_0$ ; let  $\sigma(x_0) = S_0$ . Proceeding by induction assume that  $n \in \omega$  and the strategy  $\sigma$  is constructed for the first  $n$  moves in such a way that

(4) for any  $\sigma$ -admissible segment  $\langle x_0, S_0, \dots, x_n, S_n \rangle$  we have defined a family  $\{U_0, \dots, U_n\}$  of non-empty open subsets of  $Y$  such that  $U_i \cap S_i = \emptyset$  for every  $i \leq n$  and  $\bar{U}_{i+1} \subset U_i$  if  $i < n$ .

If the move of Player  $P$  is a point  $x_{n+1} \in X$ , then (3) can be applied again to find a set  $S_{n+1} \in \mathcal{S}$  and  $U_{n+1} \in \tau^*(Y)$  such that  $x_{n+1} \in S_{n+1}$ ,  $\bar{U}_{n+1} \subset U_n$  and  $U_{n+1} \cap S_{n+1} = \emptyset$ . Letting  $\sigma(x_0, S_0, \dots, x_n, S_n, x_{n+1}) = S_{n+1}$  we complete the definition of the strategy  $\sigma$  and it is immediate that (4) holds for all  $n \in \omega$ .

Finally, assume that  $\{x_i, S_i : i \in \omega\}$  is a  $\sigma$ -admissible play. The definition of  $\sigma$  implies existence of a sequence  $\{U_i : i \in \omega\} \subset \tau^*(Y)$  such that  $\bar{U}_{i+1} \subset U_i$  and  $U_i \cap S_i = \emptyset$  for every  $i \in \omega$ . It follows from pseudocompactness of  $Y$  that  $\bigcap_{n \in \omega} U_n \neq \emptyset$ . The property (4) guarantees that  $(\bigcap_{n \in \omega} U_n) \cap (\bigcup_{n \in \omega} S_n) = \emptyset$  so  $\bigcup_{n \in \omega} S_n \neq X$  and hence  $\sigma$  is a winning strategy.  $\square$

**3.7 Corollary.** *If  $X$  is a compact space and  $\mathcal{S}$  is a subbase of  $X$ , then the following conditions are equivalent:*

- (a)  $X$  is scattered;
- (b) Player  $P$  has a winning strategy in the game  $PO(\mathcal{S}, X)$ ;
- (c) Player  $O$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ .

PROOF: We have already mentioned that the implication (a) $\implies$ (b) is true for the game  $PO(X)$  (see [8, Corollary 9.5]) so it is true for  $PO(\mathcal{S}, X)$  by Theorem 3.3. The implication (b) $\implies$ (c) is trivial and (c) $\implies$ (a) is an immediate consequence of Theorem 3.6. □

It follows from Theorem 2 of the paper [4] and Theorem 3.3 that Player  $P$  has no winning strategy in the game  $PO(\mathcal{S}, X)$  if  $X$  is an uncountable space of countable pseudocharacter and  $\mathcal{S}$  is a subbase of  $X$ . The same conclusion follows from the main result of the paper of Pawlikowski [7].

In particular, if  $X$  is a discrete uncountable space, then Player  $P$  has no winning strategy in the game  $PO(\mathcal{S}, X)$  for any subbase  $\mathcal{S}$  in  $X$ ; for such an  $X$ , it is easy to see that Player  $O$  always has a winning strategy in the game  $PO(X)$ . We will show that this is not the case for the game  $PO(\mathcal{S}, X)$ .

**3.8 Theorem.** *Suppose that  $X \subset \mathbb{I}$  and for any compact  $K \subset \mathbb{I}$ , if  $K \subset X$  or  $K \subset \mathbb{I} \setminus X$ , then  $K$  is countable. Such  $X$  are called Bernstein sets and it is well known that they exist. Consider the families  $\mathcal{S}_0 = \{[0, x] \cap X : x \in X\}$  and  $\mathcal{S}_1 = \{[x, 1] \cap X : x \in X\}$ ; then  $\mathcal{S} = \mathcal{S}_0 \cup \mathcal{S}_1$  is a subbase for the discrete topology on  $X$  and neither of the players has a winning strategy in the game  $PO(\mathcal{S}, X)$ . In particular the discrete space  $X$  of cardinality  $\mathfrak{c}$  admits a subbase  $\mathcal{S}$  such that the game  $PO(\mathcal{S}, X)$  is undetermined on  $X$ .*

PROOF: It is trivial that  $\mathcal{S}$  is a subbase for the discrete topology on  $X$  so, from now on we provide  $X$  with the discrete topology. Observe first that Player  $P$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ , due to Theorem 3.3 and the fact that  $P$  has no winning strategy in the game  $PO(X)$  by [4, Theorem 2]. Striving for a contradiction, assume that Player  $O$  has a winning strategy  $\sigma$  in the game  $PO(\mathcal{S}, X)$ . In what follows “initial segment” or simply “segment” will mean “a  $\sigma$ -admissible segment of a play in  $PO(\mathcal{S}, X)$ .”

Given initial segments  $I = \langle x_0, S_0, \dots, x_n, S_n \rangle$  and  $I' = \langle y_0, T_0, \dots, y_m, T_m \rangle$  of a play in  $PO(\mathcal{S}, X)$ , we say that  $I'$  extends  $I$  if  $I \subset I'$ . For any initial segment  $I$  of a play in  $PO(\mathcal{S}, X)$  let  $\mathcal{E}(I) = \{J : J \supset I \text{ is an initial segment}\}$ . Let  $\mathcal{E} = \mathcal{E}(\emptyset)$  be the family of all initial segments of the game  $PO(\mathcal{S}, X)$ . Since the strategy  $\sigma$  is winning,

- (5) if  $I = \langle x_0, S_0, \dots, x_n, S_n \rangle \in \mathcal{E}$ , then  $H(I) = X \setminus \bigcup \{S_i : i \leq n\}$  is dense (with respect to the natural topology) in a non-trivial closed interval.

We claim that

- (6) for any initial segment  $I \in \mathcal{E}$ , there exist segments  $I_0, I_1 \in \mathcal{E}(I)$  such that  $\overline{H(I_0)} \cap \overline{H(I_1)} = \emptyset$  (the bar denotes the closure in  $\mathbb{I}$ ).

To see that the statement (6) is true assume that there exists a segment  $I = \langle x_0, S_0, \dots, x_n, S_n \rangle \in \mathcal{E}$  such that  $\overline{H(I_0)} \cap \overline{H(I_1)} \neq \emptyset$  for any  $I_0, I_1 \in \mathcal{E}(I)$ . It is

easy to see that this implies that  $F = \bigcap \{ \overline{H(J)} : J \in \mathcal{E}(I) \} \neq \emptyset$ ; fix a point  $r \in F$ . We have two cases to consider.

*Case 1.*  $r \in X$ . Let  $x_{n+1} = r$  and  $S_{n+1} = \sigma(x_0, S_0, \dots, x_n, S_n, x_{n+1})$ . We will inductively extend the segment  $I_0 = \langle x_0, S_0, \dots, x_{n+1}, S_{n+1} \rangle$  to a play  $\mathcal{P}$  in which  $O$  applies the strategy  $\sigma$ . We will only have to choose a point  $x_i$  and then the strategy  $\sigma$  will automatically give us the set  $S_i = \sigma(x_0, S_0, \dots, x_{i-1}, S_{i-1}, x_i)$  for any  $i > n + 1$ .

If  $i \geq n + 1$  and we have the segment  $I = \langle x_0, S_0, \dots, x_i, S_i \rangle$ , then it follows from  $I \in \mathcal{E}(I_0)$  and the fact that the strategy  $\sigma$  is winning, that the set  $H(I)$  is uncountable; since also  $r \in \overline{H(I)}$ , we can choose a point  $x_{i+1} \in H(I)$  such that  $|x_{i+1} - r| < 2^{-i}$ . If  $S_{i+1} = \sigma(x_0, S_0, \dots, x_i, S_i, x_{i+1})$  and  $S_{n+1}$  both belong to  $S_j$  for some  $j \in \mathbb{D}$ , then it follows from  $x_{i+1} \in S_{i+1} \setminus \overline{S_{n+1}}$  that  $S_{n+1} \subset S_{i+1}$  and  $r$  is not the endpoint of the set  $S_{i+1}$ ; this implies  $r \notin \overline{H(J)}$  for the segment  $J = \langle x_0, S_0, \dots, x_{i+1}, S_{i+1} \rangle$  which is a contradiction. Therefore, for some element  $j \in \mathbb{D}$ , we have  $S_{n+1} \in S_j$  and  $S_{i+1} \in S_{1-j}$ ; if  $S_{n+1} \cap S_{i+1} \neq \emptyset$ , then  $S_{n+1} \cup S_{i+1} = X$  which is impossible because the strategy  $\sigma$  is winning so  $S_{n+1} \cap S_{i+1} = \emptyset$  for any  $i > n$ .

Finally observe that the sequence  $\{x_i : i > n + 1\}$  converges to  $r$  and all of its elements remain on the same side from  $r$ ; this easily implies that  $\bigcup_{i \geq n+1} S_i = X$  which is again a contradiction with the fact that  $\sigma$  is a winning strategy.

*Case 2.*  $r \notin X$ . Choose a sequence  $\{x_i : i \geq n + 1\} \subset X$  which converges to  $r$  with the additional property that both sets  $\{i \geq n + 1 : x_i > r\}$  and  $\{i \geq n + 1 : x_i < r\}$  are infinite. If  $\{x_i, S_i : i \in \omega\}$  is the play where  $O$  applies the strategy  $\sigma$ , then  $r$  cannot be the endpoint of any  $S_i$ . Therefore, if  $i > n$  and  $r \in S_i$ , then  $r$  cannot belong to the closure of the set  $H(I)$  for  $I = \langle x_0, S_0, \dots, x_i, S_i \rangle$ ; this contradiction shows that  $[x_i, 1] \cap X \subset S_i \subset (r, 1]$  if  $x_i > r$  and  $[0, x_i] \cap X \subset S_i \subset [0, r)$  if  $x_i < r$ . As an immediate consequence,  $\bigcup_{i \in \omega} S_i = X$  which is once more a contradiction with the fact that  $\sigma$  is a winning strategy so the property (6) is proved.

Given any segment  $I = \langle x_0, S_0, \dots, x_n, S_n \rangle \in \mathcal{E}$  observe that  $\overline{H(I)}$  is an interval  $[a, b]$  for some  $a, b \in \mathbb{I}$  so we can choose a point  $x_{n+1} \in H(I) \cap [a, b]$  in such a way that the length each of the intervals  $[a, x_{n+1}]$  and  $[x_{n+1}, b]$  does not exceed  $\frac{2}{3}(b - a)$ . Repeating such a choice the necessary number of times we can see that the following stronger version of the property (6) holds:

(7) for any  $\varepsilon > 0$  and any initial segment  $I \in \mathcal{E}$ , there exist initial segments  $I_0, I_1 \in \mathcal{E}(I)$  such that  $\overline{H(I_0)} \cap \overline{H(I_1)} = \emptyset$  and the diameter of the set  $\overline{H(I_j)}$  is less than  $\varepsilon$  for every  $j \in \mathbb{D}$ .

Take any point  $z \in X$  and let  $I_\emptyset = \{z, \sigma(z)\}$ . Proceeding inductively, assume that  $n \in \omega$  and we have constructed an initial segment  $I_s$  for any  $s \in \bigcup \{\mathbb{D}^m : m \leq n\}$  in such a way that

(8) for any  $m \leq n$ , the family  $\{\overline{H(I_s)} : s \in \mathbb{D}^m\}$  is disjoint;

(9) if  $m \leq n$  and  $s \in \mathbb{D}^m$ , then the diameter of  $\overline{H(I_s)}$  does not exceed  $2^{-m}$ ;



(10) if  $s \subset t$ , then  $I_t$  is an extension of  $I_s$ .

For any  $s \in \mathbb{D}^n$  apply the property (7) to find extensions  $I'$  and  $I''$  of the segment  $I_s$  such that  $\text{diam}(\overline{H(I')}) < 2^{-n-1}$  and  $\text{diam}(\overline{H(I'')}) < 2^{-n-1}$  while  $\overline{H(I')} \cap \overline{H(I'')} = \emptyset$  and let  $I_{s \smallfrown 0} = I'$  and  $I_{s \smallfrown 1} = I''$ . This gives us the family  $\{I_s : s \in \bigcup\{\mathbb{D}^m : m \leq n + 1\}\}$  and it is immediate that (8)–(10) are still fulfilled if we replace  $n$  with  $n + 1$ . Therefore our inductive procedure can be continued to construct the family  $\{I_s : s \in \mathbb{D}^{<\omega}\}$  such that the conditions (8)–(10) are satisfied for all  $n \in \omega$ .

The set  $K_n = \bigcup\{\overline{H(I_s)} : s \in \mathbb{D}^n\}$  is compact and  $K_{n+1} \subset K_n$  for all  $n \in \omega$ ; it is standard to deduce from (8)–(10) that  $K = \bigcap\{K_n : n \in \omega\}$  is homeomorphic to the Cantor set. If  $x \in K$ , then there is a unique function  $f \in \mathbb{D}^\omega$  such that  $\{x\} = \bigcap\{\overline{H(I_{f|n})} : n \in \omega\}$ . The property (10) shows that there exists a play  $\mathcal{P} = \{x_n, S_n : n \in \omega\}$  in which  $O$  applied the strategy  $\sigma$  and  $I_{f|n}$  is an initial segment of  $\mathcal{P}$  for any  $n \in \omega$ . The equality  $\{x\} = \bigcap\{\overline{H(I_{f|n})} : n \in \omega\}$  shows that  $X \setminus \{x\} \subset \bigcup_{n \in \omega} S_n$ ; since the strategy  $\sigma$  is winning, we must have  $x \in X$ . This proves that  $K \subset X$  which is a contradiction.  $\square$

**3.9 Corollary.** *There exists a space  $X \subset \mathbb{I}$  such that  $PO(X)$  is determined on  $X$  but  $PO(\mathcal{S}, X)$  is undetermined for some subbase  $\mathcal{S}$  of  $X$ . In particular, Pawlikowski’s characterization does not hold for the game  $PO(\mathcal{S}, X)$ .*

PROOF: Let  $Z \subset \mathbb{I}$  be a set such that for any compact  $K \subset \mathbb{I}$ , if  $K \subset Z$  or  $K \subset \mathbb{I} \setminus Z$ , then  $K$  is countable. Since the Rothberger property  $C'''$  is trivially preserved by finite unions, both sets  $Z$  and  $\mathbb{I} \setminus Z$  cannot have the property  $C'''$  because  $\mathbb{I}$  does not have it. So, one of them, let us call it  $X$ , is not a  $C'''$ -set and hence Player  $O$  has a winning strategy in  $PO(X)$  by Pawlikowski’s theorem [7]. Therefore it suffices to show that Player  $O$  does not have a winning strategy in  $PO(\mathcal{S}, X)$  for some subbase  $\mathcal{S}$  in the space  $X$ .

Let  $\mathcal{Q}_0 = \{[0, x] \cap X : x \in X\}$  and  $\mathcal{Q}_1 = \{[x, 1] \cap X : x \in X\}$ ; by Theorem 3.8, Player  $O$  does not have a winning strategy in the game  $PO(\mathcal{Q}, X)$  for the family  $\mathcal{Q} = \mathcal{Q}_0 \cup \mathcal{Q}_1$ . Let  $\mathcal{S}_0 = \{[0, x] \cap X : x \in X\}$  and  $\mathcal{S}_1 = \{(x, 1] \cap X : x \in X\}$ ; since  $X$  is dense in  $\mathbb{I}$ , the family  $\mathcal{S} = \mathcal{S}_0 \cup \mathcal{S}_1$  is easily seen to be a subbase of  $X$ . Suppose that  $\sigma$  is a winning strategy in  $PO(\mathcal{S}, X)$ . If  $y \in X$  and  $y \in U = [0, x] \cap X$  for some  $x \in X$ , then let  $H(U, y) = [0, y] \cap X$ . Analogously, if  $y \in U = (x, 1] \cap X$  for some  $x \in X$ , then  $H(U, y) = [y, 1] \cap X$ .

Now, if we consider  $X$  to have the discrete topology, then it is easy to define inductively a strategy  $s$  for Player  $O$  in the game  $PO(\mathcal{Q}, X)$  in such a way that for any  $s$ -admissible segment  $I = \langle x_0, U_0, \dots, x_{n-1}, U_{n-1}, x_n \rangle$  there exists a  $\sigma$ -admissible segment  $J = \langle x_0, W_0, \dots, x_{n-1}, W_{n-1}, x_n \rangle$  for which  $U_i = H(W_i, x_i)$  for all  $i \leq n - 1$  and  $s(I) = H(\sigma(J), x_n)$ . The strategy  $s$  cannot be winning by Theorem 3.8 and hence there exists a  $\sigma$ -admissible play  $\{x_n, W_n : n \in \omega\}$  such that  $\bigcup_{n \in \omega} H(W_n, x_n) = X$ . Observing that  $H(W_n, x_n) \subset W_n$  for every  $n \in \omega$ , we conclude that  $\bigcup_{n \in \omega} W_n = X$  which is a contradiction.  $\square$

The referee observed that it would be interesting to find out for a space  $X$  of countable pseudocharacter what conditions a subbase  $\mathcal{S}$  in  $X$  must satisfy to guarantee that the game  $PO(\mathcal{S}, X)$  is undetermined on  $X$  if and only if  $X$  is a  $C''$ -space; this would generalize Pawlikowski's theorem from [7]. We do not know the answer to this question. The referee also asked what replaces the Rothberger property if there are no restrictions on a subbase  $\mathcal{S}$ . We cannot answer this question either but it is worth noting that it follows from Theorem 3.8 that for a discrete space  $X$  of cardinality  $\mathfrak{c}$ , the game  $PO(\mathcal{S}, X)$  is undetermined for some subbase  $\mathcal{S}$  in  $X$ . Therefore in this case the space  $X$  need not even be Lindelöf so if anything replaces the Rothberger property, it will be something very different.

Recall that a cardinal  $\kappa$  is called *measurable* if there exists a free  $\sigma$ -complete ultrafilter on  $\kappa$ .

**3.10 Theorem.** *If  $\kappa$  is a measurable cardinal and  $X$  is a discrete space of cardinality  $\kappa$ , then Player  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$  for any subbase  $\mathcal{S}$  of the space  $X$ .*

PROOF: Fix a free ultrafilter  $\mu$  on  $X$  which is  $\sigma$ -complete, i.e., closed under countable intersections and let  $\mathcal{S}$  be any subbase for the discrete topology on  $X$ . Given any  $n \in \omega$ , if at the  $n$ -th move Player  $P$  picks a point  $x_n \in X$ , then there is a finite family  $\mathcal{B}_n \subset \mathcal{S}$  such that  $\bigcap \mathcal{B}_n = \{x_n\}$ . If  $\mathcal{B}_n \subset \mu$  then  $\{x_n\} \in \mu$  which is a contradiction.

Therefore for any  $n \in \omega$  there exists  $S_n \in \mathcal{B}_n \setminus \mu$  and hence we can let  $\sigma(x_0, S_0, \dots, x_{n-1}, S_{n-1}, x_n) = S_n$ . If  $\{x_n, S_n : n \in \omega\}$  is a play where  $O$  applies  $\sigma$ , then  $X \setminus S_n \in \mu$  for any  $n \in \omega$ . The ultrafilter  $\mu$  being  $\sigma$ -complete, the set  $\bigcap_{n \in \omega} X \setminus S_n = X \setminus \bigcup_{n \in \omega} S_n$  belongs to  $\mu$  and hence  $X \neq \bigcup_{n \in \omega} S_n$  which shows that  $\sigma$  is a winning strategy for Player  $O$ .  $\square$

In the paper [4] Galvin introduced a game  $G^*(X)$  and proved that it is equivalent to  $PO(X)$  for both players. In  $G^*(X)$ , at the  $n$ -th move Player  $P$  chooses an open cover  $\mathcal{U}_n$  of the space  $X$  and  $O$  responds by taking a set  $U_n \in \mathcal{U}_n$ . As in  $PO(X)$ , Player  $P$  wins if  $\bigcup_{n \in \omega} U_n = X$ ; otherwise  $O$  is the winner. The following game  $CE(\mathcal{S}, X)$  is a modification of  $G^*(X)$  such that  $G^*(X) = CE(\mathcal{S}, X)$  for  $\mathcal{S} = \tau(X)$ . It follows from Theorem 3.8 that the games  $PO(X)$  and  $PO(\mathcal{S}, X)$  need not be equivalent for Player  $O$  so it is not immediately clear whether passing from  $G^*(X)$  to  $CE(\mathcal{S}, X)$  we must obtain a game equivalent to  $PO(\mathcal{S}, X)$ . However, we will show that the ideas from [4] still work for our modification and hence the game  $CE(\mathcal{S}, X)$  is equivalent to  $PO(\mathcal{S}, X)$  for both players.

**3.11 Definition.** Given a space  $X$  and a subbase  $\mathcal{S}$  in  $X$ , in the game  $CE(\mathcal{S}, X)$  we have Players  $C$  and  $E$  who at the  $n$ -th move take an open cover  $\mathcal{U}_n \subset \mathcal{S}$  of the space  $X$  and an element  $U_n \in \mathcal{U}_n$  respectively. The game stops after  $\omega$ -many moves are made and the play  $\{\mathcal{U}_n, U_n : n \in \omega\}$  is a win for Player  $E$  if  $\bigcup_{n \in \omega} U_n = X$ ; otherwise  $C$  is the winner.

**3.12 Theorem.** *Given a space  $X$  and a subbase  $\mathcal{S}$  of  $X$ ,*

- (a) Player  $P$  has a winning strategy in  $PO(\mathcal{S}, X)$  if and only if  $E$  has a winning strategy in the game  $CE(\mathcal{S}, X)$ ;
- (b) Player  $O$  has a winning strategy in  $PO(\mathcal{S}, X)$  if and only if  $C$  has a winning strategy in the game  $CE(\mathcal{S}, X)$ .

PROOF: (a) If Player  $P$  has a winning strategy in  $PO(\mathcal{S}, X)$ , then he has a winning strategy in  $PO(X)$  by Theorem 3.3. By [4, Theorem 1], Player  $E$  has a winning strategy in  $CE(\tau(X), X)$  which, evidently, implies that he has a winning strategy in  $CE(\mathcal{S}, X)$ .

Next assume that  $X$  is  $E$ -favorable and fix a winning strategy  $s$  for Player  $E$  in the game  $CE(\mathcal{S}, X)$ ; let  $\mathcal{S}(x) = \{S \in \mathcal{S} : x \in S\}$  for every  $x \in X$ . It turns out that

- (11) if  $I = \langle \mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n \rangle$  is an  $s$ -admissible initial segment of  $CE(\mathcal{S}, X)$  (which can be empty), then there exists a point  $p \in X$  such that for every set  $S \in \mathcal{S}(p)$ , there exists a cover  $\mathcal{U}(S) \subset \mathcal{S}$  of the space  $X$  such that  $S = s(I, \mathcal{U}(S))$ .

Indeed, assume that for any  $x \in X$  there exists a set  $S_x \in \mathcal{S}(x)$  such that  $S_x \neq s(I, \mathcal{U})$  for any cover  $\mathcal{U} \subset \mathcal{S}$  of the space  $X$ . Then  $\mathcal{U} = \{S_x : x \in X\} \subset \mathcal{S}$  is a cover of  $X$  and hence we have a point  $p \in X$  such that  $\sigma(I, \mathcal{U}) = S_p$ ; this contradiction proves that (11) holds.

Apply (11) to find  $x_0 \in X$  such that  $\mathcal{S}(x_0) \subset \{\sigma(\mathcal{U}) : \mathcal{U} \subset \mathcal{S} \text{ and } \bigcup \mathcal{U} = X\}$  and let  $\sigma(\emptyset) = x_0$ . If  $O$  plays  $U_0 \in \mathcal{S}(x_0)$ , then choose a cover  $\mathcal{U}_0 \subset \mathcal{S}$  such that  $U_0 = s(\mathcal{U}_0)$ . Suppose that  $n \in \omega$  and we have defined a strategy  $\sigma$  for the moves from 0 to  $n$  in such a way that for any  $\sigma$ -admissible initial segment  $\langle x_0, U_0, \dots, x_n, U_n \rangle$  of the game  $PO(\mathcal{S}, X)$  we have covers  $\mathcal{U}_0, \dots, \mathcal{U}_n$  of the space  $X$  such that the segment  $\langle \mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n \rangle$  is  $s$ -admissible. Apply (11) again to find  $x_{n+1} \in X$  such that  $\mathcal{S}(x_{n+1}) \subset \{s(\mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n, \mathcal{U}) : \mathcal{U} \subset \mathcal{S} \text{ and } \bigcup \mathcal{U} = X\}$  and let  $\sigma(x_0, U_0, \dots, x_n, U_n) = x_{n+1}$ . If Player  $O$  takes a set  $U_{n+1} \ni x_{n+1}$ , then we can choose a cover  $\mathcal{U}_{n+1} \subset \mathcal{S}$  of the space  $X$  such that  $U_{n+1} = s(\mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n, \mathcal{U}_{n+1})$ . This completes the definition of the strategy  $\sigma$ .

To see that  $\sigma$  is winning note that to any  $\sigma$ -admissible play  $\{x_n, U_n : n \in \omega\}$  we have associated an  $s$ -admissible play  $\{\mathcal{U}_n, U_n : n \in \omega\}$  so  $\bigcup_{n \in \omega} U_n = X$ , i.e., the strategy  $\sigma$  is winning. Therefore every  $E$ -favorable space is  $P$ -favorable. This completes the proof of (a).

(b) If Player  $O$  has a winning strategy  $\sigma$  in the game  $PO(\mathcal{S}, X)$ , then let  $\mathcal{U}_0 = \{\sigma(x) : x \in X\}$  and  $s(\emptyset) = \mathcal{U}_0$ . If  $E$  chooses a set  $U_0 \in \mathcal{U}_0$ , then there exists a point  $x_0 \in X$  such that  $U_0 = \sigma(x_0)$ ; consider the family  $\mathcal{U}_1 = \{\sigma(x_0, U_0, x) : x \in X\}$  and let  $s(\mathcal{U}_0, U_0) = \mathcal{U}_1$ . Proceeding inductively, assume that  $n \in \omega$  and the strategy  $s$  for Player  $C$  is defined for the moves from 0 to  $n$  in such a way that for any  $s$ -admissible initial segment  $\langle \mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n \rangle$  we have defined a set  $\{x_0, \dots, x_n\}$  such that the segment  $\langle x_0, U_0, \dots, x_n, U_n \rangle$  is  $\sigma$ -admissible.

Consider the family  $\mathcal{U}_{n+1} = \{\sigma(x_0, U_0, \dots, x_n, U_n, x) : x \in X\}$  and let  $s(\mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n) = \mathcal{U}_{n+1}$ ; if  $E$  answers with a set  $U_{n+1} \in \mathcal{U}_{n+1}$ , then choose

the point  $x_{n+1} \in X$  such that  $U_{n+1} = \sigma(x_0, U_0, \dots, x_n, U_n, x_{n+1})$ . This completes the construction of the strategy  $s$ .

To see that  $s$  is winning note that to any  $s$ -admissible play  $\{U_n, U_n : n \in \omega\}$  we have associated a  $\sigma$ -admissible play  $\{x_n, U_n : n \in \omega\}$  so  $\bigcup_{n \in \omega} U_n \neq X$ , i.e., the strategy  $s$  is winning. Therefore every  $O$ -favorable space is  $C$ -favorable.

If  $s$  is a winning strategy for Player  $C$ , then for any point  $x_0 \in X$  let  $\sigma(x_0)$  be an element  $U_0 \in \mathcal{U}_0 = s(\emptyset)$  that contains  $x_0$ . Suppose that  $n \in \omega$  and we have defined a strategy  $\sigma$  for the moves from 0 to  $n$  in such a way that for any  $\sigma$ -admissible initial segment  $\langle x_0, U_0, \dots, x_n, U_n \rangle$  of the game  $PO(\mathcal{S}, X)$  we have constructed open covers  $\mathcal{U}_0, \dots, \mathcal{U}_n \subset \mathcal{S}$  of the space  $X$  such that the segment  $\langle \mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n \rangle$  is  $s$ -admissible. For any point  $x_{n+1} \in X$  choose an element  $U_{n+1} \in \mathcal{U}_{n+1} = s(\mathcal{U}_0, U_0, \dots, \mathcal{U}_n, U_n)$  such that  $x_{n+1} \in U_{n+1}$ ; letting  $\sigma(x_0, U_0, \dots, x_n, U_n, x_{n+1}) = U_{n+1}$  we complete the definition of a strategy  $\sigma$ . To see that  $\sigma$  is winning observe that to any  $\sigma$ -admissible play  $\{x_n, U_n : n \in \omega\}$  we have associated an  $s$ -admissible play  $\{U_n, U_n : n \in \omega\}$  so  $\bigcup_{n \in \omega} U_n \neq X$ , i.e., the strategy  $\sigma$  is winning. Therefore every  $C$ -favorable space is  $O$ -favorable. This completes the proof of (b).  $\square$

**3.13 Corollary.** *Given a space  $X$  and a subbase  $\mathcal{S}$  in  $X$ , the games  $CE(\mathcal{S}, X)$  and Galvin's game  $G^*(X) = CE(\tau(X), X)$  are equivalent for Player  $E$ , i.e.,  $E$  has a winning strategy in  $CE(\mathcal{S}, X)$  if and only if he has one in  $G^*(X)$ .*

PROOF: It follows from [4, Theorem 1] that Player  $E$  has a winning strategy in the game  $G^*(X)$  if and only if  $P$  has a winning strategy in  $PO(X)$ . By Theorem 3.3 the game  $PO(X)$  is equivalent to  $PO(\mathcal{S}, X)$  for Player  $P$ . Applying Theorem 3.12 we can see that  $G^*(X)$  is equivalent to  $CE(\mathcal{S}, X)$  for Player  $E$ .  $\square$

#### 4. Open problems

A proof of a statement about discrete spaces usually involves no topology; it is all about set theory. Therefore most questions about discrete spaces belong more to set theory than to topology. In particular, this is the case when we consider the game  $PO(\mathcal{S}, X)$  on discrete spaces. The most intriguing fact is that the point-open subbase game might be useful for a purely set-theoretic task of characterizing measurable cardinals.

**4.1 Question.** *Suppose that  $X$  is a discrete space such that Player  $O$  has a winning strategy in the game  $PO(\mathcal{S}, X)$  for every subbase  $\mathcal{S}$  in  $X$ . Must the cardinality of  $X$  be measurable?*

**4.2 Question.** *Suppose that  $X$  is a discrete space of cardinality  $2^c$ . Does there exist a subbase  $\mathcal{S}$  in  $X$  for which Player  $O$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ ?*

**4.3 Question.** *Suppose that  $X$  is an uncountable discrete space whose cardinality is non-measurable. Does there exist a linear order  $<$  on the set  $X$  such that, for the subbase*

$$\mathcal{S} = \{\{y \in X : y \leq x\} : x \in X\} \cup \{\{y \in X : x \leq y\} : x \in X\},$$

Player  $O$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ ?

**4.4 Question.** Suppose that  $X$  is a discrete space of uncountable cardinality such that Player  $O$  has no winning strategy in the game  $PO(\mathcal{B}, X)$  for some subbase  $\mathcal{B}$  in  $X$ . Does there exist a linear order  $<$  on the set  $X$  such that, for the subbase

$$\mathcal{S} = \{\{y \in X : y \leq x\} : x \in X\} \cup \{\{y \in X : x \leq y\} : x \in X\},$$

Player  $O$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ ?

**4.5 Question.** Does there exist a pseudocompact space  $X$  such that the games  $PO(X)$  and  $PO(\mathcal{S}, X)$  are not equivalent for Player  $O$  for some subbase  $\mathcal{S}$  in the space  $X$ ?

**4.6 Question.** Does there exist a countably compact space  $X$  such that the games  $PO(X)$  and  $PO(\mathcal{S}, X)$  are not equivalent for Player  $O$  for some subbase  $\mathcal{S}$  in the space  $X$ ?

**4.7 Question.** Given a maximal almost disjoint family  $\mathcal{N}$  on  $\omega$  let  $X = \omega \cup \mathcal{N}$  be the Mrowka space determined by  $\mathcal{N}$  (see [3, Example 3.6.I(a)]). Does there exist a subbase  $\mathcal{S}$  in  $X$  such that Player  $O$  has no winning strategy in the game  $PO(\mathcal{S}, X)$ ?

**4.8 Question.** Suppose that  $X$  is an uncountable second countable space such that every compact subspace of  $X$  is countable. Is it true that the game  $PO(\mathcal{S}, X)$  is undetermined for some subbase  $\mathcal{S}$  of the space  $X$ ?

**4.9 Question.** Suppose that  $X$  is an uncountable space with a countable network such that every compact subspace of  $X$  is countable. Is it true that the game  $PO(\mathcal{S}, X)$  is undetermined for some subbase  $\mathcal{S}$  of the space  $X$ ?

**4.10 Question.** Suppose that  $X$  is an uncountable hereditarily Lindelöf space such that every compact subspace of  $X$  is countable. Is it true that the game  $PO(\mathcal{S}, X)$  is undetermined for some subbase  $\mathcal{S}$  of the space  $X$ ?

## REFERENCES

- [1] Baldwin S., *Possible point-open types of subsets of the reals*, Topology Appl. **38** (1991), 219–223.
- [2] Daniels P., Gruenhage G., *The point-open types of subsets of the reals*, Topology Appl. **37** (1990), no. 1, 53–64.
- [3] Engelking R., *General Topology*, PWN, Warszawa, 1977.
- [4] Galvin F., *Indeterminacy of point-open games*, Bull. Acad. Polon. Sci. Sér. Math. **26** (1978), no. 5, 445–449.
- [5] Kuratowski K., *Topology, Volume 1*, Academic Press, New York, 1966.
- [6] Laver R., *On the consistency of Borel's conjecture*, Acta Math. **137** (1976), 151–169.
- [7] Pawlikowski J., *Undetermined sets of point-open games*, Fund. Math. **144** (1994), 279–285.
- [8] Telgársky R., *Spaces defined by topological games*, Fund. Math. **88** (1975), 193–223.

- [9] Telgársky R., *Spaces defined by topological games, II*, Fund. Math. **116** (1983), no. 3, 189–207.

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA METROPOLITANA, AV. SAN RAFAEL ATLIXCO, 186, COL. VICENTINA, IZTAPALAPA, C.P. 09340, MEXICO D.F., MEXICO

*E-mail:* dgs@ciencias.unam.mx  
vova@xanum.uam.mx

(Received March 1, 2016, revised April 12, 2017)