

Mehdi Parsinia

On the mappings \mathcal{Z}_A and \mathcal{S}_A in intermediate rings of $C(X)$

Commentationes Mathematicae Universitatis Carolinae, Vol. 59 (2018), No. 3, 383–390

Persistent URL: <http://dml.cz/dmlcz/147405>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2018

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

On the mappings \mathcal{Z}_A and \mathfrak{S}_A in intermediate rings of $C(X)$

MEHDI PARSINIA

Abstract. In this article, we investigate new topological descriptions for two well-known mappings \mathcal{Z}_A and \mathfrak{S}_A defined on intermediate rings $A(X)$ of $C(X)$. Using this, coincidence of each two classes of z -ideals, \mathcal{Z}_A -ideals and \mathfrak{S}_A -ideals of $A(X)$ is studied. Moreover, we answer five questions concerning the mapping \mathfrak{S}_A raised in [J. Sack, S. Watson, *C and C^* among intermediate rings*, Topology Proc. **43** (2014), 69–82].

Keywords: z -ideal; \mathcal{Z}_A -ideal; \mathfrak{S}_A -ideal; z -filter; \mathcal{Z}_A -filter; \mathfrak{S}_A -filter; intermediate ring

Classification: 54C30, 46E25

1. Introduction

Throughout this article all topological spaces are assumed to be Tychonoff. For a given topological space X , $C(X)$ denotes the algebra of all real-valued continuous functions on X and $C^*(X)$ denotes the subalgebra of $C(X)$ consisting of all bounded elements. A subring of $C(X)$ containing $C^*(X)$ is called an intermediate ring. An intermediate ring which is isomorphic with $C(Y)$ for some Tychonoff space Y is called an intermediate C -ring. The reader is referred to [5] for undefined terms and notations concerning $C(X)$. For each element f of an intermediate ring $A(X)$, we set $S_A(f) = \{p \in \beta X : (fg)^*(p) = 0, \forall g \in A(X)\}$. As stated in [11], we could easily observe that $S_A(fg) = S_A(f) \cup S_A(g)$, $S_A(f^2 + g^2) = S_A(f) \cap S_A(g)$ and $S_A(f^n) = S_A(f)$ for each $f, g \in A(X)$ and each $n \in \mathbb{N}$. Also, $S_C(f) = \text{cl}_{\beta X} Z(f)$ for each $f \in C(X)$ and $S_{C^*}(f) = Z(f^\beta)$ for each $f \in C^*(X)$. Moreover, $\text{cl}_{\beta X} Z(f) \subseteq S_A(f) \subseteq Z(f^*)$ and thus $S_A(f) \cap X = Z(f)$ for each $f \in A(X)$. For each $p \in \beta X$, we use M_A^p (or O_A^p) to denote the set $\{f \in A(X) : p \in S_A(f)\}$ ($\{f \in A(X) : p \in \text{int}_{\beta X} S_A(f)\}$), respectively. Evidently, $M_C^p = M^p$ (or $O_C^p = O^p$) and $M_{C^*}^p = M^{*p}$ ($O_{C^*}^p = O^{*p}$), respectively. Moreover, we can see that $\text{int}_{\beta X} S_A(f) = \text{int}_{\beta X} \text{cl}_{\beta X} Z(f)$ for each $f \in A(X)$ and thus $O_A^p = O^p \cap A(X)$ for each $p \in \beta X$. An ideal I of a commutative ring R is called a z -ideal if $M_f(R) \subseteq I$ whenever $f \in I$ in which $M_f(R)$ denotes the intersection of all the maximal ideals of R containing f . It is well-known that $M_f(C(X)) = \{g \in C(X) : Z(f) \subseteq Z(g)\}$ for each $f \in C(X)$. Also, from [9, Proposition 2.7], it follows that $M_f(A(X)) = \{g \in A(X) : S_A(f) \subseteq S_A(g)\}$ for each element f of an intermediate ring $A(X)$. Therefore, an ideal I in $C(X)$

(or in an intermediate ring $A(X)$) is a z -ideal if and only if whenever $Z(f) \subseteq Z(g)$ (or $S_A(f) \subseteq S_A(g)$), where $f \in I$ and $g \in C(X)$ ($g \in A(X)$, respectively), then $g \in I$. The aim of this paper is to answer the five basic questions concerning the mapping \mathfrak{S}_A which have been raised in [13]. This is done by investigating new topological descriptions for two well-known mappings \mathcal{Z}_A and \mathfrak{S}_A defined on intermediate rings of $C(X)$. This paper consists of 3 sections. Section 1, which is already noticed, is the introduction. In Section 2, we study the mapping \mathcal{Z}_A which was first introduced in [12] and more studied in [3]. Moreover, coincidence of z -ideals of $A(X)$ with \mathcal{Z}_A -ideals and z_A -ideals is studied. In Section 3, we study the mapping \mathfrak{S}_A which was first introduced in [8] and further studied in [14] and [13]. We establish a topological description for this mapping. Coincidence of \mathfrak{S}_A -ideals with z -ideals and \mathcal{Z}_A -ideals are studied. Moreover, the five questions concerning the mapping \mathfrak{S}_A raised in [13] are answered.

2. The mapping \mathcal{Z}_A

The mapping \mathcal{Z}_A was first introduced in [12] on an intermediate ring $A(X)$ as follows: for each element f of $A(X)$,

$$\mathcal{Z}_A(f) = \{E \in Z(X) : \exists g \in A(X), fg|_{X \setminus E} = 1\}.$$

By the following statement we provide a new topological description for \mathcal{Z}_A .

Theorem 2.1. *For each element f of an intermediate ring $A(X)$, we have*

$$\mathcal{Z}_A(f) = \{E \in Z(X) : S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} E\}.$$

PROOF: If $E \in \mathcal{Z}_A(f)$, then there exists $g \in A(X)$ such that $fg|_{X \setminus E} = 1$. Thus, $(fg)^*|_{\text{cl}_{\beta X}(X \setminus E)} \neq 0$. This means that $S_A(f) \cap \text{cl}_{\beta X}(X \setminus E) = \emptyset$ and thus $S_A(f) \subseteq \beta X \setminus \text{cl}_{\beta X}(X \setminus E) = \beta X \setminus \text{cl}_{\beta X}(\beta X \setminus \text{cl}_{\beta X} E) \subseteq \text{cl}_{\beta X} E$; i.e., $S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} E$. For the reverse inclusion, let $E \in Z(X)$ be such that $S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} E$. Thus, $S_A(f) \cap (\beta X \setminus \text{int}_{\beta X} \text{cl}_{\beta X} E) = \emptyset$. Hence, there exists some $h \in C^*(X)$ such that $h^\beta(S_A(f)) = \{1\}$ and $(\beta X \setminus \text{int}_{\beta X} \text{cl}_{\beta X} E) \subseteq Z(h^\beta)$. These imply that $S_A(f) \cap S_A(h) = \emptyset$ and $X \setminus E \subseteq X \setminus \text{int}_X E \subseteq Z(h)$. Therefore, $S_A(f^2 + h^2) = S_A(f) \cap S_A(h) = \emptyset$ which implies that there exists some $k \in A(X)$ such that $(f^2 + h^2)k = 1$. Thus, clearly $fk \in A(X)$ and $(f^2k)|_{X \setminus E} = ((f^2 + h^2)k)|_{X \setminus E} = 1$; i.e., $E \in \mathcal{Z}_A(f)$. \square

The following properties of the mapping \mathcal{Z}_A follow from Theorem 2.1. Note that for each two z -filters \mathcal{F} and \mathcal{F}' , we denote by $\mathcal{F} \wedge \mathcal{F}'$ and $\mathcal{F} \vee \mathcal{F}'$ the meet and join on the lattice of z -filters, respectively.

Proposition 2.2. *The following statements hold for each two elements f, g of an intermediate ring $A(X)$ and each $n \in \mathbb{N}$.*

- (a) *If $0 \leq f \leq g$, then $\mathcal{Z}_A(f) \subseteq \mathcal{Z}_A(g)$.*
- (b) *$\bigcap \mathcal{Z}_A(f) = Z(f)$.*

- (c) $\mathcal{Z}_A(fg) = \mathcal{Z}_A(f) \wedge \mathcal{Z}_A(g)$.
- (d) $\mathcal{Z}_A(f + g) \subseteq \mathcal{Z}_A(f) \vee \mathcal{Z}_A(g)$.
- (e) $\mathcal{Z}_A(f^2 + g^2) = \mathcal{Z}_A(f) \vee \mathcal{Z}_A(g)$.
- (f) $\mathcal{Z}_A(f^n) = \mathcal{Z}_A(f)$.

PROOF: Parts (c) through (f) easily follow from Theorem 2.1.

(a) It suffices to show that $S_A(g) \subseteq S_A(f)$. Let $p \notin S_A(f)$. Thus, $f \notin M_A^p$ and hence $g \notin M_A^p$, i.e., $p \notin S_A(g)$, since $0 \leq f \leq g$ and from [3, Theorem 2.5] it follows that every maximal ideal in $A(X)$ is a convex ideal. Therefore, $S_A(g) \subseteq S_A(f)$.

(b) It is evident that $Z(f) \subseteq E$ for each $E \in \mathcal{Z}_A(f)$. Thus, $Z(f) \subseteq \bigcap \mathcal{Z}_A(f)$. Let $x \in X$ and $x \notin Z(f)$. Hence, $x \notin S_A(f)$ and thus there exists a zero-set E such that $x \notin \text{cl}_{\beta X} E$ and $S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} E$. It follows that $E \in \mathcal{Z}_A(f)$, however, $x \notin E$. This implies that $\bigcap \mathcal{Z}_A(f) \subseteq Z(f)$ and the equality follows. \square

It is shown in [8, Theorem 1.2] that the mapping \mathcal{Z}_A could characterize $C^*(X)$ among intermediate rings. The following statement shows that the mapping \mathcal{Z}_A could also characterize $C(X)$ among intermediate rings in the case that X is a P -space.

Theorem 2.3. *Let $A(X)$ be an intermediate ring of $C(X)$. Then $\mathcal{Z}_A(f) = (Z(f))$ for each $f \in A(X)$ if and only if X is a P -space and $A(X) = C(X)$.*

PROOF: (\Rightarrow) As $\mathcal{Z}_A(f) = (Z(f))$ for each $f \in A(X)$, we have $Z(f) \in \mathcal{Z}_A(f)$ and thus $S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} Z(f)$. This implies that $S_A(f) = \text{cl}_{\beta X} Z(f) = \text{int}_{\beta X} \text{cl}_{\beta X} Z(f)$ for each $f \in A(X)$. Hence, X is a P -space and, by [10, Theorem 2.3], $A(X) = C(X)$

(\Leftarrow) It is clear that $\mathcal{Z}_C(f) \subseteq (Z(f))$. If $E \in (Z(f))$, then, $Z(f) \subseteq E$ and as X is a P -space, we have $S_C(f) = \text{cl}_{\beta X} Z(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X} E$. Thus, $(Z(f)) \subseteq \mathcal{Z}_C(f)$. \square

For an ideal I of an intermediate ring $A(X)$, we denote by $\mathcal{Z}_A(I)$ the set $\bigcup_{f \in I} \mathcal{Z}_A(f)$. Also, for a z -filter \mathcal{F} on X we denote by $\mathcal{Z}_A^{-1}(\mathcal{F})$ the set $\{f \in A(X) : \mathcal{Z}_A(f) \subseteq \mathcal{F}\}$. It is clear that $I \subseteq \mathcal{Z}_A^{-1} \mathcal{Z}_A(I)$ and $\mathcal{Z}_A \mathcal{Z}_A^{-1}(\mathcal{F}) \subseteq \mathcal{F}$. We call an ideal I in $A(X)$ a \mathcal{Z}_A -ideal, if $\mathcal{Z}_A^{-1} \mathcal{Z}_A(I) = I$. Also, a z -filter \mathcal{F} on X is called a \mathcal{Z}_A -filter, if $\mathcal{Z}_A \mathcal{Z}_A^{-1}(\mathcal{F}) = \mathcal{F}$. Evidently, $\mathcal{Z}_A^{-1} \mathcal{Z}_A(M_A^p) = M_A^p$. Hence, every maximal ideal in $A(X)$ is \mathcal{Z}_A -ideal.

Remark 2.4. It easily follows from Theorem 2.1 that $\mathcal{Z}_A(M_A^p) = \mathcal{Z}_A(O_A^p)$ for each $p \in \beta X$. Moreover, $\mathcal{Z}_A^{-1}(\mathcal{U}^p) = M_A^p$ for each $p \in \beta X$. Also, one can easily prove that $\mathcal{Z}_A(M)$ is contained in a unique z -ultrafilter for each maximal ideal M in $A(X)$. These provide a new approach to [3, Theorem 3.2].

It easily follows from Theorem 2.1 that every \mathcal{Z}_A -ideal of an intermediate ring $A(X)$ is a z -ideal. However, the converse of this fact does not hold, in general. For example, let $X = \mathbb{R}$ and $A(X) = C^*(X) = C^*(\mathbb{R})$ and $p \in X$; then $O_A^p \subsetneq M_A^p$, easily verifiable. It follows that O_A^p is not a \mathcal{Z}_A -ideal in $A(X)$, however, it is clearly a z -ideal. In [7, Theorem 2.14] it is stated that whenever every ideal of

an intermediate ring $A(X)$ is a \mathcal{Z}_A -ideal, then X is a P -space. The next theorem shows that even when every z -ideal is a \mathcal{Z}_A -ideal, then we have $A(X) = C(X)$. Note that we call an ideal I in $A(X)$ a z_A -ideal, if whenever $Z(f) \subseteq Z(g)$ where $f \in I$ and $g \in A(X)$, then $g \in I$. It is easy to see that the ideals O_A^p and $M^p \cap A(X)$, for each $p \in \beta X$, are z_A -ideals in $A(X)$. Also, the ideal M_A^p , for each $p \in \beta X \setminus v_A X$, is a maximal ideal in $A(X)$ which is not a z_A -ideal, refer to [2], [1], and [10] for more details.

Theorem 2.5. *The following statements are equivalent for an intermediate ring $A(X)$.*

- (a) Every z -ideal in $A(X)$ is a \mathcal{Z}_A -ideal.
- (b) Every z_A -ideal in $A(X)$ is a \mathcal{Z}_A -ideal.
- (c) X is a P -space and $A(X) = C(X)$.

PROOF: (a) \Rightarrow (c) The proof is straightforward by using [7, Theorem 2.5 and Theorem 3.10].

(c) \Rightarrow (a) By our hypothesis and Theorem 2.3, $\mathcal{Z}_C(f) = (Z(f))$ for each $f \in C(X)$. This clearly implies that every z -ideal in $C(X)$ is a \mathcal{Z}_C -ideal.

(b) \Rightarrow (c) As O_A^p , for each $p \in \beta X$, is a z_A -ideal, by our hypothesis, O_A^p would be a \mathcal{Z}_A -ideal. Hence, $O_A^p = \mathcal{Z}_A^{-1} \mathcal{Z}_A(O_A^p) = \mathcal{Z}_A^{-1} \mathcal{Z}_A(M_A^p) = M_A^p$. This means that $A(X)$ is a regular ring. Thus, by [7, Theorem 2.5 and Theorem 3.3], X is a P -space and $A(X) = C(X)$.

(c) \Rightarrow (b) It is evident that whenever $A(X) = C(X)$, then z_A -ideals coincide with z -ideals in $A(X)$. Moreover, as X is a P -space, by [4, Theorem 3.13], every z_A -ideal is a \mathcal{Z}_A -ideal. □

3. The mapping \mathfrak{S}_A

The mapping \mathfrak{S}_A on an intermediate ring $A(X)$ was first introduced in [8] as follows: for each $f \in A(X)$

$$\mathfrak{S}_A(f) = \{E \in Z(X) : \text{for all zero-sets } H \subseteq X \setminus E, \exists g \in A(X), fg|_H = 1\}.$$

The following statement provides a new topological description to the mapping \mathfrak{S}_A which could be proved similar to Theorem 2.1.

Theorem 3.1. *Let $A(X)$ be an intermediate ring of $C(X)$. For each $f \in A(X)$ we have*

$$\mathfrak{S}_A(f) = \{E \in Z(X) : \forall H \in Z(X) \text{ and } H \subseteq X \setminus E, S_A(f) \subseteq \text{int}_{\beta X} \text{cl}_{\beta X}(X \setminus H)\}.$$

The following question concerning the basic properties of the mapping \mathfrak{S}_A has been raised in [13].

Question 1. Let $f, g \in A(X)$. Which properties analogous to those of Proposition 2.2 hold with \mathfrak{S}_A in place of \mathcal{Z}_A ?

The answer to Question 1 will be given by the next statement which is an easy consequence of Theorem 3.1.

Proposition 3.2. *For each two elements f, g of an intermediate ring $A(X)$, the following statements hold.*

- (a) *If $0 \leq f \leq g$, then $\mathfrak{S}_A(f) \subseteq \mathfrak{S}_A(g)$.*
- (b) *$\bigcap \mathfrak{S}_A(f) = Z(f)$.*
- (c) *$\mathfrak{S}_A(fg) = \mathfrak{S}_A(f) \wedge \mathfrak{S}_A(g)$.*
- (d) *$\mathfrak{S}_A(f + g) \subseteq \mathfrak{S}_A(f) \vee \mathfrak{S}_A(g)$.*
- (e) *$\mathfrak{S}_A(f^2 + g^2) = \mathfrak{S}_A(f) \vee \mathfrak{S}_A(g)$.*
- (f) *$\mathfrak{S}_A(f^n) = \mathfrak{S}_A(f)$ for each $n \in \mathbb{N}$.*

It is stated in [3, Lemma 1.3] that for a z -filter \mathcal{F} and $f \in A(X)$, we have $\lim_{\mathcal{F}} fg = 0$ for each $g \in A(X)$ if and only if $\mathcal{F} \subseteq \mathcal{Z}_A(f)$. Note that for a z -filter \mathcal{F} on X and $f \in C(X)$, we use $\lim_{\mathcal{F}} f$ to denote the limit of the filter base $f(\mathcal{F})$. The following question has been raised in [13].

Question 2. Let $f \in A(X)$ and \mathcal{F} be a z -filter on X . Is it the case that $\mathfrak{S}_A(f) \subseteq \mathcal{F}$ if and only if $\lim_{\mathcal{F}} fg = 0$ for every $g \in A(X)$?

ANSWER TO QUESTION 2: As stated in [13], if $\mathfrak{S}_A(f) \subseteq \mathcal{F}$, then $\lim_{\mathcal{F}} fg = 0$ for every $g \in A(X)$. We show that the converse of this statement does not hold, in general. Take $X = \mathbb{R}$, $f(x) = x$ for each $x \in \mathbb{R}$ and \mathcal{F} be the z -filter of all zero set neighbourhood of 0 in \mathbb{R} . Then for all $h \in C(\mathbb{R})$, $\lim_{\mathcal{F}} fh = 0$ but $\mathfrak{S}_C(f) = \{Z \in Z[X] : 0 \in Z\}$ which contains \mathcal{F} properly.

In [7, Theorem 2.8] it is stated that a topological space X is a P -space if and only if, for every intermediate ring $A(X)$, we have $\mathfrak{S}_A(M_A^p) = \mathfrak{S}_A(O_A^p)$ for each $p \in X$. The next theorem extends this fact for each $p \in \beta X$. We use the following lemma which could be proved by a little modification of the arguments of [7, Proposition 2.7] and exploiting the complete regularity of βX .

Lemma 3.3. *Let $A(X)$ be an intermediate ring of $C(X)$. Then $\mathfrak{S}_A(O_A^p) = \mathcal{Z}_A(M_A^p)$ for each $p \in \beta X$.*

Theorem 3.4. *A topological space X is a P -space if and only if for every intermediate ring $A(X)$ we have $\mathfrak{S}_A(M_A^p) = \mathfrak{S}_A(O_A^p)$ for each $p \in \beta X$.*

PROOF: (\Rightarrow) This easily follows from [7, Theorem 2.8].

(\Leftarrow) By our hypothesis, $\mathfrak{S}_C(M^p) = \mathfrak{S}_C(O^p)$. Hence, $Z(M_A^p) = \mathfrak{S}_C(O^p) = Z(O^p)$ for each $p \in \beta X$ which clearly implies that X is a P -space. \square

For an ideal I of an intermediate ring $A(X)$, we denote by $\mathfrak{S}_A(I)$ the set $\bigcup_{f \in I} \mathfrak{S}_A(f)$. Moreover, for a z -filter \mathcal{F} on X , we use $\mathfrak{S}_A^{-1}(\mathcal{F})$ to denote the set $\{f \in A(X) : \mathfrak{S}_A(f) \subseteq \mathcal{F}\}$. An ideal I in $A(X)$ is called a \mathfrak{S}_A -ideal, if $\mathfrak{S}_A^{-1}\mathfrak{S}_A(I) = I$. Also, a z -filter \mathcal{F} on X is called a \mathfrak{S}_A -filter, if $\mathfrak{S}_A\mathfrak{S}_A^{-1}(\mathcal{F}) = \mathcal{F}$. It easily follows from Theorem 2.1 and Theorem 3.1 that every \mathcal{Z}_A -ideal in $A(X)$ is a \mathfrak{S}_A -ideal and every \mathfrak{S}_A -ideal in $A(X)$ is a z -ideal. However, the converse of these facts does not hold, in general, see the next example.

Example 3.5. (a) Whenever X is not a P -space, then there exists $p \in \beta X$ such that $O^p \neq M^p$. Hence, O^p is clearly a z -ideal and thus is a \mathfrak{S}_C -ideal in $C(X)$, however, it is not a \mathcal{Z}_C -ideal, since, $\mathcal{Z}_C^{-1}\mathcal{Z}_C(O^p) = M^p$.

(b) Let $X = \mathbb{N}$ and $A(X) = C^*(\mathbb{N})$. It is clear that there exists $p \in \beta\mathbb{N}$ such that $M_A^p \neq O_A^p$. By Theorem 3.4, $\mathfrak{S}_A(M_A^p) = \mathfrak{S}_A(O_A^p)$ for each $p \in \beta\mathbb{N}$. Thus, $\mathfrak{S}_A^{-1}\mathfrak{S}_A(O_A^p) = \mathfrak{S}_A^{-1}\mathfrak{S}_A(M_A^p) = M_A^p \neq O_A^p$ which means that O_A^p is not a \mathfrak{S}_A -ideal, however, it is evidently a z -ideal.

The following theorem shows that coincidence of z_A -ideals with \mathfrak{S}_A -ideals characterizes $C(X)$ among intermediate rings.

Theorem 3.6. *Let $A(X)$ be an intermediate ring of $C(X)$. Then every \mathfrak{S}_A -ideal in $A(X)$ is a z_A -ideal if and only if $A(X) = C(X)$.*

PROOF: (\Rightarrow) Since $A(X) \neq C(X)$, there exists $f \in C(X) \setminus A(X)$. Take $g = 1/(1 + |f|)$, then g is a non-unit of $A(X)$. Consequently, $g \in M_A^p$ for some $p \in \beta X$. It is clear that, since $Z(g) = \emptyset$, M_A^p is not a z_A -ideal in $A(X)$, but M_A^p being a maximal ideal is a \mathfrak{S}_A -ideal in $A(X)$.

(\Leftarrow) This is evident, since, \mathfrak{S}_C -ideals of $C(X)$ are nothing other than z -ideals. □

Remark 3.7. It is stated in [7, Theorem 2.14] that whenever every ideal of an intermediate ring $A(X)$ is a \mathfrak{S}_A -ideal, then X is a P -space. We show that this condition implies $A(X) = C(X)$. It is evident that whenever every ideal in $A(X)$ is a \mathfrak{S}_A -ideal, then every ideal in $A(X)$ is a z -ideal and thus, by [6, Theorem 1.2], $A(X)$ is a regular ring. Hence, we have $A(X) = C(X)$, since, otherwise, there exists $f \in A(X)$ such that $Z(f) = \emptyset$ and $S_A(f) \neq \emptyset$. Hence, if we choose some $p \in S_A(f)$, then $M^p \cap A(X) \neq M_A^p$ which means that $M^p \cap A(X)$ is a prime ideal in $A(X)$ which is not a maximal ideal and thus $A(X)$ is not a regular ring. Other than this, in the proof of this theorem in [7] it is stated that $\mathfrak{S}_A^{-1}\mathcal{Z}_A(M_A^p) = M_A^p$ for each $p \in X$. We claim that this equality does not hold, in general. For example, if we let $A(X) = C(X)$ where X is not a P -space, then there exists $f \in C(X)$ and $p \in X$ such that $f \in M^p \setminus O^p$. Also, we clearly have $\mathfrak{S}_C^{-1}(\mathcal{F}) = Z^{-1}(\mathcal{F})$ for each z -filter \mathcal{F} . It follows that $\mathfrak{S}_C^{-1}\mathcal{Z}_C(M^p) = Z^{-1}\mathcal{Z}_C(M^p) = Z^{-1}Z(O^p) = O^p \neq M^p$. Recall that $\mathcal{Z}_A(M_A^p) = \mathcal{Z}_A(O_A^p)$ and $\mathcal{Z}_A(M_A^p) = \{E \in Z(X) : p \in \text{int}_{\beta X} \text{cl}_{\beta X} E\}$. Hence, $\mathcal{Z}_C(M^p) = Z(O^p)$ for each $p \in \beta X$.

The following question has been raised in [13].

Question 3. Is it the case that if \mathcal{F} is a z -filter on X , then $\mathfrak{S}_A^{-1}(\mathcal{F})$ is an ideal in $A(X)$?

We answer Question 3 by using Proposition 3.2.

ANSWER TO QUESTION 3: Let \mathcal{F} be a z -filter on X , $f, g \in \mathfrak{S}_A^{-1}(\mathcal{F})$ and $h \in A(X)$. Then, $\mathfrak{S}_A(fh) = \mathfrak{S}_A(f) \wedge \mathfrak{S}_A(h) \subseteq \mathcal{F}$, since, $f \in \mathfrak{S}_A^{-1}(\mathcal{F})$. Hence, $fh \in \mathfrak{S}_A^{-1}(\mathcal{F})$. Moreover, $\mathfrak{S}_A(f + g) \subseteq \mathfrak{S}_A(f) \vee \mathfrak{S}_A(g) \subseteq \mathcal{F}$, since, $f, g \in \mathfrak{S}_A^{-1}(\mathcal{F})$. Thus, $f + g \in \mathfrak{S}_A^{-1}(\mathcal{F})$. Therefore, $\mathfrak{S}_A^{-1}(\mathcal{F})$ is an ideal in $A(X)$ for each z -filter \mathcal{F} on X .

The following question also has been raised in [13].

Question 4: Is it the case that $A(X) = C(X)$ if and only if $\mathfrak{S}_A(M)$ is a z -ultrafilter for every maximal ideal M ?

ANSWER TO QUESTION 4. If $A(X) = C(X)$, then for each maximal ideal M of $A(X)$, $\mathfrak{S}_A(M) = Z(M)$ is a z -ultrafilter on X . On the other hand, it is easily inferred from [7, Theorem 2.10] that if X is a non pseudocompact P -space, then there exists an intermediate ring $B(X) \subsetneq C(X)$ for which for any maximal ideal M of $B(X)$, $\mathfrak{S}_B(M)$ is a z -ultrafilter on X .

The following question has been raised in [13].

Question 5. Is it the case that $A(X) = C(X)$ if and only if every z -filter on X is a \mathfrak{S}_A -filter?

The answer to Question 5 is negative as it could be inferred from [7, Example 2.13].

Acknowledgment. The author would like to express his deep gratitude to the well-informed referee for giving useful suggestions on this article.

REFERENCES

- [1] Aliabad A. R., Parsinia M., *Remarks on subrings of $C(X)$ of the form $I + C^*(X)$* , Quaest. Math. **40** (2017), no. 1, 63–73.
- [2] Aliabad A. R., Parsinia M., *z_R -ideals and z_R° -ideals in subrings of \mathbb{R}^X* , to appear in Iranian J. Math. Sci. Inform.
- [3] Byun H. L., Watson S., *Prime and maximal ideals in subrings of $C(X)$* , Topology Appl. **40** (1991), no. 1, 45–62.
- [4] Domínguez J. M., Gómez Pérez J., *Intersections of maximal ideals in algebras between $C^*(X)$ and $C(X)$* , Iberoamerican Conf. on Topology and Its Applications, Morelia, 1997, Topology Appl. **98** (1999), no. 1–3, 149–165.
- [5] Gillman L., Jerison M., *Rings of Continuous Functions*, The University Series in Higher Mathematics, D. Van Nostrand Co., Princeton, New York, 1960.
- [6] Mason G., *z -ideals and prime ideals*, J. Algebra **26** (1973), 280–297.
- [7] Murray W., Sack J., Watson S., *P -spaces and intermediate rings of continuous functions*, Rocky Mountain J. Math. **47** (2017), no. 8, 2757–2775.
- [8] Panman P., Sack J., Watson S., *Correspondence between ideals and z -filters for rings of continuous functions between C^* and C* , Comment. Math. **52** (2012), no. 1, 11–20.
- [9] Parsinia M., *Remarks on LBI-subalgebras of $C(X)$* , Comment. Math. Univ. Carolin. **57** (2016), no. 2, 261–270.
- [10] Parsinia M., *Remarks on intermediate C -rings of $C(X)$* , Quaest. Math. (online 2017), 8 pages.
- [11] Plank D., *On a class of subalgebras of $C(X)$ with applications to $\beta X \setminus X$* , Fund. Math. **64** (1969), 41–54.
- [12] Redlin L., Watson S., *Maximal ideals in subalgebras of $C(X)$* , Proc. Amer. Math. Soc. **100** (1987), no. 4, 763–766.

- [13] Sack J., Watson S., *C and C^* among intermediate rings*, Topology Proc. **43** (2014), 69–82.
- [14] Sack J., Watson S., *Characterizing $C(X)$ among intermediate C -rings on X* , Topology Proc. **45** (2015), 301–313.

M. Parsinia:

DEPARTMENT OF MATHEMATICS, SHAHID CHAMRAN UNIVERSITY OF AHVAZ,
GOLESTAN BLVD 22, AHVAZ 6135743337, KHUZESTAN, IRAN

E-mail: parsiniamehdi@gmail.com

(Received November 30, 2017, revised May 1, 2018)