

Brahim Alioune; Mohamed Boucetta; Ahmed Sid'Ahmed Lessiad
On Riemann-Poisson Lie groups

Archivum Mathematicum, Vol. 56 (2020), No. 4, 225–247

Persistent URL: <http://dml.cz/dmlcz/148390>

Terms of use:

© Masaryk University, 2020

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON RIEMANN-POISSON LIE GROUPS

BRAHIM ALIOUNE^a, MOHAMED BOUCETTA^b, AND AHMED SID'AHMED LESSIAD^c

ABSTRACT. A Riemann-Poisson Lie group is a Lie group endowed with a left invariant Riemannian metric and a left invariant Poisson tensor which are compatible in the sense introduced in [4]. We study these Lie groups and we give a characterization of their Lie algebras. We give also a way of building these Lie algebras and we give the list of such Lie algebras up to dimension 5.

1. INTRODUCTION

In this paper, we study Lie groups endowed with a left invariant Riemannian metric and a left invariant Poisson tensor satisfying a compatibility condition to be defined below. They constitute a subclass of the class of *Riemann-Poisson manifolds* introduced and studied by the second author (see [2, 3, 4, 5]).

Let $(M, \pi, \langle \cdot, \cdot \rangle)$ be smooth manifold endowed with a Poisson tensor π and a Riemannian metric $\langle \cdot, \cdot \rangle$. We denote by $\langle \cdot, \cdot \rangle^*$ the Euclidean product on T^*M naturally associated to $\langle \cdot, \cdot \rangle$. The Poisson tensor defines a Lie algebroid structure on T^*M where the anchor map is the contraction $\#_\pi: T^*M \rightarrow TM$ given by $\langle \beta, \#_\pi(\alpha) \rangle = \pi(\alpha, \beta)$ and the Lie bracket on $\Omega^1(M)$ is the Koszul bracket given by

$$(1) \quad [\alpha, \beta]_\pi = \mathcal{L}_{\#_\pi(\alpha)}\beta - \mathcal{L}_{\#_\pi(\beta)}\alpha - d\pi(\alpha, \beta), \quad \alpha, \beta \in \Omega^1(M).$$

This Lie algebroid structure and the metric $\langle \cdot, \cdot \rangle^*$ define a contravariant connection $\mathcal{D}: \Omega^1(M) \times \Omega^1(M) \rightarrow \Omega^1(M)$ by Koszul formula

$$(2) \quad \begin{aligned} 2\langle \mathcal{D}_\alpha\beta, \gamma \rangle^* &= \#_\pi(\alpha) \cdot \langle \beta, \gamma \rangle^* + \#_\pi(\beta) \cdot \langle \alpha, \gamma \rangle^* - \#_\pi(\gamma) \cdot \langle \alpha, \beta \rangle^* \\ &+ \langle [\alpha, \beta]_\pi, \gamma \rangle^* + \langle [\gamma, \alpha]_\pi, \beta \rangle^* + \langle [\gamma, \beta]_\pi, \alpha \rangle^*, \quad \alpha, \beta, \gamma \in \Omega^1(M). \end{aligned}$$

This is the unique torsionless contravariant connection which is metric, i.e., for any $\alpha, \beta, \gamma \in \Omega^1(M)$,

$$\mathcal{D}_\alpha\beta - \mathcal{D}_\beta\alpha = [\alpha, \beta]_\pi \quad \text{and} \quad \#_\pi(\alpha) \cdot \langle \beta, \gamma \rangle^* = \langle \mathcal{D}_\alpha\beta, \gamma \rangle^* + \langle \beta, \mathcal{D}_\alpha\gamma \rangle^*.$$

The notion of contravariant connection was introduced by Vaisman in [13] and studied in more details by Fernandes in the context of Lie algebroids [8]. The

2020 *Mathematics Subject Classification*: primary 53A15; secondary 53D17, 22E05.

Key words and phrases: Lie group, Poisson manifolds, Riemannian metric.

Received August 22, 2019. Editor J. Slovák.

DOI: 10.5817/AM2020-4-225

connection \mathcal{D} defined above is called *contravariant Levi-Civita connection* associated to the couple $(\pi, \langle \cdot, \cdot \rangle)$ and it appeared first in [2].

The triple $(M, \pi, \langle \cdot, \cdot \rangle)$ is called a *Riemannian-Poisson manifold* if $\mathcal{D}\pi = 0$, i.e., for any $\alpha, \beta, \gamma \in \Omega^1(M)$,

$$(3) \quad \mathcal{D}\pi(\alpha, \beta, \gamma) := \#_{\pi}(\alpha) \cdot \pi(\beta, \gamma) - \pi(\mathcal{D}_{\alpha}\beta, \gamma) + \pi(\beta, \mathcal{D}_{\alpha}\gamma) = 0.$$

This notion was introduced by the second author in [2]. Riemann-Poisson manifolds turned out to have interesting geometric properties (see[2, 3, 4, 5]). Let's mention some of them.

- (1) The condition of compatibility (3) is weaker than the condition $\nabla\pi = 0$ where ∇ is the Levi-Civita connection of $\langle \cdot, \cdot \rangle$. Indeed, the condition (3) allows the Poisson tensor to have a variable rank. For instance, linear Poisson structures which are Riemann-Poisson exist and were characterized in [5]. Furthermore, let $(M, \langle \cdot, \cdot \rangle)$ be a Riemannian manifold and (X_1, \dots, X_r) a family of commuting Killing vector fields. Put

$$\pi = \sum_{i,j} X_i \wedge X_j.$$

Then $(M, \pi, \langle \cdot, \cdot \rangle)$ is a Riemann-Poisson manifold. This example illustrates also the weakness of the condition (3) and, more importantly, it is the local model of the geometry of noncommutative deformations studied by Hawkins (see [10, Theorem 6.6]).

- (2) Riemann-Poisson manifolds can be thought of as a generalization of Kähler manifolds. Indeed, let $(M, \pi, \langle \cdot, \cdot \rangle)$ be a Poisson manifold endowed with a Riemannian metric such that π is invertible. Denote by ω the symplectic form inverse of π . Then $(M, \pi, \langle \cdot, \cdot \rangle)$ is Riemann-Poisson manifold if and only if $\nabla\omega = 0$ where ∇ is the Levi-Civita connection of $\langle \cdot, \cdot \rangle$. In this case, if we define $A: TM \rightarrow TM$ by $\omega(u, v) = \langle Au, v \rangle$ then $-A^2$ is symmetric definite positive and hence there exists a unique $Q: TM \rightarrow TM$ symmetric definite positive such that $Q^2 = -A^2$. It follows that $J = AQ^{-1}$ satisfies $J^2 = -\text{Id}_{TM}$, skew-symmetric with respect $\langle \cdot, \cdot \rangle$ and $\nabla J = 0$. Hence $(M, J, \langle \cdot, \cdot \rangle)$ is a Kähler manifold and its Kähler form $\omega_J(u, v) = \langle Ju, v \rangle$ is related to ω by the following formula:

$$(4) \quad \omega(u, v) = -\omega_J\left(\sqrt{-A^2}u, v\right), \quad u, v \in TM.$$

Having this construction in mind, we will call in this paper a Kähler manifold a triple $(M, \langle \cdot, \cdot \rangle, \omega)$ where $\langle \cdot, \cdot \rangle$ is a Riemannian metric and ω is a nondegenerate 2-form ω such that $\nabla\omega = 0$ where ∇ is the Levi-Civita connection of $\langle \cdot, \cdot \rangle$.

- (3) The symplectic foliation of a Riemann-Poisson manifold when π has a constant rank has an important property namely it is both a Riemannian foliation and a Kähler foliation.

Recall that a Riemannian foliation is a foliated manifold (M, \mathcal{F}) with a Riemannian metric $\langle \cdot, \cdot \rangle$ such that the orthogonal distribution $T^{\perp}\mathcal{F}$ is totally geodesic.

Kähler foliations are a generalization of Kähler manifolds (see [6]) and, as for the notion of Kähler manifold, we call in this paper a Kähler foliation a foliated manifold (M, \mathcal{F}) endowed with a leafwise metric $\langle \cdot, \cdot \rangle_{\mathcal{F}} \in \Gamma(\otimes^2 T^* \mathcal{F})$ and a nondegenerate leafwise differential 2-form $\omega_{\mathcal{F}} \in \Gamma(\otimes^2 T^* \mathcal{F})$ such any leaf with the restrictions of $\langle \cdot, \cdot \rangle_{\mathcal{F}}$ and $\omega_{\mathcal{F}}$ is a Kähler manifold.

Theorem 1.1 ([4]). *Let $(M, \langle \cdot, \cdot \rangle, \pi)$ be a Riemann-Poisson manifold with π of constant rank. Then its symplectic foliation is both a Riemannian and a Kähler foliation.*

Having in mind these properties particularly Theorem 1.1, it will be interesting to find large classes of examples of Riemann-Poisson manifolds. This paper will describe the rich collection of examples which are obtained by providing an arbitrary Lie group G with a Riemannian metric $\langle \cdot, \cdot \rangle$ and a Poisson tensor π invariant under left translations and such that $(G, \langle \cdot, \cdot \rangle, \pi)$ is Riemann-Poisson. We call $(G, \langle \cdot, \cdot \rangle, \pi)$ a *Riemann-Poisson Lie group*. This class of examples can be enlarged substantially, with no extra work, as follows. If $(G, \langle \cdot, \cdot \rangle, \pi)$ is a Riemann-Poisson Lie group and Γ is any discrete subgroup of G then $\Gamma \backslash G$ carries naturally a structure of Riemann-Poisson manifold.

The paper is organized as follows. In Section 2, we give the material needed in the paper and we describe the infinitesimal counterpart of Riemann-Poisson Lie groups, namely, Riemann-Poisson Lie algebras. In Section 3, we prove our main result which gives an useful description of Riemann-Poisson Lie algebras (see Theorem 3.1). We use this theorem in Section 4 to derive a method for building Riemann-Poisson Lie algebras. We explicit this method by giving the list of Riemann-Poisson Lie algebras up to dimension 5.

2. RIEMANN-POISSON LIE GROUPS AND THEIR INFINITESIMAL CHARACTERIZATION

Let G be a Lie group and $(\mathfrak{g} = T_e G, [\cdot, \cdot])$ its Lie algebra.

- (1) A left invariant Poisson tensor π on G is entirely determined by

$$\pi(\alpha, \beta)(a) = r(L_a^* \alpha, L_a^* \beta),$$

where $a \in G$, $\alpha, \beta \in T_a^* G$, L_a is the left multiplication by a and $r \in \wedge^2 \mathfrak{g}$ satisfies the classical Yang-Baxter equation

$$(5) \quad [r, r] = 0,$$

where $[r, r] \in \wedge^3 \mathfrak{g}$ is given by

$$(6) \quad [r, r](\alpha, \beta, \gamma) : = \langle \alpha, [r_{\#}(\beta), r_{\#}(\gamma)] \rangle + \langle \beta, [r_{\#}(\gamma), r_{\#}(\alpha)] \rangle + \langle \gamma, [r_{\#}(\alpha), r_{\#}(\beta)] \rangle, \quad \alpha, \beta, \gamma \in \mathfrak{g}^*,$$

and $r_{\#} : \mathfrak{g}^* \rightarrow \mathfrak{g}$ is the contraction associated to r . In this case, the Koszul bracket (1) when restricted to left invariant differential 1-forms induces a Lie bracket on \mathfrak{g}^* given by

$$(7) \quad [\alpha, \beta]_r = \text{ad}_{r_{\#}(\alpha)}^* \beta - \text{ad}_{r_{\#}(\beta)}^* \alpha, \quad \alpha, \beta \in \mathfrak{g}^*,$$

where $\prec \text{ad}_u^* \alpha, v \succ = - \prec \alpha, [u, v] \succ$. Moreover, $r_{\#}$ becomes a morphism of Lie algebras, i.e.,

$$(8) \quad r_{\#}([\alpha, \beta]_r) = [r_{\#}(\alpha), r_{\#}(\beta)], \quad \alpha, \beta \in \mathfrak{g}^*.$$

(2) A left invariant Riemannian metric $\langle \cdot, \cdot \rangle$ on G is entirely determined by

$$\langle u, v \rangle(a) = \rho(T_a L_{a^{-1}} u, T_a L_{a^{-1}} v),$$

where $a \in G$, $u, v \in T_a G$ and ρ is a scalar product on \mathfrak{g} . The Levi-Civita connection of $\langle \cdot, \cdot \rangle$ is left invariant and induces a product $A: \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ given by

$$(9) \quad 2\varrho(A_u v, w) = \varrho([u, v], w) + \varrho([w, u], v) + \varrho([w, v], u), \quad u, v, w \in \mathfrak{g}.$$

It is the unique product on \mathfrak{g} satisfying

$$A_u v - A_v u = [u, v] \quad \text{and} \quad \varrho(A_u v, w) + \varrho(v, A_u w) = 0,$$

for any $u, v, w \in \mathfrak{g}$. We call A the *Levi-Civita product* associated to $(\mathfrak{g}, [\cdot, \cdot], \rho)$.

(3) Let $(G, \langle \cdot, \cdot \rangle, \Omega)$ be a Lie group endowed with a left invariant Riemannian metric and a nondegenerate left invariant 2-form. Then $(G, \langle \cdot, \cdot \rangle, \Omega)$ is a Kähler manifold if and only if, for any $u, v, w \in \mathfrak{g}$,

$$(10) \quad \omega(A_u v, w) + \omega(u, A_u v) = 0,$$

where $\omega = \Omega(e)$, $\rho = \langle \cdot, \cdot \rangle(e)$ and A is the Levi-Civita product of $(\mathfrak{g}, [\cdot, \cdot], \rho)$. In this case we call $(\mathfrak{g}, [\cdot, \cdot], \rho, \omega)$ a Kähler Lie algebra.

As all the left invariant structures on Lie groups, Riemann-Poisson Lie groups can be characterized at the level of their Lie algebras.

Proposition 2.1. *Let $(G, \pi, \langle \cdot, \cdot \rangle)$ be a Lie group endowed with a left invariant bivector field and a left invariant metric and $(\mathfrak{g}, [\cdot, \cdot])$ its Lie algebra. Put $r = \pi(e) \in \wedge^2 \mathfrak{g}$, $\varrho = \langle \cdot, \cdot \rangle_e$ and ϱ^* the associated Euclidean product on \mathfrak{g}^* . Then $(G, \pi, \langle \cdot, \cdot \rangle)$ is a Riemann-Poisson Lie group if and only if*

$$(i) \quad [r, r] = 0,$$

$$(ii) \quad \text{for any } \alpha, \beta, \gamma \in \mathfrak{g}^*, \quad r(A_\alpha \beta, \gamma) + r(\beta, A_\alpha \gamma) = 0,$$

where A is the Levi-Civita product associated to $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$.

Proof. For any $u \in \mathfrak{g}$ and $\alpha \in \mathfrak{g}^*$, we denote by u^ℓ and α^ℓ , respectively, the left invariant vector field and the left invariant differential 1-form on G given by

$$u^\ell(a) = T_e L_a(u) \quad \text{and} \quad \alpha^\ell(a) = T_a^* L_{a^{-1}}(\alpha), \quad a \in G, \quad L_a(b) = ab.$$

Since π and $\langle \cdot, \cdot \rangle$ are left invariant, one can see easily from (1) and (2) that we have, for any $\alpha, \beta, \gamma \in \mathfrak{g}^*$,

$$\begin{cases} [\pi, \pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = [r, r](\alpha, \beta, \gamma), \quad \#_\pi(\alpha^\ell) = (r_{\#}(\alpha))^\ell, \quad \mathcal{L}_{\#_\pi(\alpha^\ell)} \beta^\ell = (\text{ad}_{r_{\#}(\alpha)}^* \beta)^\ell, \\ [\alpha^\ell, \beta^\ell]_\pi = ([\alpha, \beta]_r)^\ell, \quad \mathcal{D}_{\alpha^\ell} \beta^\ell = (A_\alpha \beta)^\ell. \end{cases}$$

The proposition follows from these formulas, (3) and the fact that $(G, \pi, \langle \cdot, \cdot \rangle)$ is a Riemann-Poisson Lie group if and only if, for any $\alpha, \beta, \gamma \in \mathfrak{g}^*$,

$$[\pi, \pi]_S(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0 \quad \text{and} \quad \mathcal{D}\pi(\alpha^\ell, \beta^\ell, \gamma^\ell) = 0. \quad \square$$

Conversely, given a triple $(\mathfrak{g}, r, \varrho)$ where \mathfrak{g} is a real Lie algebra, $r \in \wedge^2 \mathfrak{g}$ and ϱ a Euclidean product on \mathfrak{g} satisfying the conditions (i) and (ii) in Proposition 2.1 then, for any Lie group G whose Lie algebra is \mathfrak{g} , if π and $\langle \cdot, \cdot \rangle$ are the left invariant bivector field and the left invariant metric associated to (r, ϱ) then $(G, \pi, \langle \cdot, \cdot \rangle)$ is a Riemann-Poisson Lie group.

Definition 2.1. A *Riemann-Poisson Lie algebra* is a triple $(\mathfrak{g}, r, \varrho)$ where \mathfrak{g} is a real Lie algebra, $r \in \wedge^2 \mathfrak{g}$ and ϱ a Euclidean product on \mathfrak{g} satisfying the conditions (i) and (ii) in Proposition 2.1.

To end this section, we give another characterization of the solutions of the classical Yang-Baxter equation (5) which will be useful later.

We observe that $r \in \wedge^2 \mathfrak{g}$ is equivalent to the data of a vector subspace $S \subset \mathfrak{g}$ and a nondegenerate 2-form $\omega_r \in \wedge^2 S^*$.

Indeed, for $r \in \wedge^2 \mathfrak{g}$, we put $S = \text{Im}r_\#$ and $\omega_r(u, v) = r(r_\#^{-1}(u), r_\#^{-1}(v))$ where $u, v \in S$ and $r_\#^{-1}(u)$ is any antecedent of u by $r_\#$.

Conversely, let (S, ω) be a vector subspace of \mathfrak{g} with a non-degenerate 2-form. The 2-form ω defines an isomorphism $\omega^b: S \rightarrow S^*$ by $\omega^b(u) = \omega(u, \cdot)$, we denote by $\#: S^* \rightarrow S$ its inverse and we put $r_\# = \# \circ i^*$ where $i^*: \mathfrak{g}^* \rightarrow S^*$ is the dual of the inclusion $i: S \hookrightarrow \mathfrak{g}$.

With this observation in mind, the following proposition gives another description of the solutions of the Yang-Baxter equation.

Proposition 2.2. *Let $r \in \wedge^2 \mathfrak{g}$ and (S, ω_r) its associated vector subspace. The following assertions are equivalent:*

- (1) $[r, r] = 0$.
- (2) S is a subalgebra of \mathfrak{g} and

$$\delta\omega_r(u, v, w) := \omega_r(u, [v, w]) + \omega_r(v, [w, u]) + \omega_r(w, [u, v]) = 0$$

for any $u, v, w \in S$.

Proof. The proposition follows from the following formulas:

$$\langle \gamma, r_\#([\alpha, \beta]_r) - [r_\#(\alpha), r_\#(\beta)] \rangle = -[r, r](\alpha, \beta, \gamma), \quad \alpha, \beta, \gamma \in \mathfrak{g}^*$$

and, if S is a subalgebra,

$$[r, r](\alpha, \beta, \gamma) = -\delta\omega_r(r_\#(\alpha), r_\#(\beta), r_\#(\gamma)). \quad \square$$

This proposition shows that there is a correspondence between the set of solutions of the Yang-Baxter equation the set of symplectic subalgebras of \mathfrak{g} . We recall that a symplectic algebra is a Lie algebra S endowed with a non-degenerate 2-form ω such that $\delta\omega = 0$.

3. A CHARACTERIZATION OF RIEMANN-POISSON LIE ALGEBRAS

In this section, we combine Propositions 2.1 and 2.2 to establish a characterization of Riemann-Poisson Lie algebras which will be used later to build such Lie algebras. We establish first an intermediary result.

Proposition 3.1. *Let $(\mathfrak{g}, r, \varrho)$ be a Lie algebra endowed with $r \in \wedge^2 \mathfrak{g}$ and a Euclidean product ϱ . Denote by $\mathcal{I} = \ker r_{\#}$, \mathcal{I}^{\perp} its orthogonal with respect to ϱ^* and A the Levi-Civita product associated to $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$. Then $(\mathfrak{g}, r, \varrho)$ is a Riemann-Poisson Lie algebra if and only if:*

- (c₁) $[r, r] = 0$.
- (c₂) For all $\alpha \in \mathcal{I}, A_{\alpha} = 0$.
- (c₃) For all $\alpha, \beta, \gamma \in \mathcal{I}^{\perp}, A_{\alpha}\beta \in \mathcal{I}^{\perp}$ and

$$r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0.$$

Proof. By using the splitting $\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^{\perp}$, on can see that the conditions (i) and (ii) in Proposition 2.1 are equivalent to

$$(11) \quad \begin{cases} [r, r] = 0, \\ r(A_{\alpha}\beta, \gamma) = 0, \alpha \in \mathcal{I}, \beta \in \mathcal{I}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0, \alpha \in \mathcal{I}, \beta \in \mathcal{I}^{\perp}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) = 0, \alpha \in \mathcal{I}^{\perp}, \beta \in \mathcal{I}, \gamma \in \mathcal{I}^{\perp}, \\ r(A_{\alpha}\beta, \gamma) + r(\beta, A_{\alpha}\gamma) = 0, \alpha \in \mathcal{I}^{\perp}, \beta \in \mathcal{I}^{\perp}, \gamma \in \mathcal{I}^{\perp}. \end{cases}$$

Suppose that the conditions (c₁)-(c₃) hold. Then for any $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^{\perp}$, $A_{\beta}\alpha = [\beta, \alpha]_r$ and hence $r_{\#}(A_{\beta}\alpha) = [r_{\#}(\beta), r_{\#}(\alpha)] = 0$ and hence the equations in (11) holds.

Conversely, suppose that (11) holds. Then (c₁) holds obviously.

For any $\alpha, \beta \in \mathcal{I}$, the second equation in (11) is equivalent to $A_{\alpha}\beta \in \mathcal{I}$ and we have from (7) and (9) $[\alpha, \beta]_r = 0$ and $A_{\alpha}\beta \in \mathcal{I}^{\perp}$. Thus $A_{\alpha}\beta = 0$.

Take now $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^{\perp}$. For any $\gamma \in \mathcal{I}$, $\varrho^*(A_{\alpha}\beta, \gamma) = -\varrho^*(\beta, A_{\alpha}\gamma) = 0$ and hence $A_{\alpha}\beta \in \mathcal{I}^{\perp}$. On the other hand,

$$r_{\#}([\alpha, \beta]_r) = r_{\#}(A_{\alpha}\beta) - r_{\#}(A_{\beta}\alpha) \stackrel{(8)}{=} [r_{\#}(\alpha), r_{\#}(\beta)] = 0.$$

So, for any $\gamma \in \mathcal{I}^{\perp}$,

$$\prec \gamma, r_{\#}(A_{\alpha}\beta) \succ = \prec \gamma, r_{\#}(A_{\beta}\alpha) \succ = r(A_{\beta}\alpha, \gamma) \stackrel{(11)}{=} 0.$$

This shows that $A_{\alpha}\beta \in \mathcal{I}$ and hence $A_{\alpha}\beta = 0$. Finally, (c₂) is true. Now, for any $\alpha \in \mathcal{I}^{\perp}$, the fourth equation in (11) implies that A_{α} leaves invariant \mathcal{I} and since it is skew-symmetric it leaves invariant \mathcal{I}^{\perp} and (c₃) follows. This completes the proof. □

Proposition 3.2. *Let $(\mathfrak{g}, \varrho, r)$ be a Lie algebra endowed with a solution of classical Yang-Baxter equation and a bi-invariant Euclidean product, i.e.,*

$$\varrho(\text{ad}_u v, w) + \varrho(v, \text{ad}_u w) = 0, \quad u, v, w \in \mathfrak{g}.$$

Then $(\mathfrak{g}, \varrho, r)$ is Riemann-Poisson Lie algebra if and only if $\text{Im} r_{\#}$ is an abelian subalgebra.

Proof. Since ϱ is bi-invariant, one can see easily that for any $u \in \mathfrak{g}$, ad_u^* is skew-symmetric with respect to ϱ^* and hence the Levi-Civita product A associated to $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$ is given by $A_{\alpha}\beta = \text{ad}_{r_{\#}(\alpha)}^*\beta$. So, $(\mathfrak{g}, \varrho, r)$ is Riemann-Poisson Lie algebra if and only if, for any $\alpha, \beta, \gamma \in \mathfrak{g}^*$,

$$\begin{aligned} 0 &= r(\text{ad}_{r_{\#}(\alpha)}^*\beta, \gamma) + r(\beta, \text{ad}_{r_{\#}(\alpha)}^*\gamma) \\ &= \langle \beta, [r_{\#}(\alpha), r_{\#}(\gamma)] \rangle - \langle \gamma, [r_{\#}(\alpha), r_{\#}(\beta)] \rangle \\ &\stackrel{(5)}{=} \langle \alpha, [r_{\#}(\beta), r_{\#}(\gamma)] \rangle \end{aligned}$$

and the result follows. □

Let $(\mathfrak{g}, [\cdot, \cdot])$ be a Lie algebra, $r \in \wedge^2 \mathfrak{g}$ and ϱ a Euclidean product on \mathfrak{g} . Denote by (S, ω_r) the symplectic vector subspace associated to r and by $\#: \mathfrak{g}^* \rightarrow \mathfrak{g}$ the isomorphism given by ϱ . Note that the Euclidean product on \mathfrak{g}^* is given by $\varrho^*(\alpha, \beta) = \varrho(\#(\alpha), \#(\beta))$. We have

$$\mathfrak{g}^* = \mathcal{I} \oplus \mathcal{I}^\perp \quad \text{and} \quad \mathfrak{g} = S \oplus S^\perp,$$

where $\mathcal{I} = \ker r_{\#}$. Moreover, $r_{\#}: \mathcal{I}^\perp \rightarrow S$ is an isomorphism, we denote by $\tau: S \rightarrow \mathcal{I}^\perp$ its inverse. From the relation

$$\varrho(\#(\alpha), r_{\#}(\beta)) = \langle \alpha, r_{\#}(\beta) \rangle = r(\beta, \alpha),$$

we deduce that $\#: \mathcal{I} \rightarrow S^\perp$ is an isomorphism and hence $\#: \mathcal{I}^\perp \rightarrow S$ is also an isomorphism.

Consider the isomorphism $J: S \rightarrow S$ linking ω_r to $\varrho|_S$, i.e.,

$$\omega_r(u, v) = \rho(Ju, v), \quad u, v \in S.$$

One can see easily that $J = -\# \circ \tau$.

Theorem 3.1. *With the notations above, $(\mathfrak{g}, r, \varrho)$ is a Riemann-Poisson Lie algebra if and only if the following conditions hold:*

- (1) $(S, \varrho|_S, \omega_r)$ is a Kähler Lie subalgebra, i.e., for all $s_1, s_2, s_3 \in S$,
- (12)
$$\omega_r(\nabla_{s_1}s_2, s_3) + \omega_r(s_2, \nabla_{s_1}s_3) = 0,$$

where ∇ is the Levi-Civita product associated to $(S, [\cdot, \cdot], \varrho|_S)$.
- (2) for all $s \in S$ and all $u, v \in S^\perp$,
- (13)
$$\varrho(\phi_S(s)(u), v) + \varrho(u, \phi_S(s)(v)) = 0,$$

where $\phi_S: S \rightarrow \text{End}(S^\perp)$, $u \mapsto \text{pr}_{S^\perp} \circ \text{ad}_u$ and $\text{pr}_{S^\perp}: \mathfrak{g} \rightarrow S^\perp$ is the orthogonal projection.
- (3) For all $s_1, s_2 \in S$ and all $u \in S^\perp$,
- (14)
$$\omega_r(\phi_{S^\perp}(u)(s_1), s_2) + \omega_r(s_1, \phi_{S^\perp}(u)(s_2)) = 0,$$

where $\phi_{S^\perp}: S^\perp \rightarrow \text{End}(S)$, $u \mapsto \text{pr}_S \circ \text{ad}_u$ and $\text{pr}_S: \mathfrak{g} \rightarrow S$ is the orthogonal projection.

Proof. Suppose first that $(\mathfrak{g}, r, \varrho)$ is a Riemann-Poisson Lie algebra. According to Propositions 3.1 and 2.2, this is equivalent to

$$(15) \quad \begin{cases} (S, \omega_r) \text{ is a symplectic subalgebra,} \\ \forall \alpha \in \mathcal{I}, A_\alpha = 0, \\ \forall \alpha, \beta, \gamma \in \mathcal{I}^\perp, A_\alpha \beta \in \mathcal{I}^\perp \quad \text{and} \quad r(A_\alpha \beta, \gamma) + r(\beta, A_\alpha \gamma) = 0, \end{cases}$$

where A is the Levi-Civita product of $(\mathfrak{g}^*, [\cdot, \cdot]_r, \varrho^*)$.

For $\alpha, \beta \in \mathcal{I}$ and $\gamma \in \mathcal{I}^\perp$,

$$(16) \quad \begin{aligned} 2\varrho^*(A_\alpha \beta, \gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= \varrho^*(\text{ad}_{r_\#(\gamma)}^* \beta, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \alpha, \beta) \\ &= -\prec \beta, [r_\#(\gamma), \#(\alpha)] \succ -\prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) - \varrho(\#(\alpha), [r_\#(\gamma), \#(\beta)]). \end{aligned}$$

Since $\#: \mathcal{I} \rightarrow S^\perp$ and $r_\#: \mathcal{I}^\perp \rightarrow S$ are isomorphisms, we deduce from (16) that $A_\alpha \beta = 0$ for any $\alpha, \beta \in \mathcal{I}$ is equivalent to (13).

For $\alpha \in \mathcal{I}$ and $\beta, \gamma \in \mathcal{I}^\perp$,

$$(17) \quad \begin{aligned} 2\varrho^*(A_\alpha \beta, \gamma) &= \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta) \\ &= -\varrho^*(\text{ad}_{r_\#(\beta)}^* \alpha, \gamma) - \varrho^*(\text{ad}_{r_\#(\beta)}^* \gamma, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \beta, \alpha) + \varrho^*(\text{ad}_{r_\#(\gamma)}^* \alpha, \beta) \\ &= -\prec \alpha, [r_\#(\beta), \#(\gamma)] \succ + \prec \gamma, [r_\#(\beta), \#(\alpha)] \succ - \prec \beta, [r_\#(\gamma), \#(\alpha)] \succ \\ &\quad - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = \varrho(\#(\gamma), [r_\#(\beta), \#(\alpha)]) \\ &\quad - \varrho(\#(\beta), [r_\#(\gamma), \#(\alpha)]) + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\varrho(J \circ r_\#(\gamma), [r_\#(\beta), \#(\alpha)]) + \varrho(J \circ r_\#(\beta), [r_\#(\gamma), \#(\alpha)]) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ \\ &= -\omega_r(r_\#(\gamma), \text{pr}_S([r_\#(\beta), \#(\alpha)])) - \omega_r(\text{pr}_S([r_\#(\gamma), \#(\alpha)]), r_\#(\beta)) \\ &\quad + \prec \alpha, [r_\#(\beta), \#(\gamma)] \succ - \prec \alpha, [r_\#(\gamma), \#(\beta)] \succ. \end{aligned}$$

Now, $\#(\beta), \#(\gamma) \in S$ and $r_\#(\beta), r_\#(\gamma) \in S$ and since S is a subalgebra we deduce that $[r_\#(\beta), \#(\gamma)], [r_\#(\gamma), \#(\beta)] \in S$ and hence

$$-\prec \alpha, [r_\#(\beta), \#(\gamma)] \succ = -\prec \alpha, [r_\#(\gamma), \#(\beta)] \succ = 0.$$

We have also $\#: \mathcal{I} \rightarrow S^\perp$ and $r_\#: \mathcal{I}^\perp \rightarrow S$ are isomorphisms so that, by virtue of (17), $A_\alpha \beta = 0$ for any $\alpha \in \mathcal{I}$ and $\beta \in \mathcal{I}^\perp$ is equivalent to (14).

On the other hand, for any $\alpha, \beta, \gamma \in \mathcal{I}^\perp$, since $\# = -J \circ r_\#$, the relation

$$2\varrho^*(A_\alpha \beta, \gamma) = \varrho^*([\alpha, \beta]_r, \gamma) + \varrho^*([\gamma, \beta]_r, \alpha) + \varrho^*([\gamma, \alpha]_r, \beta)$$

can be written

$$\begin{aligned} 2\varrho(J \circ r_\#(A_\alpha \beta), J \circ r_\#(\gamma)) &= \varrho(J \circ r_\#([\alpha, \beta]_r), J \circ r_\#(\gamma)) \\ &\quad + \varrho(J \circ r_\#([\gamma, \beta]_r), J \circ r_\#(\alpha)) \\ &\quad + \varrho(J \circ r_\#([\gamma, \alpha]_r), J \circ r_\#(\beta)). \end{aligned}$$

But $r_{\#}([\alpha, \beta]_r) = [r_{\#}(\alpha), r_{\#}(\beta)]$ and hence

$$2\langle r_{\#}(A_{\alpha}, \beta), r_{\#}(\gamma) \rangle_J = \langle [r_{\#}(\alpha), r_{\#}(\beta)], r_{\#}(\gamma) \rangle_J + \langle [r_{\#}(\mathfrak{g}), r_{\#}(\beta)], r_{\#}(\alpha) \rangle_J + \langle [r_{\#}(\gamma), r_{\#}(\alpha)], r_{\#}(\beta) \rangle_J,$$

where $\langle u, v \rangle_J = \varrho(Ju, Jv)$. This shows that $r_{\#}(A_{\alpha}\beta) = \nabla_{r_{\#}(\alpha)}r_{\#}(\beta)$ where ∇ is the Levi-Civita product of $(S, [\ , \], \langle \ , \ \rangle_J)$ and the third relation in (15) is equivalent to

$$\omega_r(\nabla_u v, w) + \omega_r(v, \nabla_u w) = 0, \quad u, v, w \in S.$$

This is equivalent to $\nabla_u Jv = J\nabla_u v$. Let us show that ∇ is actually the Levi-Civita product of $(S, [\ , \], \varrho)$. Indeed, for any $u, v, w \in S$, $\nabla_u v - \nabla_v u = [u, v]$ and

$$\begin{aligned} \varrho(\nabla_u v, w) + \varrho(\nabla_u w, v) &= \langle J^{-1}\nabla_u v, J^{-1}w \rangle_J + \langle J^{-1}\nabla_u w, J^{-1}v \rangle_J \\ &= \langle \nabla_u J^{-1}v, J^{-1}w \rangle_J + \langle \nabla_u J^{-1}w, J^{-1}v \rangle_J \\ &= 0. \end{aligned}$$

So we have shown the direct part of the theorem. The converse can be deduced easily from the relations we established in the proof of the direct part. \square

Example 1. Let G be a compact connected Lie group, \mathfrak{g} its Lie algebra and T an even dimensional torus of G . Choose a bi-invariant Riemannian metric $\langle \ , \ \rangle$ on G , a nondegenerate $\omega \in \wedge^2 S^*$ where S is the Lie algebra of T and put $\varrho = \langle \ , \ \rangle(e)$. Let $r \in \wedge^2 \mathfrak{g}$ be the solution of the classical Yang-Baxter associated to (S, ω) . By using either Proposition 3.2 or Theorem 3.1, one can see easily that $(\mathfrak{g}, \varrho, r)$ is a Riemann-Poisson Lie algebra and hence $(G, \langle \ , \ \rangle, \pi)$ is a Riemann-Poisson Lie group where π is the left invariant Poisson tensor associated to r . According to Theorem 1.1, the orbits of the right action of T on G defines a Riemannian and Kähler foliation. For instance, $G = \text{SO}(2n)$, $T = \text{Diagonal}(D_1, \dots, D_n)$ where $D_i = \begin{pmatrix} \cos(\theta_i) & \sin(\theta_i) \\ -\sin(\theta_i) & \cos(\theta_i) \end{pmatrix}$ and $\langle \ , \ \rangle = -K$ where K is the Killing form.

4. CONSTRUCTION OF RIEMANN-POISSON LIE ALGEBRAS

In this section, we give a general method for building Riemann-Poisson Lie algebras and we use it to give all Riemann-Poisson Lie algebras up to dimension 5.

According to Theorem 3.1, to build Riemann-Poisson Lie algebras one needs to solve the following problem.

Problem 1. We look for:

- (1) A Kähler Lie algebra $(\mathfrak{h}, [\ , \], \varrho_{\mathfrak{h}}, \omega)$,
- (2) a Euclidean vector space $(\mathfrak{p}, \varrho_{\mathfrak{p}})$,
- (3) a bilinear skew-symmetric map $[\ , \]_{\mathfrak{p}} : \mathfrak{p} \times \mathfrak{p} \longrightarrow \mathfrak{p}$,
- (4) a bilinear skew-symmetric map $\mu : \mathfrak{p} \times \mathfrak{p} \longrightarrow \mathfrak{h}$,
- (5) two linear maps $\phi_{\mathfrak{p}} : \mathfrak{p} \longrightarrow \text{sp}(\mathfrak{h}, \omega)$ and $\phi_{\mathfrak{h}} : \mathfrak{h} \longrightarrow \text{so}(\mathfrak{p})$ where $\text{sp}(\mathfrak{h}, \omega) = \{J : \mathfrak{h} \longrightarrow \mathfrak{h}, J^{\omega} + J = 0\}$ and $\text{so}(\mathfrak{p}) = \{A : \mathfrak{p} \longrightarrow \mathfrak{p}, A^* + A = 0\}$, J^{ω} is the adjoint with respect to ω and A^* is the adjoint with respect to $\varrho_{\mathfrak{p}}$,

such that the bracket $[\ , \]$ on $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ given, for any $a, b \in \mathfrak{p}$ and $u, v \in \mathfrak{h}$, by

$$(18) \quad [u, v] = [u, v]_{\mathfrak{h}}, [a, b] = \mu(a, b) + [a, b]_{\mathfrak{p}}, [a, u] = -[u, a] = \phi_{\mathfrak{p}}(a)(u) - \phi_{\mathfrak{h}}(u)(a)$$

is a Lie bracket.

In this case, $(\mathfrak{g}, [\ , \])$ endowed with $r \in \wedge^2 \mathfrak{g}$ associated to (\mathfrak{h}, ω) and the Euclidean product $\varrho = \varrho_{\mathfrak{h}} \oplus \varrho_{\mathfrak{p}}$ becomes, by virtue of Theorem 3.1, a Riemann-Poisson Lie algebra.

Proposition 4.1. *With the data and notations of Problem 1, the bracket given by (18) is a Lie bracket if and only if, for any $u, v \in \mathfrak{h}$ and $a, b, c \in \mathfrak{p}$,*

$$(19) \quad \begin{cases} \phi_{\mathfrak{p}}(a)([u, v]_{\mathfrak{h}}) = [u, \phi_{\mathfrak{p}}(a)(v)]_{\mathfrak{h}} + [\phi_{\mathfrak{p}}(a)(u), v]_{\mathfrak{h}} + \phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(v)(a))(u) - \phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(u)(a))(v), \\ \phi_{\mathfrak{h}}(u)([a, b]_{\mathfrak{p}}) = [a, \phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}} + [\phi_{\mathfrak{h}}(u)(a), b]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b), \\ \phi_{\mathfrak{h}}([u, v]_{\mathfrak{h}}) = [\phi_{\mathfrak{h}}(u), \phi_{\mathfrak{h}}(v)], \\ \phi_{\mathfrak{p}}([a, b]_{\mathfrak{p}})(u) = [\phi_{\mathfrak{p}}(a), \phi_{\mathfrak{p}}(b)](u) + [u, \mu(a, b)]_{\mathfrak{h}} - \mu(a, \phi_{\mathfrak{h}}(u)(b)) - \mu(\phi_{\mathfrak{h}}(u)(a), b), \\ \mathcal{F}[a, [b, c]_{\mathfrak{p}}]_{\mathfrak{p}} = \mathcal{F} \phi_{\mathfrak{h}}(\mu(b, c))(a), \\ \mathcal{F} \phi_{\mathfrak{p}}(a)(\mu(b, c)) = \mathcal{F} \mu([b, c]_{\mathfrak{p}}, a), \end{cases}$$

where \mathcal{F} stands for the circular permutation.

Proof. The equations follow from the Jacobi identity applied to (a, u, v) , (a, b, u) and (a, b, c) . □

We tackle now the task of determining the list of all Riemann-Poisson Lie algebras up to dimension 5. For this purpose, we need to solve Problem 1 in the following four cases: (a) $\dim \mathfrak{p} = 1$, (b) $\dim \mathfrak{h} = 2$ and \mathfrak{h} non abelian, (c) $\dim \mathfrak{h} = \dim \mathfrak{p} = 2$ and \mathfrak{h} abelian, (d) $\dim \mathfrak{h} = 2$, $\dim \mathfrak{p} = 3$ and \mathfrak{h} abelian.

It is easy to find the solutions of Problem 1 when $\dim \mathfrak{p} = 1$ since in this case $\text{so}(\mathfrak{p}) = 0$ and the three last equations in (19) hold obviously.

Proposition 4.2. *If $\dim \mathfrak{p} = 1$ then the solutions of Problem 1 are a Kähler Lie algebra $(\mathfrak{h}, \varrho, \omega)$, $\phi_{\mathfrak{h}} = 0$, $[\ , \]_{\mathfrak{p}} = 0$, $\mu = 0$ and $\phi_{\mathfrak{p}}(a) \in \text{sp}(\mathfrak{h}, \omega) \cap \text{Der}(\mathfrak{h})$ where a is a generator of \mathfrak{p} and $\text{Der}(\mathfrak{h})$ the Lie algebra of derivations of \mathfrak{h} .*

Let us solve Problem 1 when \mathfrak{h} is 2-dimensional non abelian.

Proposition 4.3. *Let $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\ , \]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ be a solution of Problem 1 with \mathfrak{h} is 2-dimensional non abelian. Then there exists an orthonormal basis $\mathbb{B} = (e_1, e_2)$ of \mathfrak{h} , $b_0 \in \mathfrak{p}$ and two constants $\alpha \neq 0$ and $\beta \neq 0$ such that:*

- (i) $[e_1, e_2]_{\mathfrak{h}} = \alpha e_1, \omega = \beta e_1^* \wedge e_2^*$,
- (ii) $(\mathfrak{p}, [\ , \]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is a Euclidean Lie algebra,
- (iii) $\phi_{\mathfrak{h}}(e_1) = 0, \phi_{\mathfrak{h}}(e_2) \in \text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$ and, for any $a \in \mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(a, b_0) \\ 0 & 0 \end{pmatrix}$,
- (iv) for any $a, b \in \mathfrak{p}$, $\mu(a, b) = \mu_0(a, b)e_1$ with μ_0 is a 2-cocycle of $(\mathfrak{p}, [\ , \]_{\mathfrak{p}})$ satisfying

$$(20) \quad \mu_0(a, \phi_{\mathfrak{h}}(e_2)b) + \mu_0(\phi_{\mathfrak{h}}(e_2)a, b) = -\varrho_{\mathfrak{p}}([a, b]_{\mathfrak{p}}, b_0) - \alpha \mu_0(a, b).$$

Proof. Note first that from the third relation in (19) we get that $\phi_{\mathfrak{h}}(\mathfrak{h})$ is a solvable subalgebra of $\mathfrak{so}(\mathfrak{p})$ and hence must be abelian. Since \mathfrak{h} is 2-dimensional non abelian then $\dim \phi_{\mathfrak{h}}(\mathfrak{h}) = 1$ and $[\mathfrak{h}, \mathfrak{h}] \subset \ker \phi_{\mathfrak{h}}$. So there exists an orthonormal basis (e_1, e_2) of \mathfrak{h} such that $[e_1, e_2]_{\mathfrak{h}} = \alpha e_1$, $\phi_{\mathfrak{h}}(e_1) = 0$ and $\omega = \beta e_1^* \wedge e_2^*$. If we identify the endomorphisms of \mathfrak{h} with their matrices in the basis (e_1, e_2) , we get that $\mathfrak{sp}(\mathfrak{h}, \omega) = \mathfrak{sl}(2, \mathbb{R})$ and there exists $a_0, b_0, c_0 \in \mathfrak{p}$ such that, for any $a \in \mathfrak{p}$,

$$\phi_{\mathfrak{p}}(a) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}.$$

The first equation in (19) is equivalent to

$$\begin{aligned} \alpha(\varrho_{\mathfrak{p}}(a_0, a)e_1 + \varrho_{\mathfrak{p}}(c_0, a)e_2) &= -\alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 + \alpha\varrho_{\mathfrak{p}}(a_0, a)e_1 \\ &\quad + \varrho_{\mathfrak{p}}(a_0, \phi_{\mathfrak{h}}(e_2)(a))e_1 + \varrho_{\mathfrak{p}}(c_0, \phi_{\mathfrak{h}}(e_2)(a))e_2, \end{aligned}$$

for any $a \in \mathfrak{p}$. Since $\phi_{\mathfrak{h}}(e_2)$ is sekw-symmetric, this is equivalent to

$$\phi_{\mathfrak{h}}(e_2)(a_0) = -\alpha a_0 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(c_0) = -\alpha c_0.$$

This implies that $a_0 = c_0 = 0$. The second equation in (19) implies that $\phi_{\mathfrak{h}}(e_2)$ is a derivation of $[\cdot, \cdot]_{\mathfrak{p}}$. If we take $u = e_1$ in the forth equation in (19), we get that $[e_1, \mu(a, b)] = 0$, for any $a, b \in \mathfrak{p}$ and hence $\mu(a, b) = \mu_0(a, b)e_1$. If we take $u = e_2$ in the forth equation in (19) we get (20). The two last equations are equivalent to $[\cdot, \cdot]_{\mathfrak{p}}$ is a Lie bracket and μ_0 is 2-cocycle of $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$. \square

The following proposition gives the solutions of Problem 1 when \mathfrak{h} is 2-dimensional abelian and $\dim \mathfrak{p} = 2$.

Proposition 4.4. *Let $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ be a solution of Problem 1 with \mathfrak{h} is 2-dimensional abelian and $\dim \mathfrak{p} = 2$. Then one of the following situations occurs:*

- (1) $\phi_{\mathfrak{h}} = 0$, $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is a 2-dimensional Euclidean Lie algebra, there exists $a_0 \in \mathfrak{p}$ and $D \in \mathfrak{sp}(\mathfrak{h}, \omega)$ such that, for any $a \in \mathfrak{p}$, $\phi_{\mathfrak{p}}(a) = \varrho_{\mathfrak{p}}(a_0, a)D$ and there is no restriction on μ . Moreover, $a_0 \in [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}^{\perp}$ if $D \neq 0$.
- (2) $\phi_{\mathfrak{h}} = 0$, $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is a 2-dimensional non abelian Euclidean Lie algebra, $\phi_{\mathfrak{p}}$ identifies \mathfrak{p} to a two dimensional subalgebra of $\mathfrak{sp}(\mathfrak{h}, \omega)$ and there is no restriction on μ .
- (3) $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is a Euclidean abelian Lie algebra and there exists an orthonormal basis $\mathbb{B} = (e_1, e_2)$ of \mathfrak{h} and $b_0 \in \mathfrak{p}$ such that $\omega = \alpha e_1^* \wedge e_2^*$, $\phi_{\mathfrak{h}}(e_1) = 0$, $\phi_{\mathfrak{h}}(e_2) \neq 0$ and, for any $a \in \mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$ and there is no restriction on μ .

Proof. Note first that since $\dim \mathfrak{p} = 2$ the last two equations in (19) hold obviously and $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$ is a Lie algebra. We distinguish two cases:

- (i) $\phi_{\mathfrak{h}} = 0$. Then (19) is equivalent to $\phi_{\mathfrak{p}}$ is a representation of \mathfrak{p} in $\mathfrak{sp}(\mathfrak{h}, \omega) \simeq \mathfrak{sl}(2, \mathbb{R})$. Since $\mathfrak{sl}(2, \mathbb{R})$ doesn't contain any abelian two dimensional subalgebra, if \mathfrak{p} is an abelian Lie algebra then $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) \leq 1$ and the first situation occurs. If \mathfrak{p} is not abelian then the first or the second situation occurs depending on $\dim \phi_{\mathfrak{p}}(\mathfrak{p})$.

(ii) $\phi_{\mathfrak{h}} \neq 0$. Since $\dim \text{so}(\mathfrak{p}) = 1$ there exists an orthonormal basis $\mathbb{B} = (e_1, e_2)$ of \mathfrak{h} such that $\phi_{\mathfrak{h}}(e_1) = 0$ and $\phi_{\mathfrak{h}}(e_2) \neq 0$. We have $\text{sp}(\mathfrak{h}, \omega) = \text{sl}(2, \mathbb{R})$ and hence, for any $a \in \mathfrak{p}$, $M(\phi_{\mathfrak{p}}(a), \mathbb{B}) = \begin{pmatrix} \varrho_{\mathfrak{p}}(a_0, a) & \varrho_{\mathfrak{p}}(b_0, a) \\ \varrho_{\mathfrak{p}}(c_0, a) & -\varrho_{\mathfrak{p}}(a_0, a) \end{pmatrix}$. Choose an orthonormal basis (a_1, a_2) of \mathfrak{p} . Then there exists $\lambda \neq 0$ such that $\phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2$ and $\phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1$.

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1) = 0, \quad a \in \mathfrak{p}.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1) = \phi_{\mathfrak{p}}(a_2)(e_1) = 0.$$

Then $a_0 = c_0 = 0$ and hence $\phi_{\mathfrak{p}}(a) = \begin{pmatrix} 0 & \varrho_{\mathfrak{p}}(b_0, a) \\ 0 & 0 \end{pmatrix}$. The second equation in (19) gives

$$\begin{aligned} \phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) &= [a_1, \phi_{\mathfrak{h}}(e_2)(a_2)]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(e_2)(a_1), a_2]_{\mathfrak{p}} \\ &\quad + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_2)(e_2))(a_2) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_1)(e_2))(a_2), \end{aligned}$$

and hence $\phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = 0$. Thus $[a_1, a_2]_{\mathfrak{p}} = 0$. All the other equations in (19) hold obviously. □

To tackle the last case, we need the determination of 2-dimensional subalgebras of $\text{sl}(2, \mathbb{R})$.

Proposition 4.5. *The 2-dimensional subalgebras of $\text{sl}(2, \mathbb{R})$ are*

$$\begin{aligned} \mathfrak{g}_1 &= \left\{ \begin{pmatrix} \alpha & \beta \\ 0 & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\}, \quad \mathfrak{g}_2 = \left\{ \begin{pmatrix} \alpha & 0 \\ \beta & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\}, \\ \mathfrak{g}_x &= \left\{ \begin{pmatrix} \alpha & \frac{2\beta - \alpha}{x} \\ (\alpha + 2\beta)x & -\alpha \end{pmatrix}, \alpha, \beta \in \mathbb{R} \right\} \end{aligned}$$

where $x \in \mathbb{R} \setminus \{0\}$. Moreover, $\mathfrak{g}_x = \mathfrak{g}_y$ if and only if $x = y$.

Proof. Let \mathfrak{g} be a 2-dimensional subalgebra of $\text{sl}(2, \mathbb{R})$. We consider the basis $\mathbb{B} = (h, e, f)$ of $\text{sl}(2, \mathbb{R})$ given by

$$e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad f = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad h = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Then

$$[h, e] = 2e, \quad [h, f] = -2f \quad \text{and} \quad [e, f] = h.$$

If $h \in \mathfrak{g}$ then ad_h leaves \mathfrak{g} invariant. But ad_h has three eigenvalues $(0, 2, -2)$ with the associated eigenvectors (h, e, f) and hence its restriction to \mathfrak{g} has $(0, 2)$ or $(0, -2)$ as eigenvalues. Thus $\mathfrak{g} = \mathfrak{g}_1$ or $\mathfrak{g} = \mathfrak{g}_2$.

Suppose now that $h \notin \mathfrak{g}$. By using the fact that $\text{sl}(2, \mathbb{R})$ is unimodular, i.e., for any $w \in \text{sl}(2, \mathbb{R})$ $\text{tr}(\text{ad}_w) = 0$, we can choose a basis (u, v) of \mathfrak{g} such that (u, v, h) is a basis of $\text{sl}(2, \mathbb{R})$ and

$$[u, v] = u, \quad [h, u] = au + v \quad \text{and} \quad [h, v] = du - av - h.$$

If (x_1, x_2, x_3) and (y_1, y_2, y_3) are the coordinates of u and v in \mathbb{B} , the brackets above gives

$$\begin{cases} -2(x_1y_3 - x_3y_1) - x_1 = 0, \\ 2(x_2y_3 - x_3y_2) - x_2 = 0, \\ x_1y_2 - x_2y_1 - x_3 = 0, \end{cases} \quad \begin{cases} y_1 = (2-a)x_1, \\ y_2 = -(a+2)x_2, \\ y_3 = -ax_3, \end{cases} \quad \text{and} \quad \begin{cases} dx_1 = (a+2)y_1, \\ dx_2 = (a-2)y_2, \\ dx_3 = ay_3 + 1. \end{cases}$$

Note first that if $x_1 = 0$ then $(x_2, x_3) = (0, 0)$ which impossible so we must have $x_1 \neq 0$ and hence $d = 4 - a^2$. If we replace in the third equation in the second system and the last equation, we get $x_3 = \frac{1}{4}$ and $y_3 = -\frac{a}{4}$. The third equation in the first system gives $x_2 = -\frac{1}{16x_1}$ and hence $y_1 = (2-a)x_1$ and $y_2 = \frac{(a+2)}{16x_1}$. Thus

$$\begin{aligned} \mathfrak{g} &= \text{span} \left\{ \begin{pmatrix} \frac{1}{4} & -\frac{1}{16x_1} \\ x_1 & -\frac{1}{4} \end{pmatrix}, \begin{pmatrix} -\frac{a}{4} & \frac{(a+2)}{16x_1} \\ (2-a)x_1 & \frac{a}{4} \end{pmatrix} \right\} \\ &= \text{span} \left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix} \right\}; \quad x = 4x_1. \end{aligned}$$

But

$$\begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} = a \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix} + \begin{pmatrix} -a & \frac{(a+2)}{x} \\ (2-a)x & a \end{pmatrix}$$

and hence

$$\mathfrak{g} = \text{span} \left\{ \begin{pmatrix} 1 & -\frac{1}{x} \\ x & -1 \end{pmatrix}, \begin{pmatrix} 0 & \frac{2}{x} \\ 2x & 0 \end{pmatrix} \right\} = \mathfrak{g}_x.$$

One can check easily that $\mathfrak{g}_x = \mathfrak{g}_y$ if and only if $x = y$. This completes the proof. \square

The following two propositions give the solutions of Problem 1 when \mathfrak{h} is 2-dimensional abelian and $\dim \mathfrak{p} = 3$.

Proposition 4.6. *Let $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ be a solution of Problem 1 with \mathfrak{h} is 2-dimensional abelian and $\dim \mathfrak{p} = 3$ and $\phi_{\mathfrak{h}} = 0$. Then one of the following situations occurs:*

- (i) $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is 3-dimensional Euclidean Lie algebra, $\phi_{\mathfrak{p}} = 0$ and μ is 2-cocycle for the trivial representation.
- (ii) $\phi_{\mathfrak{p}}$ is an isomorphism of Lie algebras between $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$ and $\mathfrak{sl}(2, \mathbb{R})$ and there exists an endomorphism $L: \mathfrak{p} \rightarrow \mathfrak{h}$ such that for any $a, b \in \mathfrak{p}$,

$$\mu(a, b) = \phi_{\mathfrak{p}}(a)(L(b)) - \phi_{\mathfrak{p}}(b)(L(a)) - L([a, b]_{\mathfrak{p}}).$$

- (iii) There exists a basis $\mathbb{B}_{\mathfrak{p}} = (a_1, a_2, a_3)$ of \mathfrak{p} , $\alpha \neq 0, \beta \neq 0, \gamma, \tau \in \mathbb{R}$ such that $[\cdot, \cdot]_{\mathfrak{p}}$ has one of the two following forms

$$\begin{cases} [a_1, a_2]_{\mathfrak{p}} = 0, [a_1, a_3]_{\mathfrak{p}} = \beta a_1, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \alpha \neq 0, \beta \neq 0 \\ M(\varrho_{\mathfrak{p}}, \mathbb{B}_{\mathfrak{p}}) = I_3 \end{cases} \quad \text{or} \quad \begin{cases} [a_1, a_2]_{\mathfrak{p}} = [a_1, a_3]_{\mathfrak{p}} = 0, \\ [a_2, a_3]_{\mathfrak{p}} = \alpha a_2, \alpha \neq 0, \\ M(\varrho_{\mathfrak{p}}, \mathbb{B}_{\mathfrak{p}}) = \begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}. \end{cases}$$

In both cases, there exists an orthonormal basis $\mathbb{B}_{\mathfrak{h}} = (e_1, e_2)$ of \mathfrak{h} , $x \neq 0$, $u \neq 0$ and $v \in \mathbb{R}$ such that $\phi_{\mathfrak{p}}$ has one of the following forms

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0, \end{cases} \quad \text{or} \quad \begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0, \end{cases}$$

$$\begin{cases} M(\phi_{\mathfrak{p}}(a_2), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \\ M(\phi_{\mathfrak{p}}(a_3), \mathbb{B}_{\mathfrak{h}}) = \begin{pmatrix} v & -\frac{2v+\alpha}{2x} \\ \frac{2v-\alpha}{2}x & -v \end{pmatrix}, \\ \phi_{\mathfrak{p}}(a_1) = 0. \end{cases}$$

Moreover, μ is a 2-cocycle for $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

(iv) There exists an orthonormal basis $\mathbb{B} = (a_1, a_2, a_3)$ of \mathfrak{p} such that $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$, $\phi_{\mathfrak{p}}(a_3)$ is a non zero element of $\text{sp}(\mathfrak{h}, \omega)$ and

$$\begin{cases} [a_1, a_2]_{\mathfrak{p}} = 0, [a_1, a_3]_{\mathfrak{p}} = \beta a_1 + \rho a_2, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \end{cases} \quad \text{or} \quad \begin{cases} [a_1, a_2]_{\mathfrak{p}} = \alpha a_2, [a_1, a_3]_{\mathfrak{p}} = \rho a_2, \\ [a_2, a_3]_{\mathfrak{p}} = \gamma a_2, \alpha \neq 0. \end{cases}$$

Moreover, μ is a 2-cocycle for $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

Proof. In this case, (19) is equivalent to $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}})$ is a Lie algebra and $\phi_{\mathfrak{p}}$ is a representation and μ is a 2-cocycle of $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \phi_{\mathfrak{p}})$.

We distinguish four cases:

- (1) $\phi_{\mathfrak{p}} = 0$ and the case (i) occurs.
- (2) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 3$ and hence \mathfrak{p} is isomorphic to $\text{sp}(\mathfrak{h}, \omega) \simeq \text{sl}(2, \mathbb{R})$ and hence μ is a coboundary. Thus (ii) occurs.
- (3) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 2$ then $\ker \phi_{\mathfrak{p}}$ is a one dimensional ideal of \mathfrak{p} . But $\phi_{\mathfrak{p}}(\mathfrak{p})$ is a 2-dimensional subalgebra of $\text{sp}(\mathfrak{h}, \omega) \simeq \text{sl}(2, \mathbb{R})$, therefore it is non abelian so $\mathfrak{p}/\ker \mathfrak{p}$ is non abelian.

If $\ker \mathfrak{p} \subset [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$ then $\dim[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = 2$ so there exists an orthonormal basis (a_1, a_2, a_3) of \mathfrak{p} such that $a_1 \in \ker \mathfrak{p}$ and

$$[a_1, a_2]_{\mathfrak{p}} = \xi a_1, [a_1, a_3]_{\mathfrak{p}} = \beta a_1 \quad \text{and} \quad [a_2, a_3]_{\mathfrak{p}} = \gamma a_1 + \alpha a_2, \alpha \neq 0, \beta \neq 0$$

and we must have $\xi = 0$ in order to have the Jacobi identity.

If $\ker \mathfrak{p} \not\subset [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$ then $\ker \mathfrak{p} \subset Z(\mathfrak{p})$ and $\dim[\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}} = 1$. Then there exists a basis (a_1, a_2, a_3) of \mathfrak{p} such that $a_1 \in \ker \mathfrak{p}$, $a_2 \in [\mathfrak{p}, \mathfrak{p}]_{\mathfrak{p}}$, $a_3 \in \{a_1, a_2\}^{\perp}$ and

$$[a_2, a_3]_{\mathfrak{p}} = \alpha a_2, [a_3, a_1]_{\mathfrak{p}} = [a_1, a_2]_{\mathfrak{p}} = 0, \alpha \neq 0.$$

The matrix of $\varrho_{\mathfrak{p}}$ in (a_1, a_2, a_3) is given by

$$\begin{pmatrix} 1 & \tau & 0 \\ \tau & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

We choose an orthonormal basis (e_1, e_2) of \mathfrak{h} and identify $\text{sp}(\mathfrak{h}, \omega)$ to $\text{sl}(2, \mathbb{R})$. Now $\phi_{\mathfrak{p}}(\mathfrak{p}) = \{\phi_p(a_2), \phi_p(a_3)\}$ is a subalgebra of $\text{sl}(2, \mathbb{R})$ and, according to Proposition 4.5, $\phi_{\mathfrak{p}}(\mathfrak{p}) = \mathfrak{g}_1, \mathfrak{g}_2$ or \mathfrak{g}_x . But

$$[\mathfrak{g}_1, \mathfrak{g}_1] = \mathbb{R}e, [\mathfrak{g}_2, \mathfrak{g}_2] = \mathbb{R}f \quad \text{and} \quad [\mathfrak{g}_x, \mathfrak{g}_x] = \left\{ \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix} \right\}.$$

So in order for $\phi_{\mathfrak{p}}$ to be a representation we must have

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & u \\ 0 & 0 \end{pmatrix}, \quad \text{and} \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} -\frac{\alpha}{2} & v \\ 0 & \frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}, \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} \frac{\alpha}{2} & 0 \\ v & -\frac{\alpha}{2} \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0,$$

or

$$\phi_{\mathfrak{p}}(a_2) = \begin{pmatrix} u & -\frac{u}{x} \\ ux & -u \end{pmatrix}, \quad \phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} p & -\frac{2p+\alpha}{2x} \\ \frac{2p-\alpha}{2}x & -p \end{pmatrix} \quad \text{and} \quad \phi_{\mathfrak{p}}(a_1) = 0.$$

(4) $\dim \phi_{\mathfrak{p}}(\mathfrak{p}) = 1$ then $\ker \phi_{\mathfrak{p}}$ is a two dimensional ideal of \mathfrak{p} . Then there exists an orthonormal basis (a_1, a_2, a_3) of \mathfrak{p} such that

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_2, \quad [a_3, a_1]_{\mathfrak{p}} = pa_1 + qa_2 \quad \text{and} \quad [a_3, a_2]_{\mathfrak{p}} = ra_1 + sa_2.$$

The Jacobi identity gives $\alpha = 0$ or $(p, r) = (0, 0)$. We take $\phi_{\mathfrak{p}}(a_1) = \phi_{\mathfrak{p}}(a_2) = 0$ and $\phi_{\mathfrak{p}}(a_3) \in \text{sl}(2, \mathbb{R})$. □

Proposition 4.7. *Let $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ be a solution of Problem 1 with \mathfrak{h} is 2-dimensional abelian, $\dim \mathfrak{p} = 3$ and $\phi_{\mathfrak{h}} \neq 0$. Then there exists an orthonormal basis (e_1, e_2) of \mathfrak{h} , an orthonormal basis (a_1, a_2, a_3) of \mathfrak{p} , $\lambda > 0$, $\alpha, p, q, \mu_1, \mu_2, \mu_3 \in \mathbb{R}$ such that*

$$\phi_{\mathfrak{h}}(e_1) = 0, \quad \phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2, \quad \phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3) = 0,$$

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_3, \quad [a_1, a_3]_{\mathfrak{p}} = pa_1 + qa_2, \quad [a_2, a_3]_{\mathfrak{p}} = -qa_1 + pa_2 \quad \text{and}$$

$$\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}, \quad i = 1, 2, 3$$

and one of the following situations occurs:

(i) $p \neq 0, \alpha = 0$ and

$$\mu(a_1, a_2) = 0, \quad \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \quad \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(ii) $p = 0, \mu_3 \neq 0, \alpha = 0$ and

$$\mu(a_1, a_2) = ce_1, \quad \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \quad \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

(iii) $p = 0, \mu_3 = 0$ and

$$\begin{aligned} \mu(a_1, a_2) &= c_1e_1 + c_2e_2, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) &= \lambda^{-1}(-q\mu_1 + p\mu_2)e_1. \end{aligned}$$

Proof. Since $\phi_{\mathfrak{h}} \neq 0$ then $\phi_{\mathfrak{h}}(\mathfrak{h})$ is a non trivial abelian subalgebra of $\mathfrak{so}(\mathfrak{p})$ and hence it must be one dimensional. Then there exists an orthonormal basis (e_1, e_2) of \mathfrak{h} and an orthonormal basis (a_1, a_2, a_3) of \mathfrak{p} and $\lambda > 0$ such that $\phi_{\mathfrak{h}}(e_1) = 0$ and

$$\phi_{\mathfrak{h}}(e_2)(a_1) = \lambda a_2, \phi_{\mathfrak{h}}(e_2)(a_2) = -\lambda a_1 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)(a_3) = 0.$$

The first equation in (19) is equivalent to

$$\phi_{\mathfrak{p}}(\phi_{\mathfrak{h}}(e_2)(a))(e_1) = 0, \quad a \in \mathfrak{p}.$$

This is equivalent to

$$\phi_{\mathfrak{p}}(a_1)(e_1) = \phi_{\mathfrak{p}}(a_2)(e_1) = 0.$$

Thus $\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}$ for $i = 1, 2$ and $\phi_{\mathfrak{p}}(a_3) = \begin{pmatrix} u & v \\ w & -u \end{pmatrix}$. Consider now the second equation in (19)

$$\phi_{\mathfrak{h}}(u)([a, b]_{\mathfrak{p}}) = [a, \phi_{\mathfrak{h}}(u)(b)]_{\mathfrak{p}} + [\phi_{\mathfrak{h}}(u)(a), b]_{\mathfrak{p}} + \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(b)(u))(a) - \phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a)(u))(b).$$

This equation is obviously true when $u = e_1$ and $(a, b) = (a_1, a_2)$. For $u = e_1$ and $(a, b) = (a_1, a_3)$, we get

$$\phi_{\mathfrak{h}}(\phi_{\mathfrak{p}}(a_3)(e_1))(a_1) = 0$$

and hence $w = 0$.

For $u = e_2$ and $(a, b) = (a_1, a_2)$, we get $\phi_{\mathfrak{h}}(e_2)([a_1, a_2]_{\mathfrak{p}}) = 0$ and hence $[a_1, a_2]_{\mathfrak{p}} = \alpha a_3$.

For $u = e_2$ and $(a, b) = (a_1, a_3)$ or $(a, b) = (a_2, a_3)$, we get

$$\phi_{\mathfrak{h}}(e_2)([a_1, a_3]_{\mathfrak{p}}) = \lambda[a_2, a_3]_{\mathfrak{p}} - \lambda u a_2 \quad \text{and} \quad \phi_{\mathfrak{h}}(e_2)([a_2, a_3]_{\mathfrak{p}}) = -\lambda[a_1, a_3]_{\mathfrak{p}} + \lambda u a_1.$$

This implies that $[a_1, a_3]_{\mathfrak{p}}, [a_2, a_3]_{\mathfrak{p}} \in \text{span}\{a_1, a_2\}$ and hence

$$[a_1, a_3]_{\mathfrak{p}} = p a_1 + q a_2 \quad \text{and} \quad [a_2, a_3]_{\mathfrak{p}} = r a_1 + s a_2.$$

So

$$\begin{cases} \lambda(p a_2 - q a_1) = \lambda(r a_1 + s a_2 - u a_2), \\ \lambda(r a_2 - s a_1) = -\lambda(p a_1 + q a_2 - u a_1). \end{cases}$$

This is equivalent to

$$u = 0, p = s \quad \text{and} \quad r = -q.$$

To summarize, we get

$$[a_1, a_2]_{\mathfrak{p}} = \alpha a_3, [a_1, a_3]_{\mathfrak{p}} = p a_1 + q a_2, [a_2, a_3]_{\mathfrak{p}} = -q a_1 + p a_2 \quad \text{and}$$

$$\phi_{\mathfrak{p}}(a_i) = \begin{pmatrix} 0 & \mu_i \\ 0 & 0 \end{pmatrix}.$$

Let consider now the fourth equation in (19)

$$\phi_{\mathfrak{p}}([a, b]_{\mathfrak{p}})(u) = [\phi_{\mathfrak{p}}(a), \phi_{\mathfrak{p}}(b)](u) + [u, \mu(a, b)]_{\mathfrak{h}} - \mu(a, \phi_{\mathfrak{h}}(u)(b)) - \mu(\phi_{\mathfrak{h}}(u)(a), b).$$

This equation is obviously true for $u = e_1$.

For $u = e_2$ and $(a, b) = (a_1, a_2)$, $(a, b) = (a_1, a_3)$ or $(a, b) = (a_2, a_3)$, we get

$$\begin{cases} \alpha\mu_3 = 0, \\ (p\mu_1 + q\mu_2)e_1 = -\lambda\mu(a_2, a_3), \\ (-q\mu_1 + p\mu_2)e_1 = \lambda\mu(a_1, a_3). \end{cases}$$

The last two equations are equivalent to

$$\phi_{\mathfrak{p}}(a_3)(\mu(a_1, a_2)) = -2p\mu(a_1, a_2) \quad \text{and} \quad p[a_1, a_2]_{\mathfrak{p}} = 0.$$

• $p \neq 0$ then

$$\alpha = 0, \mu(a_1, a_2) = 0, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

• $p = 0$ and $\mu_3 \neq 0$ then $\alpha = 0$ and

$$\mu(a_1, a_2) = ce_1, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

• $p = 0$ and $\mu_3 = 0$ then

$$\mu(a_1, a_2) = c_1e_1 + c_2e_2, \mu(a_2, a_3) = -\lambda^{-1}(p\mu_1 + q\mu_2)e_1 \quad \text{and} \\ \mu(a_1, a_3) = \lambda^{-1}(-q\mu_1 + p\mu_2)e_1.$$

□

By using Propositions 4.2–4.7, we can give all the Riemann-Poisson Lie algebras of dimension 3, 4 or 5.

Let $(\mathfrak{g}, [\cdot, \cdot], \varrho, r)$ be a Riemann-Poisson Lie algebra of dimension less or equal to 5. According to what above then $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{p}$ and the Lie bracket on \mathfrak{g} is given by (18) and $((\mathfrak{h}, \omega, \varrho_{\mathfrak{h}}), (\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}}), \mu, \phi_{\mathfrak{h}}, \phi_{\mathfrak{p}})$ are solutions of Problem 1.

• $\dim \mathfrak{g} = 3$. In this case $\dim \mathfrak{h} = 2$ and $\dim \mathfrak{p} = 1$ and, by applying Proposition 4.2, the Lie bracket of \mathfrak{g} , ϱ and r are given in Table 1, where $e^{12} = e_1 \wedge e_2$.

Non vanishing Lie brackets	Bivector r	Matrix of ϱ	Conditions
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1$	αe^{12}	I_3	$a \neq 0, \alpha \neq 0$
$[e_3, e_1] = -be_1 + ce_2, [e_3, e_2] = de_1 + be_2$	αe^{12}	I_3	$\alpha \neq 0$

TAB. 1. Three dimensional Riemann-Poisson Lie algebras

• $\dim \mathfrak{g} = 4$. We have three cases:

(c41) $\dim \mathfrak{h} = 2$, $\dim \mathfrak{p} = 2$ and \mathfrak{h} is non abelian and we can apply Proposition 4.3 to get the Lie brackets on \mathfrak{g} , ϱ and r . They are described in rows 1 and 2 in Table 2.

(c42) $\dim \mathfrak{h} = 2$, $\dim \mathfrak{p} = 2$ and \mathfrak{h} is abelian and we can apply Propositions 4.4 and 4.5 to get the Lie brackets on \mathfrak{g} , ϱ and r . They are described in rows 3 and 8 in Table 2.

(c43) $\dim \mathfrak{h} = 4$. In this case \mathfrak{g} is a Kähler Lie algebra. We have used [12] to derive all four dimensional Kähler Lie algebra together with their symplectic derivations. The results are given in Table 3. The notation $\text{Der}^s(\mathfrak{h})$ stands

for the vector spaces of derivations which are skew-symmetric with respect the symplectic form. The vector space $\text{Der}^s(\mathfrak{h})$ is described by a family of generators and E_{ij} is the matrix with 1 in the i row and j column and 0 elsewhere.

Non vanishing Lie brackets	Bivector r	Matrix of ϱ	Conditions
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1 + ce_4,$ $[e_4, e_2] = de_1 - ce_3$	αe^{12}	I_4	$a \neq 0, \alpha \neq 0$
$[e_1, e_2] = ae_1, [e_3, e_2] = be_1,$ $[e_4, e_2] = de_1, [e_3, e_4] = ce_3 - a^{-1}cbe_1$	αe^{12}	I_4	$\alpha ac \neq 0,$
$[e_3, e_4] = ae_1 + be_2$	αe^{12}	I_4	$\alpha \neq 0$
$[e_3, e_4] = ae_1 + be_2 + ce_3, [e_4, e_1] = xe_1 + ye_2,$ $[e_4, e_2] = ze_1 - xe_2$	αe^{12}	I_4	$\alpha \neq 0$
$[e_3, e_4] = ae_1 + be_2 + 2e_4, [e_3, e_1] = e_1,$ $[e_3, e_2] = -e_2, [e_4, e_2] = e_1$	αe^{12}	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2$
$[e_3, e_4] = ae_1 + be_2 - 2e_4, [e_3, e_1] = e_1,$ $[e_3, e_2] = -e_2, [e_4, e_1] = e_2$	αe^{12}	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2$
$[e_3, e_4] = ae_1 + be_2 - 2e_3, [e_3, e_1] = e_1 + xe_2,$ $[e_3, e_2] = -\frac{1}{x}e_1 - e_2, [e_4, e_1] = xe_2, [e_4, e_2] = \frac{1}{x}e_1$	αe^{12}	$\text{Diag}\left(1, 1, \begin{pmatrix} \mu & \nu \\ \nu & \rho \end{pmatrix}\right)$	$\alpha \neq 0, \mu, \rho > 0$ $\mu\rho > \nu^2, x \neq 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_2] = xe_1 + ye_4,$ $[e_4, e_2] = ze_1 - ye_3$	αe^{12}	I_4	$\alpha y \neq 0$

TAB. 2. Four dimensional Riemann-Poisson Lie algebras of rank 2

Non vanishing Lie brackets	Bivector r	Matrix of ϱ	$\text{Der}^s(\mathfrak{h})$
$[e_1, e_2] = e_2,$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d)$	$\{E_{21}, E_{33} - E_{44}, E_{43}, E_{34}\}$
$[e_1, e_2] = -e_3, [e_1, e_3] = e_2,$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c)$	$\{E_{23} - E_{32}, E_{41}\}$
$[e_1, e_2] = e_2, [e_3, e_4] = e_4,$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d)$	$\{E_{21}, E_{43}\}$
$[e_4, e_1] = e_1, [e_4, e_2] = -\delta e_3,$ $[e_4, e_3] = \delta e_2$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c)$	$\{E_{14}, E_{23} - E_{32}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = \frac{1}{2}e_1, [e_4, e_2] = \frac{1}{2}e_2,$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, \mu b, \mu a, b)$	$\{E_{34}, E_{22} - E_{11}, E_{12} + E_{21}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = 2e_1, [e_4, e_2] = -e_2,$	$\alpha(e^{23} + e^{14})$	$\text{Diag}(a, a, 2a, 2a)$	$\{2E_{14} - E_{32}\}$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3,$ $[e_4, e_1] = \frac{1}{2}e_1 - e_2,$ $[e_4, e_2] = e_1 + \frac{1}{2}e_2,$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, a, a, a)$	$\{E_{34}, E_{12} - E_{21}\}$

TAB. 3. Four-dimensional Kähler Lie algebras and their symplectic derivations, $a, b, c, d > 0, \alpha\beta \neq 0$

• $\dim \mathfrak{g} = 5$. We have:

- (c51) $\dim \mathfrak{h} = 4$ and \mathfrak{h} abelian and hence a symplectic vector space. We can apply Proposition 4.2 and \mathfrak{g} is semi-direct product.
- (c52) $\dim \mathfrak{h} = 4$ and \mathfrak{h} non abelian. We can apply Proposition 4.2 and Table 3 to get the Lie brackets on \mathfrak{g}, ϱ and r . The result is summarized in Table 4.
- (c53) $\dim \mathfrak{h} = 2$ and \mathfrak{h} non abelian. We apply Proposition 4.3. In this case $(\mathfrak{p}, [\cdot, \cdot]_{\mathfrak{p}}, \varrho_{\mathfrak{p}})$ is a 3-dimensional Euclidean Lie algebra and one must compute $\text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$ and solve (20). Three dimensional Euclidean Lie algebras were classified in [9]. For each of them we have computed $\text{Der}(\mathfrak{p}) \cap \text{so}(\mathfrak{p})$

and solved (20) by using Maple. The result is summarized in Table 5 when \mathfrak{p} is unimodular and Table 6 when \mathfrak{p} is nonunimodular.

(c54) $\dim \mathfrak{h} = 2$ and \mathfrak{h} abelian and $\phi_{\mathfrak{h}} = 0$. We apply Proposition 4.6 and we perform all the needed computations. We use the classification of 3-dimensional Euclidean Lie algebras given in [9]. The results are given in Tables 7-8.

(c55) $\dim \mathfrak{h} = 2$ and \mathfrak{h} abelian and $\phi_{\mathfrak{h}} \neq 0$. We apply Proposition 4.7 and we perform all the needed computations. The results are given in Table 9.

Non vanishing Lie brackets	Bivector r	Matrix of ρ	Conditions
$[e_1, e_2] = e_2, [e_5, e_1] = xe_2,$ $[e_5, e_3] = ye_3 + te_4, [e_5, e_4] = ze_3 - ye_4$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d, e)$	$\alpha\beta \neq 0$ $a, b, c, d, e > 0$
$[e_1, e_2] = -e_3, [e_1, e_3] = e_2,$ $[e_5, e_1] = ye_4, [e_5, e_2] = -xe_3, [e_5, e_3] = xe_2$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c, d)$	$\alpha\beta \neq 0$ $a, b, c, d > 0$
$[e_1, e_2] = e_2, [e_3, e_4] = e_4,$ $[e_5, e_1] = xe_2, [e_5, e_3] = ye_4$	$\alpha e^{12} + \beta e^{34}$	$\text{Diag}(a, b, c, d, e)$	$\alpha\beta \neq 0$ $a, b, c, d, e > 0$
$[e_4, e_1] = e_1, [e_4, e_2] = -\delta e_3, [e_4, e_3] = \delta e_2$ $[e_5, e_2] = -ye_3, [e_5, e_3] = ye_2, [e_5, e_4] = xe_1$	$\alpha e^{14} + \beta e^{23}$	$\text{Diag}(a, b, b, c, d)$	$\alpha\beta \neq 0, \delta > 0$ $a, b, c, d > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = \frac{1}{2}e_1$ $[e_4, e_2] = \frac{1}{2}e_2, [e_5, e_1] = xe_1 + ye_2,$ $[e_5, e_2] = ye_1 - xe_2, [e_5, e_4] = ze_3$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, \mu b, \mu a, b, c)$	$\alpha \neq 0$ $a, b, c, \mu > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = 2e_1$ $[e_4, e_2] = -e_2, [e_5, e_2] = xe_3, [e_5, e_4] = -2xe_1$	$\alpha(e^{23} + e^{14})$	$\text{Diag}(a, a, 2a, 2a, b)$	$\alpha \neq 0$ $a, b > 0$
$[e_1, e_2] = e_3, [e_4, e_3] = e_3, [e_4, e_1] = \frac{1}{2}e_1 - e_2$ $[e_4, e_2] = e_1 + \frac{1}{2}e_2, [e_5, e_1] = -xe_2, [e_5, e_2] = xe_1$ $[e_5, e_4] = ye_3$	$\alpha(e^{12} - e^{34})$	$\text{Diag}(a, a, a, a, b)$	$\alpha \neq 0$ $a, b > 0$

TAB. 4. Five-dimensional Riemann-Poisson Lie algebras of rank 4

Non vanishing Lie brackets	r	Matrix of ρ	Conditions
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1 - ce_4, [e_4, e_2] = d\mu e_1 + ce_3$ $[e_5, e_2] = fe_1, [e_3, e_4] = -fe_1 + e_5$	αe^{12}	$\text{Diag}(1, \rho, \mu, \mu, 1)$	$c\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = ce_1$ $[e_5, e_2] = d\mu e_1, [e_3, e_5] = be_1 - e_2, [e_4, e_5] = -ce_1 + e_4$	αe^{12}	$\text{Diag}(1, \rho, 1, 1, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b+c)e_1, [e_4, e_2] = (cx+b)e_1$ $[e_5, e_2] = d\mu e_1, [e_3, e_5] = (b+c)e_1 - e_3,$ $[e_4, e_5] = -(xc+b)e_1 + e_4$	αe^{12}	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & 1 \\ 1 & x \end{pmatrix}, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = c\mu e_1$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = -\mu ce_1 + e_4, [e_4, e_5] = be_1 - e_3$	αe^{12}	$\text{Diag}(1, \rho, 1, \mu, \nu)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1$ $[e_5, e_2] = d\rho e_1, [e_3, e_4] = -2\rho de_1 + 2e_5,$ $[e_3, e_5] = 2\nu ce_1 - 2e_4, [e_4, e_5] = 2\mu be_1 - 2e_3$	αe^{12}	$\text{Diag}(1, \xi, \mu, \nu, \rho)$	$\alpha \neq 0, \nu \neq \rho$ $\mu, \nu, \rho, \xi > 0$ $\mu \neq \nu, \mu \neq \rho$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1 - \lambda e_5$ $[e_5, e_2] = d\nu e_1 + \lambda e_4, [e_3, e_4] = -\frac{2\nu(\lambda c+d)}{1+\lambda^2}e_1 + 2e_5,$ $[e_3, e_5] = \frac{2\nu(c-\lambda d)}{1+\lambda^2}e_1 - 2e_4, [e_4, e_5] = 2\mu be_1 - 2e_3$	αe^{12}	$\text{Diag}(1, \rho, \mu, \nu, \nu)$	$\lambda\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1$ $[e_5, e_2] = d\rho e_1, [e_3, e_4] = -\rho de_1 + e_5,$ $[e_3, e_5] = \nu ce_1 - e_4, [e_4, e_5] = -\mu be_1 + e_3$	αe^{12}	$\text{Diag}(1, \xi, \mu, \nu, \rho)$	$\alpha \neq 0, \nu \neq \rho$ $\mu, \nu, \rho, \xi > 0$ $\mu \neq \nu, \mu \neq \rho$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1, [e_4, e_2] = c\nu e_1 - \lambda e_5$ $[e_5, e_2] = d\nu e_1 + \lambda e_4, [e_3, e_4] = -\frac{\nu(\lambda c+d)}{1+\lambda^2}e_1 + e_5,$ $[e_3, e_5] = \frac{\nu(c-\lambda d)}{1+\lambda^2}e_1 - e_4, [e_4, e_5] = -\mu be_1 + e_3$	αe^{12}	$\text{Diag}(1, \rho, \mu, \nu, \nu)$	$\lambda\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = b\mu e_1 - ue_4 - ve_5,$ $[e_4, e_2] = c\mu e_1 + ue_3 - we_5, [e_5, e_2] = d\mu e_1 + ve_3 + we_4,$ $[e_3, e_4] = xe_1 + e_5, [e_3, e_5] = ye_1 - e_4, [e_4, e_5] = ze_1 + e_3$ $x = -\frac{\mu(buw-cv+du^2+bv+cw+d)}{1+u^2+v^2+w^2}$ $y = \frac{\mu(-bv+cv^2-duw+bu-dw+c)}{1+u^2+v^2+w^2}$ $z = -\frac{\mu(bw^2-cv+duw-cu-dv+b)}{1+u^2+v^2+w^2}$	αe^{12}	$\text{Diag}(1, \rho, \mu, \mu, \mu)$	$\alpha \neq 0$ $\mu, \rho > 0$

TAB. 5. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and unimodular complement

Non vanishing Lie brackets	r	Matrix of ϱ
$[e_1, e_2] = e_1, [e_3, e_2] = (f + c\lambda + f\lambda^2)e_1 - \lambda e_4, \lambda \neq 0,$ $[e_4, e_2] = ce_1 + \lambda e_3, [e_5, e_2] = d\mu e_1,$ $[e_3, e_5] = fe_1 - e_3, [e_4, e_5] = (\lambda f + c)e_1 - e_4,$	αe^{12}	$\text{Diag}(1, \rho, 1, 1, \mu)$ $\mu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = be_1, [e_4, e_2] = c\mu e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = \mu ce_1 - e_4,$ $[e_4, e_5] = (-fb + 2\mu c)e_1 + fe_3 - 2e_4, f = 1 \text{ or } f \leq 0$	αe^{12}	$\text{Diag}(1, \rho, 1, \mu, \nu)$ $0 < \mu < f , \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + c\mu)e_1, [e_4, e_2] = (c + b\mu)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (\mu b + c)e_1 - e_4,$ $[e_4, e_5] = ((2 - \mu)c + (2\mu - 1)b)e_1 + e_3 - 2e_4$	αe^{12}	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, \nu)$ $\mu, \nu, \rho > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + c)e_1, [e_4, e_2] = (b + c\mu)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (b + c\mu)e_1 - e_4,$ $[e_4, e_5] = ((2 - f)b + (2\mu - f)c)e_1 + fe_3 - 2e_4$	αe^{12}	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & 1 \\ 1 & \mu \end{pmatrix}, \nu)$ $\nu, \rho > 0, c > \mu > 1$
$[e_1, e_2] = e_1, [e_3, e_2] = (b + \frac{1}{2}c)e_1, [e_4, e_2] = (c + \frac{1}{2}b)e_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = (c + \frac{1}{2}b)e_1 - e_4,$ $[e_4, e_5] = (b + 2c)e_1 - 2e_4$	αe^{12}	$\text{Diag}(1, \rho, \begin{pmatrix} 1 & \frac{1}{2} \\ \frac{1}{2} & 1 \end{pmatrix}, \nu)$ $\rho, \nu > 0$
$[e_1, e_2] = e_1, [e_3, e_2] = xe_1, [e_4, e_2] = ye_1,$ $[e_5, e_2] = d\nu e_1, [e_3, e_5] = ze_1 - e_4,$ $[e_4, e_5] = te_1 + fe_3 - 2e_4, 0 < f < 1,$ $x = \frac{((\mu+1)b + (\mu-1)c)f - 2b}{2f^2(f-1)}, y = z = \frac{(\mu-1)(cf+b)}{2f(f-1)}$ $t = \frac{(1-\mu)cf + ((f-2)\mu + f)b}{2f(1-f)}$	αe^{12}	$A^t B A$ $A = \begin{pmatrix} \frac{1+s}{-2fs} & -\frac{1}{2s} & 0 \\ \frac{1-s}{2fs} & \frac{1}{2s} & 0 \\ 0 & 0 & 1 \end{pmatrix}$ $B = \text{Diag}(1, \rho, \begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, \nu)$ $\nu, \rho > 0$ $s = \sqrt{1-f}, 0 \leq \mu < 1$

TABLE 6. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with non abelian Kähler subalgebra and non unimodular complement. ($\alpha \neq 0$)

Non vanishing Lie brackets	Bivector r	Matrix of ϱ	Conditions
$[e_3, e_4] = ae_1 + be_2 + e_5, [e_3, e_5] = ce_1 + de_2$ $[e_4, e_5] = fe_1 + ge_2$	αe^{12}	$\text{Diag}(1, 1, \mu, \mu, 1)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_5] = ce_1 + de_2 - e_3$ $[e_4, e_5] = fe_1 + ge_2 + e_4$	αe^{12}	$\text{Diag}(1, 1, 1, 1, \mu)$ $\text{Diag}(1, 1, \begin{pmatrix} 1 & 1 \\ 1 & x \end{pmatrix}, \mu)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_4] = ae_1 + be_2, [e_3, e_5] = ce_1 + de_2 + e_4$ $[e_4, e_5] = fe_1 + ge_2 - e_3$	αe^{12}	$\text{Diag}(1, 1, 1, \mu, \nu)$	$\alpha \neq 0$ $\mu, \nu > 0$
$[e_3, e_4] = ae_1 + be_2 + 2e_5, [e_3, e_5] = ce_1 + de_2 - 2e_4$ $[e_4, e_5] = fe_1 + ge_2 - 2e_3$	αe^{12}	$\text{Diag}(1, 1, \mu, \nu, \rho)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_3, e_4] = ae_1 + be_2 + e_5, [e_3, e_5] = ce_1 + de_2 - e_4$ $[e_4, e_5] = fe_1 + ge_2 + e_3$	αe^{12}	$\text{Diag}(1, 1, \mu, \nu, \rho)$	$\alpha \neq 0$ $\mu, \nu, \rho > 0$
$[e_3, e_5] = ce_1 + de_2 - e_3$ $[e_4, e_5] = fe_1 + ge_2 - e_4$	αe^{12}	$\text{Diag}(1, 1, 1, 1, \mu)$	$\alpha \neq 0$ $\mu > 0$
$[e_3, e_5] = ce_1 + de_2 - e_4$ $[e_4, e_5] = fe_1 + ge_2 + xe_3 - 2e_4$	αe^{12}	There are many cases See [9]	$\alpha \neq 0$

TABLE 7. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra

Non vanishing Lie brackets	r	Matrix of ϱ
$[e_3, e_1] = -e_2, [e_3, e_2] = e_1, [e_4, e_1] = e_2, [e_4, e_2] = e_1$ $[e_5, e_1] = e_1, [e_5, e_2] = -e_2,$ $[e_3, e_4] = 2e_5 + (l_{22} - l_{21} - 2l_{13})e_1 - (l_{12} + l_{11} + 2l_{23})e_2$ $[e_3, e_5] = -2e_4 + (l_{23} - l_{11} + 2l_{12})e_1 - (l_{13} - l_{21} - 2l_{22})e_2,$ $[e_4, e_5] = -2e_3 + (l_{23} - l_{12} + 2l_{11})e_1 + (l_{13} + l_{22} + 2l_{21})e_2$	αe^{12}	Diag(1, 1, μ, ν, ρ) $\mu, \nu, \rho > 0$
$[e_4, e_2] = ue_1, [e_5, e_1] = -\frac{\alpha}{2}e_1, [e_5, e_2] = ve_1 + \frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $(a + 2b)x - 2tu + 2yv = 0, a \neq 0, b \neq 0, (3a + 2b)y = 0$	αe^{12}	Diag(1, 1, 1, 1, 1)
$[e_4, e_2] = ue_1, [e_5, e_1] = -\frac{\alpha}{2}e_1, [e_5, e_2] = ve_1 + \frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0,$ $ax - 2tu = 0$	αe^{12}	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_4, e_1] = ue_2, [e_5, e_1] = \frac{\alpha}{2}e_1 + ve_2, [e_5, e_2] = -\frac{\alpha}{2}e_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $(3a + 2b)x = 0, a \neq 0, b \neq 0$ $(a + 2b)y - 2zu + 2xv = 0$	αe^{12}	Diag(1, 1, 1, 1, 1)
$[e_4, e_1] = ue_2, [e_5, e_1] = \frac{\alpha}{2}e_1 + ve_2, [e_5, e_2] = -\frac{\alpha}{2}e_2,$ $[e_3, e_4] = ye_2, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0, ay - 2zu = 0$	αe^{12}	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_4, e_1] = ue_1 + upe_2, [e_4, e_2] = -\frac{u}{p}e_1 - ue_2,$ $[e_5, e_1] = ve_1 + \frac{(2v-a)p}{2}e_2, [e_5, e_2] = -\frac{(2v+a)}{2p}e_1 - ve_2$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = be_3 + ze_1 + te_2, a \neq 0, b \neq 0$ $[e_4, e_5] = ce_3 + ae_4 + re_1 + se_2,$ $((2a + 2b + 2v)x - 2zu)p - ay + 2tu - 2yv = 0$ $(2xv - ax - 2zu)p + (2a + 2b - 2v)y + 2tu = 0$	αe^{12}	Diag(1, 1, 1, 1, 1)
$[e_4, e_1] = ue_1 + upe_2, [e_4, e_2] = -\frac{u}{p}e_1 - ue_2,$ $[e_5, e_1] = ve_1 + \frac{(2v-a)p}{2}e_2, [e_5, e_2] = -\frac{(2v+a)}{2p}e_1 - ve_2$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = ze_1 + te_2,$ $[e_4, e_5] = ae_4 + re_1 + se_2, a \neq 0, b \neq 0$ $((2a + 2v)x - 2zu)p - ay + 2tu - 2yv = 0$ $(2xv - ax - 2zu)p + (2a - 2v)y + 2tu = 0$	αe^{12}	Diag(1, 1, $\begin{pmatrix} 1 & \mu \\ \mu & 1 \end{pmatrix}, 1)$
$[e_5, e_1] = ue_1 + ve_2, [e_5, e_2] = we_1 - ue_2,$ $[e_3, e_4] = xe_1 + ye_2, [e_3, e_5] = ae_3 + be_4 + ze_1 + te_2,$ $[e_4, e_5] = ce_3 + de_4 + re_1 + se_2,$ $(a + d + u)x + yw = 0, xv + (a + d - u)y = 0$	αe^{12}	Diag(1, 1, 1, 1, 1)
$[e_5, e_1] = ue_1 + ve_2, [e_5, e_2] = we_1 - ue_2,$ $[e_3, e_4] = xe_1 + ye_2 + ae_4, [e_3, e_5] = be_4 + ze_1 + te_2,$ $[e_4, e_5] = ce_4 + re_1 + se_2, a \neq 0$ $(c + u)x - ar + yw = 0$ $(c - u)y - as + xv = 0$	αe^{12}	Diag(1, 1, 1, 1, 1)

TAB. 8. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra ($\alpha \neq 0$) (Continued)

Non vanishing Lie brackets	Bivector r	Matrix of ϱ	Conditions
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3, [e_5, e_2] = ze_1$ $[e_3, e_5] = pe_3 + qe_4 + a^{-1}(-qx + py)e_1,$ $[e_3, e_5] = -qe_3 + pe_4 - a^{-1}(px + qy)e_1$	αe^{12}	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0$
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3, [e_5, e_2] = ze_1$ $[e_3, e_4] = be_1$ $[e_3, e_5] = qe_4 - a^{-1}qx e_1,$ $[e_3, e_5] = -qe_3 - a^{-1}qye_1$	αe^{12}	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0, z \neq 0$
$[e_3, e_2] = xe_1 - ae_4, [e_4, e_2] = ye_1 + ae_3,$ $[e_3, e_4] = be_1 + ce_2$ $[e_3, e_5] = qe_4 - a^{-1}qx e_1,$ $[e_3, e_5] = -qe_3 - a^{-1}qye_1$	αe^{12}	$\text{Diag}(1, 1, 1, 1, 1)$	$\alpha \neq 0$ $a \neq 0$

TAB. 9. Five-dimensional Riemann-Poisson Lie algebras of rank 2 with abelian Kähler subalgebra (Continued)

This theorem unknown to our knowledge can be used to build examples of Riemann-Poisson Lie algebras.

Theorem 4.1. *Let (G, \langle , \rangle) be an even dimensional flat Riemannian Lie group. Then there exists a left invariant differential Ω on G such that $(G, \langle , \rangle, \Omega)$ is a Kähler Lie group.*

Proof. Let \mathfrak{g} be the Lie algebra of G and $\varrho = \langle , \rangle(e)$. According to Milnor’s Theorem [11, Theorem 1.5] and its improved version [1, Theorem 3.1] the flatness of the metric on G is equivalent to $[\mathfrak{g}, \mathfrak{g}]$ is even dimensional abelian, $[\mathfrak{g}, \mathfrak{g}]^\perp = \{u \in \mathfrak{g}, \text{ad}_u + \text{ad}_u^* = 0\}$ is also even dimensional abelian and $\mathfrak{g} = [\mathfrak{g}, \mathfrak{g}] \oplus [\mathfrak{g}, \mathfrak{g}]^\perp$. Moreover, the Levi-Civita product is given by

$$(21) \quad L_a = \begin{cases} \text{ad}_a & \text{if } a \in [\mathfrak{g}, \mathfrak{g}]^\perp, \\ 0 & \text{if } a \in [\mathfrak{g}, \mathfrak{g}] \end{cases}$$

and there exists a basis $(e_1, f_1, \dots, e_r, f_r)$ of $[\mathfrak{g}, \mathfrak{g}]$ and $\lambda_1, \dots, \lambda_r \in [\mathfrak{g}, \mathfrak{g}]^\perp \setminus \{0\}$ such that for any $a \in [\mathfrak{g}, \mathfrak{g}]^\perp$,

$$[a, e_i] = \lambda_i(a) f_i \quad \text{and} \quad [a, f_i] = -\lambda_i(a) e_i.$$

We consider a nondegenerate skew-symmetric 2-form ω_0 on $[\mathfrak{g}, \mathfrak{g}]^\perp$ and ω_1 the nondegenerate skew-symmetric 2-form on $[\mathfrak{g}, \mathfrak{g}]$ given by $\omega_1 = \sum_{i=1}^r e_i^* \wedge f_i^*$. One can see easily that $\omega = \omega_0 \oplus \omega_1$ is a Kähler form on \mathfrak{g} . \square

REFERENCES

- [1] Ait Haddou, M., Boucetta, M., Lebzioui, H., *Left-invariant Lorentzian flat metrics on Lie groups*, J. Lie Theory **22** (1) (2012), 269–289.
- [2] Boucetta, M., *Compatibilité des structures pseudo-riemanniennes et des structures de Poisson*, C.R. Acad. Sci. Paris Sér. I **333** (2001), 763–768.
- [3] Boucetta, M., *Riemann-poisson manifolds and kähler-riemann foliations*, C.R. Acad. Sci. Paris, Sér. I **336** (2003), 423–428.
- [4] Boucetta, M., *Poisson manifolds with compatible pseudo-metric and pseudo-Riemannian Lie algebras*, Differential Geom. Appl. **20** (2004), 279–291.
- [5] Boucetta, M., *On the Riemann-Lie algebras and Riemann-Poisson Lie groups*, J. Lie Theory **15** (1) (2005), 183–195.

- [6] Deninger, C., Singhof, W., *Real polarizable hodge structures arising from foliation*, Ann. Global Anal. Geom. **21** (2002), 377–399.
- [7] Dufour, J.P., Zung, N.T., *Poisson Structures and Their Normal Forms*, Progress in Mathematics, vol. 242, Birkhäuser Verlag, 2005.
- [8] Fernandes, R.L., *Connections in Poisson Geometry 1: Holonomy and invariants*, J. Differential Geom. **54** (2000), 303–366.
- [9] Ha, K.Y., Lee, J.B., *Left invariant metrics and curvatures on simply connected three dimensional Lie groups*, Math. Nachr. **282** (2009), 868–898.
- [10] Hawkin, E., *The structure of noncommutative deformations*, J. Differential Geom. **77** (2007), 385–424.
- [11] Milnor, J., *Curvatures of left invariant metrics on Lie Groups*, Adv. Math. **21** (1976), 293–329.
- [12] Ovando, G., *Invariant pseudo-Kähler metrics in dimension four*, J. Lie Theory **16** (2006), 371–391.
- [13] Vaisman, I., *Lectures on the Geometry of Poisson Manifolds*, Progress in Mathematics, vol. 118, Birkhäuser, Berlin, 1994.

^aUNIVERSITÉ DE NOUAKCHOTT,
FACULTÉ DES SCIENCES ET TECHNIQUES
E-mail: pacha.ali86@gmail.com

^bUNIVERSITÉ CADI-AYYAD,
FACULTÉ DES SCIENCES ET TECHNIQUES,
BP 549 MARRAKECH MAROC
E-mail: m.boucetta@uca.ac.ma

^cUNIVERSITÉ DE NOUAKCHOTT,
FACULTÉ DES SCIENCES ET TECHNIQUES
E-mail: lessiadahmed@gmail.com