

Angkana Sripayap; Pattira Ruengsinsub; Teerapat Srichan  
On generalized square-full numbers in an arithmetic progression

*Czechoslovak Mathematical Journal*, Vol. 72 (2022), No. 1, 149–163

Persistent URL: <http://dml.cz/dmlcz/149578>

## Terms of use:

© Institute of Mathematics AS CR, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON GENERALIZED SQUARE-FULL NUMBERS  
IN AN ARITHMETIC PROGRESSION

ANGKANA SRIPAYAP, PATTIRA RUENGINSUB, TEERAPAT SRICHAN, Bangkok

Received August 20, 2020. Published online October 14, 2021.

*Abstract.* Let  $a$  and  $b \in \mathbb{N}$ . Denote by  $R_{a,b}$  the set of all integers  $n > 1$  whose canonical prime representation  $n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_r^{\alpha_r}$  has all exponents  $\alpha_i$  ( $1 \leq i \leq r$ ) being a multiple of  $a$  or belonging to the arithmetic progression  $at + b$ ,  $t \in \mathbb{N}_0 := \mathbb{N} \cup \{0\}$ . All integers in  $R_{a,b}$  are called generalized square-full integers. Using the exponent pair method, an upper bound for character sums over generalized square-full integers is derived. An application on the distribution of generalized square-full integers in an arithmetic progression is given.

*Keywords:* arithmetic progression; character sum; exponent pair method; square-full number

*MSC 2020:* 11B50, 11N25, 11N69

1. INTRODUCTION AND RESULTS

An integer  $n > 1$  is called *square-full*, if in the canonical representation of  $n$  each exponent is not less than 2. Let  $Q_2$  denote the set of all square-full integers,  $q_2(n)$  the characteristic function of the set  $Q_2$  and  $Q_2(x)$  the number of square-full integers  $n \leq x$ . The investigation of square-full integers has a long and rich history. Erdős and Szekeres in [5] proved that

$$(1.1) \quad Q_2(x) = \frac{\zeta(3/2)}{\zeta(3)} x^{1/2} + O(x^{1/3}).$$

In 1958, Bateman and Grosswald in [1] improved (1.1) and showed that

$$(1.2) \quad Q_2(x) = \frac{\zeta(3/2)}{\zeta(3)} x^{1/2} + \frac{\zeta(2/3)}{\zeta(2)} x^{1/3} + O(x^{1/6} \exp(-C(\log x)^{3/5} (\log \log x)^{-1/5}))$$

for some absolute constant  $C > 0$ . For a given integer  $q \geq 2$  and an integer  $l$  such that  $\gcd(l, q) = 1$ ,  $Q_2(x; l, q)$  denotes the number of square-full numbers  $n \leq x$  and

$n \equiv l \pmod{q}$ . The study on the distribution of square-full numbers in an arithmetic progression attracts the interests of many authors; see, e.g. [2], [3], [6], [7] and [10].

It would be interesting to extend this problem on generalized square-full integers. Let  $a$  and  $b$  be fixed positive integers and  $R_{a,b}$  the set of all integers  $n > 1$  with the property that, for the canonical representation of  $n = p_1^{\alpha_1} p_2^{\alpha_2} \dots p_r^{\alpha_r}$ , each exponent  $\alpha_i$  ( $1 \leq i \leq r$ ) is either a multiple of  $a$  or is contained in the progression  $at + b$ ,  $t \geq 0$ . Let  $r_{a,b}$  be the characteristic function for the element in the set  $R_{a,b}$  and  $R_{a,b}(x)$  the number of integers  $n \leq x$  such that  $n \in R_{a,b}$ . Cohen was the first mathematician to introduce generalized square-full integers and he derived asymptotic formulas for  $R_{a,b}(x)$  in various cases, see [4]. It is clear that the set  $R_{2,3}$  is the set of square-full integers. Thus, all integers in  $R_{a,b}$  may be called *generalized square-full integers*.

In this paper, we use the exponent pair method of [8] and [9] to bound the character sums over generalized square-full numbers and apply it to study the distribution for a generalized square-full numbers in an arithmetic progression.

In [10], Srichan used the exponent pair method to study the appearance of square-full numbers in an arithmetic progression and showed that

$$(1.3) \quad Q_2(x; l, q) = \frac{L(3/2, \chi_0) + \sum_{\chi_1} \bar{\chi}_1(l) L(3/2, \chi_1)}{qL(3, \chi_0)} x^{1/2} \\ + \frac{L(2/3, \chi_0) + \sum_{\chi_2} \bar{\chi}_2(l) L(2/3, \chi_2)}{qL(2, \chi_0)} x^{1/3} + O(q^{1/2+\varepsilon} x^{1/6}),$$

where  $\chi_0$ ,  $\chi_1$  and  $\chi_2$  denote the principal, quadratic and cubic character modulo  $q$ , respectively. However, there is a mistake in the  $O$ -term of (1.3). Srichan overlooked the value of  $\varphi(q)$ . In particular, the  $O$ -term in the formula for  $Q_2(x; l, q)$  should be corrected to  $q^{4/3+\varepsilon} x^{1/6}$ . In the present paper, we also give this correction.

Let  $a$  and  $b$  be fixed positive integers, and  $a < b < 2a$ . For given integers  $q \geq 2$  and  $0 < l < q$  such that  $\gcd(l, q) = 1$ , let  $R_{a,b}(x; l, q)$  be the number of integers  $n \leq x$  such that  $n \in R_{a,b}$  and  $n \equiv l \pmod{q}$ . We obtain the following theorem.

**Theorem 1.1.** *Let  $a$  and  $b$  be fixed positive integers and  $a < b < 2a$ . For given integers  $q \geq 2$ ,  $0 < l < q$ , such that  $\gcd(l, q) = 1$  and  $x \geq q^{a+b}$ , we have*

$$R_{a,b}(x; l, q) = \left( \frac{L(b/a, \chi_0)}{qL(2b/a, \chi_0)} + \sum_{\chi \in A_\chi} \bar{\chi}(l) \frac{L(b/a, \chi^b)}{qL(2b/a, \chi^{2b})} \right) x^{1/a} \\ + \left( \frac{L(a/b, \chi_0)}{qL(2, \chi_0)} + \sum_{\chi \in B_\chi} \bar{\chi}(l) \frac{L(a/b, \chi^a)}{qL(2, \chi^{2b})} \right) x^{1/b} + O(q^{4/3+\varepsilon} x^{1/(2b)}),$$

where  $A_\chi$  and  $B_\chi$  denote the set of all non-principal characters modulo  $q$  of order  $d$  such that  $d \mid a$  and  $d \mid b$ , respectively.

**Remark 1.1.** Using Theorem 1.1 we obtain the correct error term in [10] for  $a = 2$  and  $b = 3$ .

## 2. PRELIMINARY RESULTS AND NOTATIONS

As usual, let  $\mu(n)$  and  $\varphi(n)$  denote the Möbius function and the Euler phi function, respectively, and  $\tau_{a,b}(n)$  denotes the number of ordered pairs  $(d, \delta)$  of positive integers  $d$  and  $\delta$  such that  $d^a \delta^b = n$ . Let  $\psi(x) = x - \lfloor x \rfloor - \frac{1}{2}$ . The following lemmas are well known and the proofs can be found in their references.

**Lemma 2.1** ([4], Lemma 2.1). *If  $a \nmid b$ , then*

$$r_{a,b}(n) = \sum_{d^{2b}\delta=n} \mu(d)\tau_{a,b}(\delta).$$

**Lemma 2.2** ([8], Lemma 2). *Let  $\omega$  and  $\kappa$  be two real numbers,  $\omega > 0$ ,  $0 < \kappa \neq 1$ . Then*

$$\sum_{n \leq \omega} n^{-\kappa} = \zeta(\kappa) - \frac{1}{\kappa - 1} \omega^{1-\kappa} - \psi(\omega) \omega^{-\kappa} + O(\omega^{-\kappa-1}).$$

**Lemma 2.3.** *For real  $x > 1$  and any non-principal character  $\chi$  modulo  $q$ ,*

$$(2.1) \quad \sum_{k \leq x} \chi(k) = \sum_{j \leq q} \chi(j) \left[ \frac{x}{q} - \frac{j}{q} + 1 \right]$$

and

$$(2.2) \quad \sum_{k \leq x} \chi(k) [k] = \sum_{j \leq q} \chi(j) \sum_{\substack{k \leq x \\ k \equiv j \pmod{q}}} [k].$$

*Proof.* Equations (2.1) and (2.2) follow from the periodicity of the character  $\chi$  modulo  $q$ . □

**Lemma 2.4** ([9], Lemma 13). *If  $f(n)$  is an arithmetic function, then*

$$\sum_{\substack{n \leq \omega, \\ \gcd(n,q)=1}} f(n) = \sum_{d|q} \mu(d) \sum_{m \leq \omega/d} f(md).$$

**Lemma 2.5.** For  $\alpha > 0$ ,  $\alpha \neq 1$ , and  $0 < \beta \leq 1$ , we have

$$\sum_{\substack{n \leq x \\ n \equiv l \pmod{q}}} n^{-\alpha} = q^{-\alpha} \zeta\left(\alpha, \frac{l}{q}\right) + \frac{1}{1-\alpha} \frac{x^{1-\alpha}}{q} - \psi\left(\frac{x-l}{q}\right) x^{-\alpha} + O(qx^{-\alpha-1}),$$

where

$$\zeta(s, \beta) = \sum_{n=0}^{\infty} (n + \beta)^{-s}, \quad \Re(s) > 1.$$

*Proof.* It follows from Lemma 14 in [9]. □

**Lemma 2.6** ([9], Lemma 17). Let  $x, \eta, \alpha, \omega$  be real numbers,  $j$  and  $q$  positive numbers, where  $x \geq 1$ ,  $\alpha > 0$ ,  $\eta \geq 1$ ,  $1 \leq j \leq q$ , and  $(k, l)$  an exponent pair with  $k > 0$  and

$$R(x, \eta, \alpha; q, j; \omega) = \sum_{\substack{n \leq \eta \\ n \equiv j \pmod{q}}} \psi\left(\frac{x}{n^\alpha} + \omega\right),$$

if  $\omega$  is independent on  $n$ . Then

$$R(x, \eta, \alpha; q, j; \omega) = O(1) + O(x^{-1/2} \eta^{1+\alpha/2} q^{-1}) + \begin{cases} O(x^{k/(k+1)} \eta^{(l-\alpha k)/(k+1)} q^{-l/(k+1)}) & \text{for } l > \alpha k, \\ O(x^{k/(k+1)} \log \eta q^{-\alpha k/(k+1)}) & \text{for } l = \alpha k, \\ O((xq^{-\alpha})^{k/(1+(1+\alpha)k-l)}) & \text{for } l < \alpha k, \end{cases}$$

where the  $O$ -constants are dependent on only  $\alpha$ .

### 3. PROOF OF THEOREM 1.1

*Proof.* Let  $a$  and  $b$  be fixed positive integers and  $a < b < 2a$ . Let  $q$  and  $l$  be integers such that  $q \geq 2$ ,  $0 < l < q$  and  $\gcd(l, q) = 1$ . By the orthogonality relation for Dirichlet characters modulo  $q$ , we have

$$(3.1) \quad R_{a,b}(x; l, q) = \sum_{\substack{n \leq x \\ n \equiv l \pmod{q}}} r_{a,b}(n) = \frac{1}{\varphi(q)} \sum_{\chi} \bar{\chi}(l) C_{a,b}(x, \chi),$$

where

$$C_{a,b}(x, \chi) := \sum_{n \leq x} r_{a,b}(n) \chi(n).$$

In view of Lemma 2.1, we have

$$\begin{aligned}
(3.2) \quad C_{a,b}(x, \chi) &= \sum_{d^{2b} \delta \leq x} \mu(d) \chi^{2b}(d) \tau_{a,b}(\delta) \chi(\delta) \\
&= \sum_{d \leq (xq^{-(a+b)})^{1/(2b)}} \mu(d) \chi^{2b}(d) \sum_{\delta \leq xd^{-2b}} \tau_{a,b}(\delta) \chi(\delta) \\
&\quad + \sum_{(xq^{-(a+b)})^{1/(2b)} < d \leq x^{1/(2b)}} \mu(d) \chi^{2b}(d) \sum_{\delta \leq xd^{-2b}} \tau_{a,b}(\delta) \chi(\delta) \\
&:= U_1 + U_2.
\end{aligned}$$

First, we bound  $U_2$ . For any character modulo  $q$ , we have

$$\begin{aligned}
(3.3) \quad U_2 &= \sum_{(xq^{-(a+b)})^{1/(2b)} < d \leq x^{1/(2b)}} \mu(d) \chi^{2b}(d) \sum_{m^a n^b \leq xd^{-2b}} \chi(m^a n^b) \\
&\ll \sum_{(xq^{-(a+b)})^{1/(2b)} < d \leq x^{1/(2b)}} \frac{1}{q} (xd^{-2b})^{1/a} \\
&\ll x^{1/(2b)} q^{-(a^2+ab-2b^2)/(2ab)}.
\end{aligned}$$

Now we calculate  $U_1$ . Put

$$T_{a,b}(x, \chi) := \sum_{n \leq x} \tau_{a,b}(n) \chi(n).$$

Thus

$$(3.4) \quad U_1 = \sum_{d \leq (xq^{-(a+b)})^{1/(2b)}} \mu(d) \chi^{2b}(d) T_{a,b}(xd^{-2b}, \chi).$$

Then we need the asymptotic formula for  $T_{a,b}(x, \chi)$ . We consider the corresponding Dirichlet series of  $r_{a,b}(n) \chi(n)$ , namely,

$$\sum_{n=1}^{\infty} \frac{r_{a,b}(n) \chi(n)}{n^s} = \frac{L(as, \chi^a) L(bs, \chi^b)}{L(2bs, \chi^{2b})},$$

which converges absolutely for  $\Re(s) > 1/a$ . Since

$$\frac{1}{L(2bs, \chi^{2b})} = \sum_{n=1}^{\infty} \frac{\mu(n) \chi^{2b}(n)}{n^{2bs}} \quad \text{for } \Re(s) \geq \frac{1}{2b}$$

has a comparatively small abscissa of absolute convergence with  $1 \leq r < 2b$ , we put

$$D_b(x, \chi, r) := \sum_{d \leq x^{1/(2b)}} \frac{\mu(d) \chi^{2b}(d)}{d^{2b/r}}.$$

Thus, for  $1 \leq r < 2b$ , we have

$$(3.5) \quad D_b(x, \chi, r) = \frac{1}{L(2bs, \chi^{2b})} - \sum_{d > x^{1/(2b)}} \frac{\mu(d)\chi^{2b}(d)}{d^{2b/r}} = \frac{1}{L(2bs, \chi^{2b})} + O(x^{1/(2b)-1/r}).$$

Since

$$\sum_{n=1}^{\infty} \frac{\tau_{a,b}(n)\chi(n)}{n^s} = L(as, \chi^a)L(bs, \chi^b)$$

and from the Euler product for  $L(s, \chi)$ , we can deduce

$$(3.6) \quad T_{a,b}(x, \chi) = \sum_{m^a n^b \leq x} \chi^a(m)\chi^b(n) = \sum_{m \leq x^{1/(a+b)}} \chi^a(m) \sum_{n \leq (x/m^a)^{1/b}} \chi^b(n) \\ + \sum_{n \leq x^{1/(a+b)}} \chi^b(n) \sum_{m \leq (x/n^b)^{1/a}} \chi^a(m) \\ - \sum_{m \leq x^{1/(a+b)}} \chi^a(m) \sum_{n \leq x^{1/(a+b)}} \chi^b(n).$$

Let  $A_\chi$  and  $B_\chi$  be the set of all non-principal characters mod  $q$  of order  $d$  such that  $d \mid a$  and  $d \mid b$ , respectively. Now we consider the sum (3.6) in the following five cases:

*Case 1:*  $\chi = \chi_0$ , the principal character modulo  $q$ . In view of (3.6), we have

$$T_{a,b}(x, \chi_0) = \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} \sum_{\substack{n \leq (x/m^a)^{1/b} \\ \gcd(n,q)=1}} 1 + \sum_{\substack{n \leq x^{1/(a+b)} \\ \gcd(n,q)=1}} \sum_{\substack{m \leq (x/n^b)^{1/a} \\ \gcd(m,q)=1}} 1 \\ - \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} 1 \sum_{\substack{n \leq x^{1/(a+b)} \\ \gcd(n,q)=1}} 1.$$

In view of Lemma 2.4, we have

$$T_{a,b}(x, \chi_0) = \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/b}/(dm^{a/b}) \\ \gcd(n,q)=1}} 1 + \sum_{\substack{n \leq x^{1/(a+b)} \\ \gcd(n,q)=1}} \sum_{d|q} \mu(d) \sum_{\substack{m \leq x^{1/a}/(dn^{b/a}) \\ \gcd(m,q)=1}} 1 \\ - \sum_{d|q} \mu(d) \sum_{\substack{m \leq x^{1/(a+b)}/d \\ \gcd(m,q)=1}} 1 \sum_{t|q} \mu(t) \sum_{\substack{m \leq x^{1/(a+b)}/t \\ \gcd(m,q)=1}} 1 \\ = \sum_{d|q} \mu(d) \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} \left\lfloor \frac{x^{1/b}}{dm^{a/b}} \right\rfloor + \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ \gcd(n,q)=1}} \left\lfloor \frac{x^{1/a}}{dn^{b/a}} \right\rfloor \\ - \sum_{d|q} \mu(d) \left\lfloor \frac{x^{1/(a+b)}}{d} \right\rfloor \sum_{t|q} \mu(t) \left\lfloor \frac{x^{1/(a+b)}}{t} \right\rfloor.$$

We apply Lemma 2.4 again to the first two inner sums and note that the last term is the product of the same sums. Thus, we have

$$\begin{aligned}
T_{a,b}(x, \chi_0) &= \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \left\lfloor \frac{x^{1/b}}{d(tm)^{a/b}} \right\rfloor \\
&+ \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \left\lfloor \frac{x^{1/a}}{d(tm)^{b/a}} \right\rfloor \\
&- \left( \sum_{d|q} \mu(d) \left\lfloor \frac{x^{1/(a+b)}}{d} \right\rfloor \right)^2.
\end{aligned}$$

From  $\lfloor x \rfloor = x - \psi(x) - \frac{1}{2}$ , we have

$$\begin{aligned}
T_{a,b}(x, \chi_0) &= \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \left( \frac{x^{1/b}}{d(tm)^{a/b}} - \psi\left(\frac{x^{1/b}}{d(tm)^{a/b}}\right) - \frac{1}{2} \right) \\
&+ \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \left( \frac{x^{1/a}}{d(tm)^{b/a}} - \psi\left(\frac{x^{1/a}}{d(tm)^{b/a}}\right) - \frac{1}{2} \right) \\
&- \left( \sum_{d|q} \mu(d) \left( \frac{x^{1/(a+b)}}{d} - \psi\left(\frac{x^{1/(a+b)}}{d}\right) - \frac{1}{2} \right) \right)^2.
\end{aligned}$$

In view of the well known identities  $\sum_{d|q} \mu(d) = 0$  for  $q > 1$  and  $\sum_{d|q} \mu(d)/d = \varphi(q)/q$ , we have

$$\begin{aligned}
T_{a,b}(x, \chi_0) &= x^{1/b} \frac{\varphi(q)}{q} \sum_{t|q} \frac{\mu(t)}{t^{a/b}} \sum_{m \leq t^{-1}x^{1/(a+b)}} m^{-a/b} \\
&- \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq t^{-1}x^{1/(a+b)}} \psi\left(\frac{x^{1/b}}{d(tm)^{a/b}}\right) \\
&+ x^{1/a} \frac{\varphi(q)}{q} \sum_{t|q} \frac{\mu(t)}{t^{b/a}} \sum_{m \leq t^{-1}x^{1/(a+b)}} m^{-b/a} \\
&- \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq t^{-1}x^{1/(a+b)}} \psi\left(\frac{x^{1/a}}{d(tm)^{b/a}}\right) \\
&+ 2 \frac{\varphi(q)}{q} \sum_{d|q} \mu(d) \psi\left(\frac{x^{1/(a+b)}}{d}\right) x^{1/(a+b)} \\
&- \frac{\varphi^2(q)}{q^2} x^{2/(a+b)} + O(q^\varepsilon).
\end{aligned}$$



Now we use Lemma 2.2 to get

$$\begin{aligned}
T_{a,b}(x, \chi_0) &= x^{1/b} \frac{\varphi(q)}{q} \sum_{t|q} \frac{\mu(t)}{t^{a/b}} \left( \zeta\left(\frac{a}{b}\right) + b \frac{x^{1/b(a+b)}}{t^{1/b}} \right. \\
&\quad \left. - \psi\left(\frac{x^{1/(a+b)}}{t}\right) x^{-a/(a+b)b} t^{a/b} + O\left(x^{-1/b} t^{(a+b)/b}\right) \right) \\
&\quad - \sum_{d|q} \sum_{t|q} \mu(d) \mu(t) \sum_{m \leq t^{-1} x^{1/(a+b)}} \psi\left(\frac{x^{1/b}}{d(tm)^{a/b}}\right) \\
&\quad + x^{1/a} \frac{\varphi(q)}{q} \sum_{t|q} \frac{\mu(t)}{t^{b/a}} \left( \zeta\left(\frac{b}{a}\right) - a \frac{x^{-1/a(a+b)}}{t^{-1/a}} \right. \\
&\quad \left. - \psi\left(\frac{x^{1/(a+b)}}{t}\right) x^{-b/a(a+b)} t^{b/a} + O\left(x^{-1/a} t^{(a+b)/a}\right) \right) \\
&\quad - \sum_{d|q} \sum_{t|q} \mu(d) \mu(t) \sum_{m \leq t^{-1} x^{1/(a+b)}} \psi\left(\frac{x^{1/a}}{d(tm)^{b/a}}\right) \\
&\quad - \frac{\varphi^2(q)}{q^2} x^{2/(a+b)} + 2 \frac{\varphi(q)}{q} \sum_{d|q} \mu(d) \psi\left(\frac{x^{1/(a+b)}}{d}\right) x^{1/(a+b)} + O(q^\varepsilon).
\end{aligned}$$

In view of  $\sum_{t|q} \mu(t)/t^\alpha = \prod_{p|q} (1 - p^{-\alpha})$ ,  $\alpha \in \mathbb{R}$ , and  $\zeta(s) \prod_{p|q} (1 - p^{-s}) = L(s, \chi_0)$ ,  $s \neq 1$ , we have

$$\begin{aligned}
(3.7) \quad T_{a,b}(x, \chi_0) &= \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} + b \frac{\varphi^2(q)}{q^2} x^{2/(a+b)} \\
&\quad - \frac{\varphi(q)}{q} \sum_{t|q} \mu(t) \psi\left(\frac{x^{1/(a+b)}}{t}\right) x^{1/(a+b)} + O(q^{1+\varepsilon}) \\
&\quad - \sum_{d|q} \sum_{t|q} \mu(d) \mu(t) \sum_{m \leq t^{-1} x^{1/(a+b)}} \psi\left(\frac{x^{1/b}}{d(tm)^{a/b}}\right) \\
&\quad + \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} - a \frac{\varphi^2(q)}{q^2} x^{2/(a+b)} \\
&\quad - \frac{\varphi(q)}{q} \sum_{t|q} \mu(t) \psi\left(\frac{x^{1/(a+b)}}{t}\right) x^{1/(a+b)} + O(q^{1+\varepsilon}) \\
&\quad - \sum_{d|q} \sum_{t|q} \mu(d) \mu(t) \sum_{m \leq t^{-1} x^{1/(a+b)}} \psi\left(\frac{x^{1/a}}{d(tm)^{b/a}}\right) - \frac{\varphi^2(q)}{q^2} x^{2/(a+b)} \\
&\quad + 2 \frac{\varphi(q)}{q} \sum_{d|q} \mu(d) \psi\left(\frac{x^{1/(a+b)}}{d}\right) x^{1/(a+b)} + O(q^\varepsilon) \\
&:= \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} + \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} + O(q^{1+\varepsilon}) - S_1 - S_2,
\end{aligned}$$

where

$$S_1 = \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \psi\left(\frac{x^{1/b}}{d(tm)^{a/b}}\right)$$

and

$$S_2 = \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) \sum_{m \leq x^{1/(a+b)}/t} \psi\left(\frac{x^{1/a}}{d(tm)^{b/a}}\right).$$

If we use Lemma 2.6 with the exponent pair  $(\frac{2}{7}, \frac{4}{7})$ , then for  $q \leq x^{1/(a+b)}$  we have

$$\begin{aligned} S_1 &= \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) R\left(\frac{x^{1/b}}{dt^{a/b}}, \frac{x^{1/(a+b)}}{t}, \frac{a}{b}, 1, 0, 0\right) \\ &\ll \sum_{d|q} \sum_{t|q} (x^{1/(2a+2b)} d^{1/2} t^{-1} + x^{2/(3a+3b)} d^{-2/9} t^{-4/9}) \\ &= O(x^{2/(3a+3b)} q^{1/3+\varepsilon}) \end{aligned}$$

and

$$\begin{aligned} S_2 &= \sum_{d|q} \sum_{t|q} \mu(d)\mu(t) R\left(\frac{x^{1/a}}{dt^{b/a}}, \frac{x^{1/(a+b)}}{t}, \frac{b}{a}, 1, 0, 0\right) \\ &\ll \sum_{d|q} \sum_{t|q} (x^{1/(2a+2b)} d^{1/2} t^{-1} + x^{2/(3a+3b)} d^{-2/9} t^{-4/9}) \\ &= O(x^{2/(a+b)} q^{1/3+\varepsilon}). \end{aligned}$$

Substituting for  $S_1$  and  $S_2$  in (3.7), we have, for  $q \leq x^{1/(a+b)}$ ,

$$(3.8) \quad T_{a,b}(x, \chi_0) = \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} + \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} + O(x^{2/(3a+3b)} q^{1/3+\varepsilon}).$$

*Case 2:* the non-principal characters  $\chi$  modulo  $q$  such that  $\chi \in A_\chi - B_\chi$ . In view of (3.6), we have

$$\begin{aligned} T_{a,b}(x, \chi) &= \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} \sum_{n \leq (x/m^a)^{1/b}} \chi^b(n) \\ &\quad + \sum_{n \leq x^{1/(a+b)}} \chi^b(n) \sum_{\substack{m \leq (x/n^b)^{1/a} \\ \gcd(m,q)=1}} 1 \\ &\quad - \sum_{\substack{m \leq x^{1/(a+b)} \\ \gcd(m,q)=1}} 1 \sum_{n \leq x^{1/(a+b)}} \chi^b(n). \end{aligned}$$

We use (2.1) of Lemma 2.3 for the inner sum of the first term and then apply Lemma 2.1. For the second term we first use Lemma 2.1 for its inner sum and then

use (2.2) of Lemma 2.3. We apply Lemmas 2.1 and 2.3 separately for the last term. Thus, we have

$$\begin{aligned}
T_{a,b}(x, \chi) &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{n \leq x^{1/(a+b)}/d} \left[ \frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q} + 1 \right] \\
&\quad + \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} \left[ \frac{x^{1/a}}{dn^{b/a}} \right] \\
&\quad - \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \left[ \frac{x^{1/(a+b)}}{d} \right] \left[ \frac{x^{1/(a+b)}}{q} - \frac{j}{q} + 1 \right].
\end{aligned}$$

We write  $[x]$  as  $x - \psi(x) - \frac{1}{2}$  and  $\psi(x+1) = \psi(x)$ . We obtain

$$\begin{aligned}
T_{a,b}(x, \chi) &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{n \leq x^{1/(a+b)}/d} \left( \frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q} + \frac{1}{2} - \psi\left(\frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q}\right) \right) \\
&\quad + \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} \left( \frac{x^{1/a}}{dn^{b/a}} - \psi\left(\frac{x^{1/a}}{dn^{b/a}}\right) - \frac{1}{2} \right) \\
&\quad - \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \left[ \frac{x^{1/(a+b)}}{d} \right] \left( \frac{x^{1/(a+b)}}{q} - \frac{j}{q} + \frac{1}{2} - \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) \right).
\end{aligned}$$

Using the following identities,  $\sum_{j \leq q} \chi(j) = 0$  for a non-principal character  $\chi$ , and  $\sum_{d|q} \mu(d) = 0$ ,  $q > 1$ , we have

$$\begin{aligned}
(3.9) \quad T_{a,b}(x, \chi) &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{n \leq x^{1/(a+b)}/d} \left( -\frac{j}{q} - \psi\left(\frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q}\right) \right) \\
&\quad + \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} \left( \frac{x^{1/a}}{dn^{b/a}} - \psi\left(\frac{x^{1/a}}{dn^{b/a}}\right) \right) \\
&\quad - \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \left[ \frac{x^{1/(a+b)}}{d} \right] \left( -\frac{j}{q} - \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) \right) \\
&= - \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{n \leq x^{1/(a+b)}/d} \psi\left(\frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q}\right) \\
&\quad + \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} \left( \frac{x^{1/a}}{dn^{b/a}} - \psi\left(\frac{x^{1/a}}{dn^{b/a}}\right) \right) \\
&\quad + \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \left[ \frac{x^{1/(a+b)}}{d} \right] \left( \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) \right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{\varphi(q)}{q} x^{1/a} \sum_{j \leq q} \chi^b(j) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} n^{-b/a} \\
&\quad + \frac{\varphi(q)}{q} x^{1/(a+b)} \sum_{j \leq q} \chi^b(j) \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) + O(q^2) - S_3 - S_4,
\end{aligned}$$

where

$$\begin{aligned}
S_3 &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{n \leq x^{1/(a+b)}/d} \psi\left(\frac{x^{1/b}}{q(nd)^{a/b}} - \frac{j}{q}\right), \\
S_4 &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} \psi\left(\frac{x^{1/a}}{dn^{b/a}}\right).
\end{aligned}$$

Using Lemma 2.5 and writing the Hurwitz zeta-function as the Dirichlet  $L$ -function  $q^{-s} \sum_{j \leq q} \chi(j) \zeta(s, j/q) = L(s, \chi)$  for  $s \neq 1$ , we have

$$\begin{aligned}
(3.10) \quad &x^{1/a} \frac{\varphi(q)}{q} \sum_{j \leq q} \chi(j)^b \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv j \pmod{q}}} n^{-b/a} \\
&= x^{1/a} \frac{\varphi(q)}{q} \sum_{j \leq q} \chi^b(j) \left( q^{-b/a} \zeta\left(\frac{b}{a}, \frac{j}{q}\right) + \frac{a}{a-b} x^{(a-b)/a} \right. \\
&\quad \left. - \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) x^{-b/(a(a+b))} + O(qx^{-1/a}) \right) \\
&= x^{1/a} \frac{\varphi(q)}{q} \sum_{j \leq q} \chi^b(j) \left( q^{-b/a} \zeta\left(\frac{b}{a}, \frac{j}{q}\right) \right. \\
&\quad \left. - \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) x^{-b/(a(a+b))} + O(qx^{-1/a}) \right) \\
&= -\frac{\varphi(q)}{q} \sum_{j \leq q} \chi^b(j) \psi\left(\frac{x^{1/(a+b)}}{q} - \frac{j}{q}\right) x^{1/(a+b)} \\
&\quad + \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi^b\right) x^{1/a} + O(q^2).
\end{aligned}$$

Substituting (3.10) into (3.9), we get

$$(3.11) \quad T_{a,b}(x, \chi) = \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi^b\right) x^{1/a} + O(q^2) - S_3 - S_4.$$

If we use Lemma 2.6 with the exponent pair  $(\frac{2}{7}, \frac{4}{7})$ , then for  $q \leq x^{1/(a+b)}$  we have

$$\begin{aligned} S_3 &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) R\left(\frac{x^{1/b}}{qd^{a/b}}, \frac{x^{1/(a+b)}}{d}, \frac{a}{b}, 1, 1, \frac{-j}{q}\right) \\ &\ll \sum_{j \leq q} \sum_{d|q} \left(x^{1/(2a+2b)} q^{1/2} d^{-1} + x^{2/(3a+3b)} q^{-2/9} d^{-4/9}\right) \\ &= O(x^{2/(3a+3b)} q^{4/3+\varepsilon}) \end{aligned}$$

and similarly

$$\begin{aligned} S_4 &= \sum_{j \leq q} \chi^b(j) \sum_{d|q} \mu(d) R\left(\frac{x^{1/a}}{d}, x^{1/(a+b)}, \frac{b}{a}, q, j, 0\right) \\ &\ll \sum_{j \leq q} \sum_{d|q} \left(x^{1/(2a+2b)} q^{1/2} d^{-1} + x^{2/(3a+3b)} q^{-2/9} d^{-4/9}\right) \\ &= O(x^{2/(3a+3b)} q^{4/3+\varepsilon}). \end{aligned}$$

Replacing  $S_3$  and  $S_4$  in (3.11), we have, for  $q \leq x^{1/(a+b)}$ ,

$$(3.12) \quad T_{a,b}(x, \chi) = \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi^b\right) x^{1/a} + O(x^{2/(3a+3b)} q^{4/3+\varepsilon}).$$

*Case 3:* non-principal characters  $\chi$  modulo  $q$  such that  $\chi \in B_\chi - A_\chi$ . We prove this case by the same steps as Case 2. We obtain, for  $q \leq x^{1/(a+b)}$ ,

$$(3.13) \quad T_{a,b}(x, \chi) = \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi^a\right) x^{1/b} + O(x^{2/(3a+3b)} q^{4/3+\varepsilon}).$$

*Case 4:* non-principal characters  $\chi$  modulo  $q$  such that  $\chi \in A_\chi \cap B_\chi$ . In this case we obtain the same result as in Case 1. Thus we have, for  $q \leq x^{1/(a+b)}$ ,

$$(3.14) \quad T_{a,b}(x, \chi) = \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} + \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} + O(x^{2/(3a+3b)} q^{1/3+\varepsilon}).$$

*Case 5:* non-principal characters  $\chi$  modulo  $q$  such that  $\chi \notin A_\chi \cup B_\chi$ . In view of (3.6), we use (2.1) of Lemma 2.3 and get that

$$\begin{aligned} T_{a,b}(x, \chi) &= \sum_{j \leq q} \chi^b(j) \sum_{n \leq x^{1/(a+b)}} \chi^a(n) \left[ \frac{x^{1/b}}{qn^{a/b}} - \frac{j}{q} + 1 \right] \\ &\quad + \sum_{j \leq q} \chi^a(j) \sum_{m \leq x^{1/(a+b)}} \chi^b(m) \left[ \frac{x^{1/a}}{qm^{b/a}} - \frac{j}{q} + 1 \right] \\ &\quad - \sum_{j \leq q} \sum_{h \leq q} \chi^a(j) \chi^b(h) \left[ \frac{x^{1/(a+b)}}{q} - \frac{j}{q} + 1 \right] \left[ \frac{x^{1/(a+b)}}{q} - \frac{h}{q} + 1 \right]. \end{aligned}$$

Since  $[x] = x - \psi(x) - \frac{1}{2}$ ,  $\sum_{j \leq q} \chi(j) = 0$  for non-principal characters and  $\psi(x) = \psi(x+1)$ , we have

$$\begin{aligned}
T_{a,b}(x, \chi) &= \sum_{j \leq q} \chi^b(j) \sum_{n \leq x^{1/(a+b)}} \chi^a(n) \left( \frac{x^{1/b}}{qn^{a/b}} - \frac{j}{q} + \frac{1}{2} - \psi \left( \frac{x^{1/b}}{qn^{a/b}} - \frac{j}{q} \right) \right) \\
&\quad + \sum_{j \leq q} \chi^a(j) \sum_{m \leq x^{1/(a+b)}} \chi^b(m) \left( \frac{x^{1/a}}{qm^{b/a}} - \frac{j}{q} + \frac{1}{2} - \psi \left( \frac{x^{1/a}}{qm^{b/a}} - \frac{j}{q} \right) \right) \\
&\quad - \sum_{j \leq q} \sum_{h \leq q} \chi^a(j) \chi^b(h) \left( \frac{x^{1/(a+b)}}{q} - \frac{j}{q} + \frac{1}{2} - \psi \left( \frac{x^{1/(a+b)}}{q} - \frac{j}{q} \right) \right) \\
&\quad \times \left( \frac{x^{1/(a+b)}}{q} - \frac{h}{q} + \frac{1}{2} - \psi \left( \frac{x^{1/(a+b)}}{q} - \frac{h}{q} \right) \right) \\
&= - \sum_{j \leq q} \chi^b(j) \sum_{n \leq x^{1/(a+b)}} \chi^a(n) \left( \frac{j}{q} + \psi \left( \frac{x^{1/b}}{qn^{a/b}} - \frac{j}{q} \right) \right) \\
&\quad - \sum_{j \leq q} \chi^a(j) \sum_{m \leq x^{1/(a+b)}} \chi^b(m) \left( \frac{j}{q} + \psi \left( \frac{x^{1/a}}{qm^{b/a}} - \frac{j}{q} \right) \right) \\
&\quad - \sum_{j \leq q} \sum_{h \leq q} \chi^a(j) \chi^b(h) \left( \frac{j}{q} + \psi \left( \frac{x^{1/(a+b)}}{q} - \frac{j}{q} \right) \right) \left( \frac{h}{q} + \psi \left( \frac{x^{1/(a+b)}}{q} - \frac{h}{q} \right) \right).
\end{aligned}$$

In view of (2.2) of Lemma 2.3, we have

$$\begin{aligned}
T_{a,b}(x, \chi) &= - \sum_{j \leq q} \sum_{h \leq q} \chi^b(j) \chi^a(h) \sum_{\substack{n \leq x^{1/(a+b)} \\ n \equiv h \pmod{q}}} \psi \left( \frac{x^{1/(a+b)}}{qn^{a/b}} - \frac{j}{q} \right) \\
&\quad - \sum_{j \leq q} \sum_{h \leq q} \chi^a(j) \chi^b(h) \sum_{\substack{m \leq x^{1/(a+b)} \\ m \equiv j \pmod{q}}} \psi \left( \frac{x^{1/(a+b)}}{qm^{b/a}} - \frac{h}{q} \right) + O(q^2) \\
&=: -S_5 - S_6 + O(q^2).
\end{aligned}$$

If we use Lemma 2.6 with the exponent pair  $(\frac{2}{7}, \frac{4}{7})$ , then, for  $q \leq x^{1/(a+b)}$ , we have

$$\begin{aligned}
S_5 &= \sum_{j \leq q} \sum_{h \leq q} \chi^b(j) \chi^a(h) R \left( \frac{x^{1/b}}{q}, x^{1/(a+b)}, \frac{a}{b}, q, h, \frac{-j}{q} \right) \\
&\ll \sum_{j \leq q} \sum_{h \leq q} (x^{1/(2a+2b)} q^{-1/2} + x^{2/(3a+3b)} q^{-2/3}) \\
&= O(x^{2/(3a+3b)} q^{4/3})
\end{aligned}$$

and similarly

$$S_6 = O(x^{2/(3a+3b)} q^{4/3}).$$

Thus,

$$(3.15) \quad T_{a,b}(x, \chi) = O(x^{2/(3a+3b)} q^{4/3}).$$

Considering (3.8), (3.12), (3.13), (3.14) and (3.15), we deduce that

$$T_{a,b}(x, \chi) = \begin{cases} \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} + \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} \\ \quad + O(x^{1/(2b)} q^{1/3+\varepsilon}) & \text{for } \chi = \chi_0, \\ \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi^b\right) x^{1/a} + O(x^{1/(2b)} q^{4/3+\varepsilon}) & \text{for } \chi \in A_\chi - B_\chi, \\ \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi^a\right) x^{1/b} + O(x^{1/(2b)} q^{4/3+\varepsilon}) & \text{for } \chi \in B_\chi - A_\chi, \\ \frac{\varphi(q)}{q} L\left(\frac{b}{a}, \chi_0\right) x^{1/a} + \frac{\varphi(q)}{q} L\left(\frac{a}{b}, \chi_0\right) x^{1/b} \\ \quad + O(x^{1/(2b)} q^{1/3+\varepsilon}) & \text{for } \chi \in A_\chi \cap B_\chi, \\ O(x^{1/(2b)} q^{4/3}) & \text{for } \chi \notin A_\chi \cup B_\chi. \end{cases}$$

We return to the asymptotics for (3.4). In view of (3.8), we obtain the asymptotic formula for the first case  $\chi = \chi_0$ ,

$$(3.16) \quad \begin{aligned} C_{a,b}(x, \chi_0) &= \sum_{d \leq x^{1/(2b)}} \mu(d) \chi_0^{2b}(d) T_{a,b}(x d^{-2b}, \chi_0) \\ &= \frac{\varphi(q)}{q} \frac{L(b/a, \chi_0)}{L(2b/a, \chi_0)} x^{1/a} + \frac{\varphi(q)}{q} \frac{L(a/b, \chi_0)}{L(2, \chi_0)} x^{1/b} + O(x^{1/(2b)} q^{1/3+\varepsilon}). \end{aligned}$$

For other non-principal characters, we proceed in a similar way. For  $q \leq x^{1/(a+b)}$ , we have

$$(3.17) \quad C_{a,b}(x, \chi) = \begin{cases} \frac{\varphi(q)}{q} \frac{L(b/a, \chi^b)}{L(2b/a, \chi^{2b})} x^{1/a} + O(x^{1/(2b)} q^{4/3+\varepsilon}) & \text{for } \chi \in A_\chi - B_\chi, \\ \frac{\varphi(q)}{q} \frac{L(a/b, \chi^a)}{L(2, \chi^{2b})} x^{1/b} + O(x^{1/(2b)} q^{4/3+\varepsilon}) & \text{for } \chi \in B_\chi - A_\chi, \\ \frac{\varphi(q)}{q} \frac{L(b/a, \chi_0)}{L(2b/a, \chi^{2b})} x^{1/a} + \frac{\varphi(q)}{q} \frac{L(a/b, \chi_0)}{L(2, \chi^{2b})} x^{1/b} \\ \quad + O(x^{1/(2b)} q^{1/3+\varepsilon}) & \text{for } \chi \in A_\chi \cap B_\chi, \\ O(x^{1/(2b)} q^{4/3}) & \text{for } \chi \notin A_\chi \cup B_\chi. \end{cases}$$

Since  $a < b < 2a$ ,  $\frac{1}{4} < a/(2b) < \frac{1}{2}$  and  $1 < b/a < 2$ , then  $(2b^2 - ab - a^2)/(2ab) < \frac{4}{3}$ . Thus,  $O(x^{1/(2b)} q^{4/3})$  dominates the bound of  $U_2$  in (3.3) and Theorem 1.1 follows from (3.1), (3.16) and (3.17).  $\square$

**Acknowledgements.** The authors express their gratitude to the referee for his/her helpful and detailed comments.

### References

- [1] *P. T. Bateman, E. Grosswald*: On a theorem of Erdős and Szekeres. III. *J. Math.* *2* (1958), 88–98. [zbl](#) [MR](#) [doi](#)
- [2] *T. H. Chan*: Squarefull numbers in arithmetic progression. II. *J. Number Theory* *152* (2015), 90–104. [zbl](#) [MR](#) [doi](#)
- [3] *T. H. Chan, K. M. Tsang*: Squarefull numbers in arithmetic progressions. *Int. J. Number Theory* *9* (2013), 885–901. [zbl](#) [MR](#) [doi](#)
- [4] *E. Cohen*: Arithmetical notes. II: An estimate of Erdős and Szekeres. *Scripta Math.* *26* (1963), 353–356. [zbl](#) [MR](#)
- [5] *P. Erdős, S. Szekeres*: Über die Anzahl der Abelschen Gruppen gegebener Ordnung und über ein verwandtes zahlentheoretisches Problem. *Acta Szeged* *7* (1934), 95–102. (In German.) [zbl](#)
- [6] *H. Liu, T. Zhang*: On the distribution of square-full numbers in arithmetic progressions. *Arch. Math.* *101* (2013), 53–64. [zbl](#) [MR](#) [doi](#)
- [7] *M. Munsch*: Character sums over squarefree and squarefull numbers. *Arch. Math.* *102* (2014), 555–563. [zbl](#) [MR](#) [doi](#)
- [8] *H.-E. Richert*: Über die Anzahl Abelscher Gruppen gegebener Ordnung. I. *Math. Z.* *56* (1952), 21–32. (In German.) [zbl](#) [MR](#) [doi](#)
- [9] *H.-E. Richert*: Über die Anzahl Abelscher Gruppen gegebener Ordnung. II. *Math. Z.* *58* (1953), 71–84. (In German.) [zbl](#) [MR](#) [doi](#)
- [10] *T. Srichan*: Square-full and cube-full numbers in arithmetic progressions. *Šiauliai Math. Semin.* *8* (2013), 223–248. [zbl](#) [MR](#)

*Authors' addresses:* Angkana Sripayap, Department of Mathematics, Faculty of Science, Kasetsart University, 50 Thanon Ngam Wong Wan, Lat Yao, Chatuchak, Bangkok 10900, Thailand, e-mail: [fscianr@ku.ac.th](mailto:fscianr@ku.ac.th); Pattira Ruengsinsub (corresponding author), Department of Mathematics, Faculty of Science, Kasetsart University, 50 Thanon Ngam Wong Wan, Lat Yao, Chatuchak, Bangkok 10900, Thailand, e-mail: [fscipar@ku.ac.th](mailto:fscipar@ku.ac.th); Teerapat Srichan, Department of Mathematics, Faculty of Science, Kasetsart University, 50 Thanon Ngam Wong Wan, Lat Yao, Chatuchak, Bangkok 10900, Thailand, e-mail: [fscitrp@ku.ac.th](mailto:fscitrp@ku.ac.th).