

Jonathan David Farley

Does the endomorphism poset  $P^P$  determine whether a finite poset  $P$  is connected? An issue Duffus raised in 1978

*Mathematica Bohemica*, Vol. 148 (2023), No. 4, 435–446

Persistent URL: <http://dml.cz/dmlcz/151965>

## Terms of use:

© Institute of Mathematics AS CR, 2023

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

DOES THE ENDOMORPHISM POSET  $P^P$  DETERMINE  
 WHETHER A FINITE POSET  $P$  IS CONNECTED?  
 AN ISSUE DUFFUS RAISED IN 1978

JONATHAN DAVID FARLEY, Baltimore

Received January 19, 2022. Published online August 29, 2022.  
 Communicated by Radomír Halaš

*Abstract.* Duffus wrote in his 1978 Ph.D. thesis, “It is not obvious that  $P$  is connected and  $P^P \cong Q^Q$  imply that  $Q$  is connected”, where  $P$  and  $Q$  are finite nonempty posets. We show that, indeed, under these hypotheses  $Q$  is connected and  $P \cong Q$ .

*Keywords:* (partially) ordered set; exponentiation; connected

*MSC 2020:* 06A07

## 1. INTRODUCTION

In the 1979 *Proceedings of the American Mathematical Society*, Duffus and Wille proved that  $P^P \cong Q^Q$  implies  $P \cong Q$  if  $P$  and  $Q$  are both finite, nonempty, and connected (see [8], Theorem, page 14). Duffus said in 1984 in [6], page 90, “It is still an open problem to show connectedness can be dropped.” He notes in his 1978 thesis [4], page 53, “It is not obvious that  $P$  is connected and  $P^P \cong Q^Q$  imply that  $Q$  is connected.” (Note that, if this is the case, then the Duffus-Wille result implies  $P \cong Q$ .) We prove that, indeed, if  $P$  and  $Q$  are finite, nonempty posets such that  $P^P \cong Q^Q$  and  $P$  is connected, then  $Q$  is connected.

In other words, we have resolved the issue from Duffus’s 1978 thesis in that we have “half-dropped” the connectedness hypothesis used in the 1979 *Proceedings of the American Mathematical Society* paper, whereas the 1984 problem asked if it could be dropped entirely.

We assume the reader is familiar with the basic facts about the arithmetic of ordered sets, the basic consequences of Hashimoto’s refinement theorem, and Professor

Birkhoff's theorem on finite distributive lattices (e.g., [16], Propositions 3.1 and 4.1 and [3], Theorems 5.9 and 5.12 and also see [17]).

Let  $P$  and  $Q$  be posets. For  $p, p' \in P$ , we write  $p \equiv p'$  if  $p$  and  $p'$  are in the same connected component of  $P$ ;  $h(P)$  denotes the height of the finite, nonempty poset  $P$ , the largest value of  $|C| - 1$  for  $C$  a chain in  $P$ , and  $h_P(p)$  denotes the height of an element  $p$  in  $P$ . Note that  $h(P \times Q) = h(P) + h(Q)$  for  $P$  and  $Q$  finite and nonempty (see [2], Chapter I, §9, Exercise 4 (a)).

Let  $P^Q$  denote the poset of order-preserving maps from  $Q$  to  $P$ , where, for  $f, g \in P^Q$ ,

$$f \leq g \quad \text{if for all } q \in Q, \quad f(q) \leq_P g(q)$$

(see [1], page 312). For  $p \in P$ , denote the constant map  $f(q) = p$  for all  $q \in Q$  by  $\langle p \rangle$ . Let  $\mathcal{D}(P^Q)$  denote

$$\{g \in P^Q : g \text{ is constant on each connected component of } Q\}$$

and for  $Q \neq \emptyset$  let  $\mathcal{C}(P^Q)$  denote

$$\{f \in P^Q : f \equiv g \text{ for some } g \in \mathcal{D}(P^Q)\}$$

(see [16], Definition 2.1). Note that, despite the notation,  $\mathcal{C}(P^Q)$  is really a function of  $P$  and  $Q$ , not  $P^Q$  (as far as the author knows).




Figure 1.  $P^P$  where  $P$  is the 4-element crown, see [4], page 54.

The  $n$ -element chain is denoted  $\mathbf{n}$ . A poset  $P$  is *directly irreducible* if  $|P| \neq 1$  and whenever  $P \cong A \times B$  for posets  $A$  and  $B$ , then  $|A| = 1$  or  $|B| = 1$ . A finite poset  $P$  is *absolutely  $\mathcal{C}$ -indecomposable* if it is connected, directly irreducible, and, whenever  $P \cong \mathcal{C}(A^B)$  for posets  $A$  and  $B$ ,  $B \neq \emptyset$ , we have  $P \cong A$  and  $|B| = 1$ .

The following result comes from a remark McKenzie leaves to the reader after [16], Proposition 4.1, which nonetheless is true for finite, nonempty posets. (For a proof of (1), see [11], Theorem 2.8.)

**Proposition 1.** *Let  $P$ ,  $Q$ , and  $R$  be nonempty posets. Then:*

- (1)  $\mathcal{C}(P^{Q \times R}) \cong \mathcal{C}(\mathcal{C}(P^Q)^R)$  if  $P$ ,  $Q$ , and  $R$  are finite;
- (2)  $\mathcal{C}(P^{Q+R}) \cong \mathcal{C}(P^Q) \times \mathcal{C}(P^R)$ ;
- (3) if  $R$  is connected,  $\mathcal{C}((P+Q)^R) \cong \mathcal{C}(P^R) + \mathcal{C}(Q^R)$ ;
- (4)  $\mathcal{C}((Q \times R)^P) \cong \mathcal{C}(Q^P) \times \mathcal{C}(R^P)$ .

We also use the structure theorems of McKenzie (see [16], Theorems 8.1, 9.1–9.2 and [15], Theorem 5.1):

**Theorem 2.**

- (1) *Let  $A$ ,  $B$ ,  $C$ , and  $D$  be finite, nonempty, connected posets. Assume  $C$  and  $D$  are directly irreducible and  $C \not\cong D$ . Then if  $\mathcal{C}(A^C) \cong \mathcal{C}(B^D)$ , there exists a finite, nonempty, connected poset  $E$  such that  $A \cong \mathcal{C}(E^D)$  and  $B \cong \mathcal{C}(E^C)$ . (Note that “ $A^Q$ ” should be “ $E^Q$ ” on [16], page 211.)*
- (2) *Let  $A$ ,  $B$ , and  $C$  be finite, nonempty posets. Assume  $C$  is connected. Then  $A \cong B$  if  $\mathcal{C}(A^C) \cong \mathcal{C}(B^C)$ .*
- (3) *Let  $n \in \mathbb{N}$ . Let  $A$  and  $B$  be finite, nonempty, connected posets, and let  $C_1, \dots, C_n$  be posets such that  $\mathcal{C}(A^B) \cong C_1 \times \dots \times C_n$ . Then there exist finite, nonempty posets  $A_1, \dots, A_n$  such that  $C_i \cong \mathcal{C}(A_i^B)$  for  $i = 1, \dots, n$  and  $A \cong A_1 \times \dots \times A_n$ .*
- (4) *Let  $A$  be a finite, connected, directly irreducible poset. Then there exists an absolutely  $\mathcal{C}$ -indecomposable connected poset  $B$  and a finite, nonempty poset  $P$  such that  $A \cong \mathcal{C}(B^P)$ .*

We now extend parts of Theorem 2 to the analogue of [12], Theorem 8.2, using similar steps. The proofs of Lemma 3 and Theorem 4 first occurred in [9], but an editor asked that they be included as [10], Lemma 8 and Theorem 9.

**Lemma 3.** *Let  $A$ ,  $B$ ,  $C$ , and  $D$  be finite, nonempty, connected posets such that  $\mathcal{C}(A^C) \cong \mathcal{C}(B^D)$ . Assume that no nontrivial poset is isomorphic to both a direct factor of  $C$  and a direct factor of  $D$ . Then there exists a finite, nonempty, connected poset  $E$  such that  $A \cong \mathcal{C}(E^D)$  and  $B \cong \mathcal{C}(E^C)$ .*

**Theorem 4.** *Let  $A$ ,  $B$ ,  $C$ , and  $D$  be finite, nonempty, connected posets such that  $\mathcal{C}(A^C) \cong \mathcal{C}(B^D)$ . Then there exist finite, nonempty, connected posets  $E$ ,  $X$ ,  $Y$ , and  $Z$  such that  $A \cong \mathcal{C}(E^X)$ ,  $B \cong \mathcal{C}(E^Y)$ ,  $C \cong Y \times Z$ , and  $D \cong X \times Z$ .*

Proof. Hashimoto's refinement theorem tells us we can find finite, nonempty, connected posets  $X$ ,  $Y$ , and  $Z$  such that  $C \cong Y \times Z$ ,  $D \cong X \times Z$ , and  $X$  and  $Y$  do not share a nontrivial direct factor. Thus, by Proposition 1 (1),

$$\mathcal{C}(\mathcal{C}(A^Y)^Z) \cong \mathcal{C}(A^{Y \times Z}) \cong \mathcal{C}(A^C) \cong \mathcal{C}(B^D) \cong \mathcal{C}(B^{X \times Z}) \cong \mathcal{C}(\mathcal{C}(B^X)^Z).$$

Now  $\mathcal{C}(A^Y), \mathcal{C}(B^X) \neq \emptyset$  since  $A, B, X$ , and  $Y$  are nonempty and connected. Hence, by Theorem 2 (2),  $\mathcal{C}(A^Y) \cong \mathcal{C}(B^X)$ . By Lemma 3, there exists a finite, nonempty, connected poset  $E$  such that  $A \cong \mathcal{C}(E^X)$  and  $B \cong \mathcal{C}(E^Y)$ .  $\square$

In [4], Theorem 3.2.6, Duffus proves an outstanding version of Hashimoto's refinement theorem for posets  $A, B, C$ , and  $D$  that are sums of connected posets with a finite maximal chain such that  $A \times B \cong C \times D$  and  $A$  and  $C$  are connected. He never published this proof, although it is used in [8]. We only require one of the four posets to be connected. (We don't actually need this result, but the structure of its proof helps the reader understand the proof of the theorem we do need: this proof foreshadows the proof of Theorem 10.)

**Theorem 5.** *Let  $A, B, C$ , and  $D$  be finite posets. Let  $A$  be connected and nonempty. Assume that  $A \times B \cong C \times D$ . Then there exist posets  $W, X, Y$ , and  $Z$  such that  $A \cong W \times X$ ,  $B \cong Y \times Z$ ,  $C \cong W \times Y$ , and  $D \cong X \times Z$ .*

Proof. Note that  $B = \emptyset$  if and only if either  $C = \emptyset$  or  $D = \emptyset$ . First assume  $B = \emptyset = C$ . Then let  $W = A$ ,  $X = \mathbf{1}$ ,  $Y = \emptyset$ , and  $Z = D$ . Next, assume  $B = \emptyset = D$ . Then let  $W = \mathbf{1}$ ,  $X = A$ ,  $Y = C$ , and  $Z = \emptyset$ .

From now on, assume that  $B, C$ , and  $D$  are nonempty.

Let  $B = \sum_{h \in H} B_h$ ,  $C = \sum_{i \in I} C_i$ , and  $D = \sum_{j \in J} D_j$  be the decompositions of  $B, C$ , and  $D$ , respectively, into connected components. Since  $A \times B \cong \sum_{h \in H} A \times B_h$  has the same number of components as  $C \times D \cong \sum_{i \in I, j \in J} C_i \times D_j$ , there is a bijection  $\Psi: I \times J \rightarrow H$  such that  $C_i \times D_j \cong A \times B_{\Psi(i,j)}$  for all  $i \in I, j \in J$ .

Let  $A_1, A_2, \dots, A_r$  ( $r \geq 0$ ) be the pairwise nonisomorphic connected, directly irreducible posets that could arise in the factorizations of any of  $A, B_h$  ( $h \in H$ ),  $C_i$  ( $i \in I$ ), and  $D_j$  ( $j \in J$ ). Say  $A \cong \prod_{l=1}^r (A_l)^{k_l}$  where  $k_l \geq 0$  ( $l = 1, \dots, r$ ). For each  $l \in \{1, \dots, r\}$ , let  $c_l \in \mathbb{N}_0$  be the highest power of  $A_l$  such that  $A_l^{c_l}$  is a factor of all of the  $C_i$  ( $i \in I$ ). Let  $W := \prod_{l=1}^r A_l^{\min\{c_l, k_l\}}$ . Let  $X := \prod_{l=1}^r A_l^{k_l - \min\{c_l, k_l\}}$ . Then  $W \times X \cong \prod_{l=1}^r A_l^{k_l} \cong A$ . Clearly, for all  $i \in I$ ,  $W$  is a factor of  $C_i$ . Hence, we may let  $\tilde{C}_i$  be  $C_i$  with  $W$  factored out, that is,  $C_i \cong \tilde{C}_i \times W$  ( $i \in I$ ). Let  $Y := \sum_{i \in I} \tilde{C}_i$ .

**Claim 6.** *For each  $j \in J$ ,  $X$  is a factor of  $D_j$ .*

Proof. Assume for a contradiction that  $X$  does not divide  $D_j$  for some  $j \in J$ . Then there exists  $l \in \{1, \dots, r\}$  such that  $A_l^{k_l - c_l}$  does not divide  $D_j$  and so  $k_l > c_l$ . Pick  $i \in I$  such that  $c_l$  is the highest power of  $A_l$  dividing  $C_i$ . Then the highest power of  $A_l$  dividing  $C_i \times D_j$  is less than  $c_l + k_l - c_l = k_l$ , a contradiction, since  $C_i \times D_j \cong A \times B_{\Psi(i,j)}$  and  $A_l^{k_l}$  divides the right-hand side.  $\square$

By Claim 6, we may let  $\tilde{D}_j$  be  $D_j$  with  $X$  factored out ( $j \in J$ ). Let  $Z = \sum_{j \in J} \tilde{D}_j$ .

**Claim 7.** For all  $(i, j) \in I \times J$ ,  $B_{\Psi(i,j)} \cong \tilde{C}_i \times \tilde{D}_j$ .

Proof. We know

$$A \times B_{\Psi(i,j)} \cong C_i \times D_j \cong W \times X \times \tilde{C}_i \times \tilde{D}_j \cong A \times \tilde{C}_i \times \tilde{D}_j,$$

so  $B_{\Psi(i,j)} \cong \tilde{C}_i \times \tilde{D}_j$ . (See [13], (4.3).)  $\square$

Thus  $B \cong Y \times Z$ . By definition,  $C \cong W \times Y$  and  $D \cong X \times Z$ .  $\square$

A “strong” or “strict” version of Theorem 5 would be useful (see [16], Proposition 3.1).

**Lemma 8.** Let  $A$  be a finite, connected, directly irreducible poset. Let  $B$  and  $P$  be posets such that  $P \neq \emptyset$ . If  $A \cong \mathcal{C}(B^P)$ , then  $P$  is finite and connected, and  $B$  is finite, connected, and directly irreducible.

Proof. We have  $B \neq \emptyset$ . Also  $|B| \geq 2$ . If  $P = C + D$  for posets  $C, D \neq \emptyset$ , then, by Proposition 1 (2),  $A \cong \mathcal{C}(B^C) \times \mathcal{C}(B^D)$ . Therefore since  $A$  is directly irreducible,  $\mathbf{1} \cong \mathcal{C}(B^C)$  or  $\mathbf{1} \cong \mathcal{C}(B^D)$ , a contradiction ( $|\mathcal{C}(B^C)|, |\mathcal{C}(B^D)| \geq 2$  since  $C, D \neq \emptyset$ ). Hence  $P$  is connected.

If  $B = E + F$  for posets  $E, F \neq \emptyset$ , then by Proposition 1 (3),  $A \cong \mathcal{C}(E^P) + \mathcal{C}(F^P)$ , a contradiction. Hence  $B$  is connected. Thus there exist  $x, y \in B$  such that  $x < y$ , so if  $P$  is infinite, then  $|P| \leq |\mathbf{2}^P| \leq |\mathcal{C}(B^P)|$ , a contradiction. Hence  $P$  is finite. Also,  $B$  is finite. If  $B \cong G \times H$  for posets  $G$  and  $H$ , then by Proposition 1 (4),  $A \cong \mathcal{C}(G^P) \times \mathcal{C}(H^P)$ , therefore, by the direct irreducibility of  $A$ , without loss of generality  $|\mathcal{C}(G^P)| = 1$ , so  $|G| = 1$ .  $\square$

**Lemma 9.** Let  $A$  be a connected, finite, directly irreducible poset. Let  $B$  be a finite, nonempty, connected poset. Then:

- (1)  $\mathcal{C}(A^B)$  is connected and directly irreducible.
- (2) Let  $C$  and  $D$  be nonempty posets. If  $\mathcal{C}(C^D) \cong \mathcal{C}(A^B)$ , then  $C$  is directly irreducible, finite, and connected, and  $D$  is finite and connected.

(3) There exist unique (up to isomorphism) nonempty posets  $E$  and  $F$  with the following two properties:

(a)  $\mathcal{C}(A^B) \cong \mathcal{C}(E^F)$  and

(b) whenever  $C$  and  $D$  are nonempty posets such that  $\mathcal{C}(A^B) \cong \mathcal{C}(C^D)$ , then there exists a poset  $G$  such that  $\mathcal{C}(E^G) \cong C$  and  $G \times D \cong F$ .

We can choose as  $E$  any absolutely  $\mathcal{C}$ -indecomposable poset  $E$  such that  $\mathcal{C}(E^J) \cong \mathcal{C}(A^B)$  for some nonempty poset  $J$ , and we can choose that  $J$  as our “ $F$ ”. If  $H$  is a finite, nonempty, connected poset, then  $E$  and  $F \times H$  are the posets that work for  $\mathcal{C}(A^{B \times H})$ .

**Proof.** (1) The poset  $\mathcal{C}(A^B)$  is nontrivial since  $A$  is nontrivial. By Theorem 2 (3),  $\mathcal{C}(A^B)$  is directly irreducible. It is connected since  $A$  and  $B$  are connected.

(2) See Lemma 8.

(3) Using (1), take the poset  $E$  given by Theorem 2(4). (We want  $E$  that is absolutely  $\mathcal{C}$ -indecomposable.) There exists a nonempty finite poset  $F$  such that  $\mathcal{C}(E^F) \cong \mathcal{C}(A^B)$ . By (2),  $F$  is connected.

If  $\mathcal{C}(A^B) \cong \mathcal{C}(C^D)$ , where  $C$  and  $D$  are nonempty posets, then by (2)  $C$  and  $D$  are finite and connected and  $\mathcal{C}(E^F) \cong \mathcal{C}(C^D)$ , so by Theorem 4, there exist nonempty, finite, connected posets  $U, G, W$ , and  $X$  such that

$$C \cong \mathcal{C}(U^G), \quad E \cong \mathcal{C}(U^W), \quad D \cong W \times X, \quad F \cong G \times X.$$

By absolute  $\mathcal{C}$ -indecomposability,  $|W| = 1$  and  $E \cong U$ , so the result follows.

If  $E'$  and  $F'$  have the same property as  $E$  and  $F$ , then there exists a poset  $G'$  such that  $\mathcal{C}(E'^{G'}) \cong E$  and  $G' \times F \cong F'$ . By absolute  $\mathcal{C}$ -indecomposability,  $E' \cong E$  and  $|G'| = 1$ , so  $F \cong F'$ .  $\square$

**Theorem 10.** Let  $A, B, C$ , and  $D$  be finite nonempty posets such that  $A$  is connected,  $B$  and  $D$  are nontrivial and connected, and  $\mathcal{C}(B^A) \cong \mathcal{C}(D^C)$ . Then there exist finite, nonempty posets  $W, X, Y$ , and  $Z$  such that  $Z$  is nontrivial and  $A \cong W \times X$ ,  $B \cong \mathcal{C}(Z^Y)$ ,  $C \cong W \times Y$ , and  $D \cong \mathcal{C}(Z^X)$ .

**Proof.** Let  $B = \prod_{h \in H} B_h$  be a representation of  $B$  as a product of connected, directly irreducible posets. Let  $D = \prod_{j \in J} D_j$  be a similar product. Let  $C = \sum_{i \in I} C_i$  be a decomposition into connected components. Let  $A_l$  ( $l \in L$ ) be the finitely many pairwise nonisomorphic connected, directly irreducible posets that could arise in the factorizations of  $A$  and  $C_i$  ( $i \in I$ ). Say  $A \cong \prod_{l \in L} A_l^{k_l}$  where  $k_l \geq 0$  ( $l \in L$ ). For each  $l \in L$ , let  $c_l \in \mathbb{N}_0$  be the maximum power of  $A_l$  such that  $A_l^{c_l}$  is a factor of all of the  $C_i$  ( $i \in I$ ). That means, of course, that it is the highest power of  $A_l$  in one

of the  $C_i$ . Let  $W = \prod_{l \in L} A_l^{\min\{c_l, k_l\}}$ . Let  $X = \prod_{l \in L} A_l^{k_l - \min\{c_l, k_l\}}$ . Then

$$W \times X \cong \prod_{l \in L} A_l^{k_l} \cong A.$$

**Claim 11.** *For all  $i \in I$ ,  $W$  is a factor of  $C_i$ .*

**Proof.** We made sure that  $A_l^{c_l}$  is a factor of every  $C_i$ , and  $W$  is a product of all of those powers of  $A_l$  or even smaller powers. Since the different  $A_l$  are pairwise nonisomorphic and we are working with connected posets, by Hashimoto's refinement theorem we have "unique factorization", so  $A_l^{c_l} \times A_{l'}^{c_{l'}}$  is a factor if and only if  $A_l^{c_l}$  is a factor, and  $A_{l'}^{c_{l'}}$  is a factor, when  $l \neq l'$ .  $\square$

By Claim 11, we may let  $\tilde{C}_i$  be  $C_i$  with  $W$  factored out ( $i \in I$ ). Let  $Y = \sum_{i \in I} \tilde{C}_i$ . Then  $W \times Y \cong \sum_{i \in I} W \times \tilde{C}_i \cong \sum_{i \in I} C_i = C$ .

Note that by Proposition 1 (1),

$$\mathcal{C}(B^A) \cong \mathcal{C}(\mathcal{C}(B^X)^W) \quad \text{and} \quad \mathcal{C}(D^C) \cong \mathcal{C}(\mathcal{C}(D^Y)^W),$$

and by Theorem 2 (2), since  $W$ , being a factor of a finite, nonempty, connected poset, is finite, nonempty, and connected,  $\mathcal{C}(B^X) \cong \mathcal{C}(D^Y)$ .

**Claim 12.**  *$X$  and  $Y$  have no nontrivial factor in common.*

**Proof.** Assume for a contradiction that  $X$  and  $Y$  do have a nontrivial factor in common. Then since  $X$  is a factor of  $A$ , we may assume the common factor is  $A_l$  for some  $l \in L$ . As  $A$  is connected, for  $A_l$  to be a factor of  $Y$ , it must be a factor of  $\tilde{C}_i$  for all  $i \in I$ . But then we would have pulled it out with  $W$ , as it were. To be precise, since  $A_l$  is a factor of  $X$ , we must have  $c_l < k_l$ , and so there is  $i \in I$  such that  $\tilde{C}_i$  has no factor of  $A_l$ , a contradiction.  $\square$

Write  $\mathcal{C}(B^X) \cong \prod_{h \in H} \mathcal{C}(B_h^X)$  and  $\mathcal{C}(D^Y) \cong \prod_{i \in I} \prod_{j \in J} \mathcal{C}(D_j^{\tilde{C}_i})$ . By Lemma 9 (1),  $\mathcal{C}(B_h^X)$  ( $h \in H$ ) and  $\mathcal{C}(D_j^{\tilde{C}_i})$  ( $i \in I, j \in J$ ) are directly irreducible. As those factors are connected, by [5], Corollary 2 (cf. [14], Theorem 6.4) there is a bijection  $\Psi: I \times J \rightarrow H$  such that  $\mathcal{C}(D_j^{\tilde{C}_i}) \cong \mathcal{C}(B_h^X)$  for all  $(i, j) \in I \times J$ . By Theorem 4 there are finite, nonempty, connected posets  $U_h, R_h, S_h$ , and  $T_h$  (where  $h = \Psi(i, j)$ ) such that

$$B_h \cong \mathcal{C}(U_h^{R_h}), \quad D_j \cong \mathcal{C}(U_h^{S_h}), \quad X \cong S_h \times T_h, \quad \text{and} \quad \tilde{C}_i \cong R_h \times T_h.$$

Moreover, by Lemma 8,  $U_h$  is directly irreducible.

**Claim 13.** *For each  $j \in J$ , choose some  $i' \in I$  and let  $E_j$  and  $F_j$  be the posets corresponding to  $D_j$  (where  $h = \Psi(i', j)$ ) given by part (3) of Lemma 9. Then  $X$  is a factor of  $F_j$ .*



Proof. By part (2) of Lemma 9,  $F_j$  is finite and connected. Assume for a contradiction that  $X$  does not divide  $F_j$ . Then there exists  $l \in L$  such that  $A_l^{k_l - c_l}$  does not divide  $F_j$ , and so  $k_l > c_l$ . Pick  $i \in I$  such that  $c_l$  is the highest power of  $A_l$  dividing  $C_i$ . Consider

$$\mathcal{C}(D_j^{C_i}) \cong \mathcal{C}(D_j^{\tilde{C}_i \times W}) \cong \mathcal{C}(\mathcal{C}(D_j^{\tilde{C}_i})^W) \cong \mathcal{C}(\mathcal{C}(B_{\Psi(i,j)}^X)^W) \cong \mathcal{C}(B_{\Psi(i,j)}^{X \times W}) \cong \mathcal{C}(B_{\Psi(i,j)}^A).$$

By part (3) of Lemma 9,  $A$  is a factor of  $F_j \times C_i$  and  $A_l^{k_l}$  divides  $A$ , so  $A_l^{k_l - c_l}$  divides  $F_j$ , a contradiction.  $\square$

By Claim 13, we may let  $\tilde{F}_j$  be  $F_j$  with  $X$  factored out ( $j \in J$ ).

**Claim 14.** For all  $(i, j) \in I \times J$ ,  $B_{\Psi(i,j)} \cong \mathcal{C}(E_j^{\tilde{C}_i \times \tilde{F}_j})$ .

Proof. Note that

$$\mathcal{C}(B_{\Psi(i,j)}^A) \cong \mathcal{C}(D_j^{C_i}) \cong \mathcal{C}(\mathcal{C}(E_j^{F_j})^{C_i}) \cong \mathcal{C}(\mathcal{C}(E_j^{X \times \tilde{F}_j})^{W \times \tilde{C}_i}) \cong \mathcal{C}(\mathcal{C}(E_j^{\tilde{C}_i \times \tilde{F}_j})^A)$$

so  $B_{\Psi(i,j)} \cong \mathcal{C}(E_j^{\tilde{C}_i \times \tilde{F}_j})$  by Theorem 2(2) since  $A$  is connected. Thus

$$B \cong \prod_{i \in I, j \in J} \mathcal{C}(\mathcal{C}(E_j^{\tilde{F}_j})^{\tilde{C}_i}) \cong \prod_{j \in J} \mathcal{C}(\mathcal{C}(E_j^{\tilde{F}_j})^Y) \cong \mathcal{C}\left(\left(\prod_{j \in J} \mathcal{C}(E_j^{\tilde{F}_j})\right)^Y\right).$$

Let  $Z = \prod_{j \in J} \mathcal{C}(E_j^{\tilde{F}_j})$ . We know

$$D \cong \prod_{j \in J} D_j \cong \prod_{j \in J} \mathcal{C}(E_j^{F_j}) \cong \prod_{j \in J} \mathcal{C}(\mathcal{C}(E_j^{\tilde{F}_j})^X) \cong \mathcal{C}\left(\left(\prod_{j \in J} \mathcal{C}(E_j^{\tilde{F}_j})\right)^X\right) \cong \mathcal{C}(Z^X).$$

$\square$

We also need a trivial extension of [15], Lemma 2.3 (although the added trivialities take up perhaps more space than is warranted).

**Lemma 15.** Let  $A$  and  $B$  be finite posets. Then:

- (1)  $A^B = \emptyset$  if and only if  $A = \emptyset$  and  $B \neq \emptyset$ ;
- (2)  $A^B$  is an antichain if and only if  $A$  is an antichain or  $B = \emptyset$ , in which case

$$|A^B| = \begin{cases} |A|^c & \text{if } A \neq \emptyset \text{ or } B \neq \emptyset, \text{ where } c \text{ is the number} \\ & \text{of connected components of } B, \\ 1 & \text{if } A = B = \emptyset; \end{cases}$$

- (3) let  $f, g \in A^B$  be such that  $f \leq g$ . Assume  $A \neq \emptyset$ . Then  $h_{A^B}([f, g]) = h(A^B)$  if and only if  $f, g \in \mathcal{D}(A^B)$  and  $h_A([f(b), g(b)]) = h(A)$  for all  $b \in B$ ;
- (4) when  $A \neq \emptyset$ ,  $h(A^B) = h(A)|B|$ .

**Proof.** (1) This is clear.

(2) Assume  $A^B$  is an antichain. If  $A$  is not an antichain and  $B \neq \emptyset$ , then  $\{\langle a \rangle : a \in A\} \cong A$  is a subposet of  $A^B$ , a contradiction. Conversely, if  $B = \emptyset$ , then  $c = 0$  and  $|A^B| = 1$ , so  $A^B$  is an antichain. If  $B \neq \emptyset$  but  $A$  is an antichain, then for all  $f, g \in A^B$ , if  $f < g$ , there exists  $b \in B$  such that  $f(b) < g(b)$  in  $A$ , a contradiction. Hence  $A^B$  is an antichain. For all  $f \in A^B$  and for every connected component  $C$  of  $B$ ,  $|f[C]| = 1$ , so  $|A^B| = |A|^c$ .

(3) This is [15], Lemma 2.3 (3) if  $f < g$ . Now assume  $f = g$ . If  $h_{A^B}([f, g]) = h(A^B)$ , then  $A^B$  is a nonempty antichain, so by (2) either  $B = \emptyset$  and the forward implication is vacuous or  $B \neq \emptyset$  and  $A$  is a nonempty antichain, so by the above,  $f, g \in \mathcal{D}(A^B)$  and for all  $b \in B$ ,  $f(b) = g(b)$ , so  $h_A([f(b), g(b)]) = 0 = h(A)$ .

Conversely, suppose  $f, g \in \mathcal{D}(A^B)$  and  $0 = h_A([f(b), g(b)]) = h(A)$  for all  $b \in B$ . If  $B = \emptyset$  then  $h(A^B) = h(\mathbf{1}) = 0 = h_{A^B}([f, g])$ . If  $B \neq \emptyset$ , let  $b_0 \in B$ . The fact  $h_A([f(b_0), g(b_0)]) = 0 = h(A)$  means  $A$  is an antichain.

By (1) and (2),  $A^B$  is a nonempty antichain, so  $h(A^B) = 0 = h_{A^B}([f, g])$ .

(4) This is true by [7], Corollary 2.2 if  $A, B \neq \emptyset$ . If  $A \neq \emptyset$  but  $B = \emptyset$ , then  $h(A^B) = h(\mathbf{1}) = 0 = h(A) \cdot 0$ . □

**Theorem 16.** *Let  $P$  and  $Q$  be finite, nonempty posets such that  $P$  is connected. Assume  $P^P \cong Q^Q$ . Then  $Q$  is connected, and therefore  $P \cong Q$ .*

**Proof.** If  $Q$  is connected, then  $P \cong Q$  by [8], Theorem. Assume now that  $Q$  is disconnected. Say  $Q = Q_0 + D$ , where  $D \neq \emptyset$  and  $Q_0$  is any component of  $Q$  of maximum height.

We show that  $Q_0$  is the unique component of  $Q$  of height  $h(Q)$ , and we will also show that  $Q_0$  is a proper direct factor of  $P$ .

**Claim 17.**  *$P^P$  and  $P$  are not antichains. Hence  $|Q_0| \neq 1$ .*

**Proof.** By our assumption,  $|Q| \geq 2$ , therefore  $|Q^Q| = |P^P| \geq 2$ . Hence  $|P| \geq 2$ . As  $P$  is connected,  $P$  is not an antichain and, thus  $P^P$  is not an antichain. (Look at two constant maps, the images of which form a two-element chain.) If  $|Q_0| = 1$ , then  $h(Q) = 0$  and thus  $Q$  is an antichain, so  $Q^Q \cong P^P$  is an antichain. □

Pick  $q_0 \in Q_0$  of maximum height in  $Q$ . Using Lemma 15,  $\langle q_0 \rangle$  corresponds to  $\langle p_0 \rangle$  for some  $p_0$  of maximum height in  $P$ . Since  $P$  is connected,  $\mathcal{C}(P^P) = \{f \in P^P : f \equiv \langle p_0 \rangle\}$ .

**Claim 18.**  $\mathcal{C}(Q_0^Q) = \{g \in Q_0^Q : g \equiv \langle q_0 \rangle\}$ .

*Proof.* Let  $g \in Q^Q$  be such that  $g \equiv \langle q_0 \rangle$ . Then  $g[Q] \subseteq Q_0$  and  $g \in \mathcal{C}(Q_0^Q)$ .

Let  $h \in \mathcal{C}(Q_0^Q)$ . Then  $h \in Q^Q$ , and, in  $Q_0^Q$ ,  $h \equiv k$  for some  $k \in \mathcal{D}(Q_0^Q)$ , and  $k \equiv \langle q \rangle$  for some  $q \in Q_0$ . But  $Q_0$  is connected, so  $\langle q \rangle \equiv \langle q_0 \rangle$ . Hence  $h \equiv \langle q_0 \rangle$ .  $\square$

We conclude that  $\mathcal{C}(P^P) \cong \mathcal{C}(Q_0^Q)$  via the original isomorphism. This is because we have described both  $\mathcal{C}(P^P)$  and  $\mathcal{C}(Q_0^Q)$  in terms just involving  $P^P$  and  $Q^Q$ , respectively, and in the same way, up to the isomorphism, since the original isomorphism maps  $\langle p_0 \rangle$  to  $\langle q_0 \rangle$ . Because the preceding argument can be repeated for any component of  $Q$  of maximum height,  $Q_0$  must be the only component with height equal to that of  $Q$ .

By Proposition 1 (2),  $\mathcal{C}(Q_0^Q) \cong \mathcal{C}(Q_0^{Q_0}) \times \mathcal{C}(Q_0^D) \cong \mathcal{C}(P^P)$ . By Theorem 2 (3), there exist finite, nonempty, connected posets  $A_1$  and  $A_2$  such that  $P \cong A_1 \times A_2$  and  $\mathcal{C}(Q_0^{Q_0}) \cong \mathcal{C}(A_1^P)$ . By Theorem 4, there exist nonempty, finite, connected posets  $E$ ,  $X$ ,  $Y$ , and  $Z$  such that

$$A_1 \cong \mathcal{C}(E^X), \quad Q_0 \cong \mathcal{C}(E^Y), \quad Q_0 \cong X \times Z, \quad P \cong Y \times Z.$$

Since  $E$  and  $Y$  are connected, there exists a finite, nonempty, connected poset  $F$  such that  $X \cong \mathcal{C}(F^Y)$  by Theorem 2 (3) applied to the second and third isomorphisms above. Now suppose  $|F| > 1$ . Since  $F$  is connected, it contains  $\mathbf{2}$  and thus we see  $\mathcal{C}(F^Y)$  and hence  $X$  contains  $\mathbf{2}^Y$ . Since the dual of  $Y$ ,  $Y^\partial$ , can be embedded in  $\mathbf{2}^Y$  ( $\mathbf{2}^Y$  is order-isomorphic to the lattice of up-sets of  $Y$ , and its poset of join-irreducibles is given by the principal up-sets; under inclusion, these are ordered like the dual of  $Y$ ), then  $Y^\partial \times Z$  can be embedded in  $\mathbf{2}^Y \times Z$  and thus in  $X \times Z \cong Q_0$ . Hence

$$h(P) = h(Y \times Z) = h(Y) + h(Z) = h(Y^\partial) + h(Z) = h(Y^\partial \times Z) \leq h(Q_0) = h(Q)$$

and

$$|P| = |Y \times Z| = |Y||Z| = |Y^\partial||Z| = |Y^\partial \times Z| \leq |Q_0| < |Q|,$$

so by Lemma 15 (4) and Claim 17,  $h(P) \neq 0$ , so  $h(P^P) = |P|h(P) < |Q|h(Q) = h(Q^Q)$ , a contradiction.

Thus  $|F| = 1$ , and  $|X| = 1$ , and so  $P \cong Y \times Q_0$ . If also  $|Y| = 1$ , then  $h(P) = h(Q)$  but  $|Q| > |P|$ , so  $h(Q^Q) > h(P^P)$ , a contradiction stemming from Lemma 15 (4), unless  $h(P) = 0$ , which would make  $P$  an antichain, contradicting Claim 17. Thus  $Q_0$  is a proper direct factor of  $P$ .

We have that  $P$  and  $Q_0$  are nonempty and connected, so Theorem 10 applies to the isomorphism we have already established,  $\mathcal{C}(P^P) \cong \mathcal{C}(Q_0^Q)$ : There exist finite, nonempty posets  $E'$ ,  $X'$ ,  $Y'$ , and  $Z'$  such that

$$P \cong \mathcal{C}(E'^{X'}), \quad Q_0 \cong \mathcal{C}(E'^{Y'}), \quad P \cong Y' \times Z', \quad \text{and} \quad Q \cong X' \times Z'.$$

Since  $P$  is connected and nonempty, so are  $Y'$  and  $Z'$ , and hence  $Z'$  divides every component of  $Q$ , in particular,  $Q_0$ —say,  $Z' \times T \cong Q_0$  for some connected, nonempty poset  $T$ . Again,  $Y'$  is connected and so is  $E'$ , by Proposition 1 (3), since  $Q_0$  is connected. By Theorem 2 (3), since  $T$  is a factor of  $Q_0 \cong \mathcal{C}(E'^{Y'})$ , there exists a finite, nonempty poset  $H$  such that  $T \cong \mathcal{C}(H^{Y'})$ ;  $H$  is connected by Proposition 1 (3).

*Case 1.*  $|H| > 1$ . Then since  $H$  is connected, it contains  $\mathbf{2}$ , and hence  $T$  contains  $\mathbf{2}^{Y'}$ , which contains  $Y'^{\partial}$  (we get the dual for the same reason as before), and  $Q_0 \cong Z' \times T$  contains  $Z' \times Y'^{\partial}$ , so  $|Q_0| \geq |Z' \times Y'^{\partial}| = |Z'| |Y'^{\partial}| = |Z'| |Y'| = |Z' \times Y'| = |P|$ . As we already know that  $P$  properly contains  $Q_0$ , we have a contradiction.

*Case 2.*  $|H| = 1$ . Then  $|T| = 1$  because  $H^{Y'}$  consists only of a constant map, and  $Z' \cong Q_0$ , because  $Z' \times \mathbf{1} \cong Z'$  and  $T \cong \mathbf{1}$ , so  $Q_0$  is a factor of  $Q$ , because  $Q \cong X' \times Z'$  and  $Z' \cong Q_0$ . But then  $h(Q_0) = h(Q) = h(Q_0) + h(X')$ , so  $h(X') = 0$  and  $X'$  is an antichain. But if  $|X'| > 1$ , then  $Q$  has two components of maximum height, a contradiction.

Hence  $X'$  is a singleton, so  $Q \cong X' \times Z' \cong Z' \cong Q_0$  and  $Q$  is connected, a contradiction.  $\square$

Perhaps one could prove that  $P^P \cong Q^Q$  implies  $P \cong Q$  if  $P$  and  $Q$  are finite and nonempty and  $P$  is directly irreducible. Of course, the remaining problem is to show that  $P^P \cong Q^Q$  implies  $P \cong Q$  if  $P$  and  $Q$  are finite and nonempty (or find a counterexample).

*Acknowledgement.* The author would like to thank Bernd Schröder for his detailed critique of the paper. The author would like to thank Dwight Duffus for sending him a reference and sections of his thesis (which the author has had since graduate school but did not at first have handy when writing this paper), and Dominic van der Zypen and Zeinab Bandpey for their willingness to look at these results. The article was accepted by the journal *Order*, but the author withdrew the paper because he did not want to sign Springer Verlag's licence to publish. (Springer Verlag publishes *Order*.) The author thanks the editor and referees for the journal *Order* for comments improving the paper, including the proof of Claim 17 of Theorem 16.

### References

- [1] *G. Birkhoff*: An extended arithmetic. *Duke Math. J.* 3 (1937), 311–316. [zbl](#) [MR](#) [doi](#)
- [2] *G. Birkhoff*: *Lattice Theory*. American Mathematical Society Colloquium Publications 25. AMS, New York, 1948. [zbl](#) [MR](#)
- [3] *B. A. Davey, H. A. Priestley*: *Introduction to Lattices and Order*. Cambridge University Press, Cambridge, 2002. [zbl](#) [MR](#) [doi](#)

- [4] *D. A. Duffus*: Toward a Theory of Finite Partially Ordered Sets: Ph.D. Thesis. University of Calgary, Calgary, 1978. [MR](#)
- [5] *D. Duffus*: Automorphisms and products of ordered sets. *Algebra Univers.* *19* (1984), 366–369. [zbl](#) [MR](#) [doi](#)
- [6] *D. Duffus*: Powers of ordered sets. *Order* *1* (1984), 83–92. [zbl](#) [MR](#) [doi](#)
- [7] *D. Duffus, I. Rival*: A logarithmic property for exponents of partially ordered sets. *Can. J. Math.* *30* (1978), 797–807. [zbl](#) [MR](#) [doi](#)
- [8] *D. Duffus, R. Wille*: A theorem on partially ordered sets of order-preserving mappings. *Proc. Am. Math. Soc.* *76* (1979), 14–16. [zbl](#) [MR](#) [doi](#)
- [9] *J. D. Farley*: An issue raised in 1978 by a then-future editor-in-chief of the Journal “Order”: Does the endomorphism poset of a finite connected poset tell us that the poset is connected? Available at <https://arxiv.org/abs/2005.03255v1> (2020), 12 pages.
- [10] *J. D. Farley*: Another problem of Jónsson and McKenzie from 1982: Refinement properties for connected powers of posets. *Algebra Univers.* *82* (2021), Article ID 48, 6 pages. [zbl](#) [MR](#) [doi](#)
- [11] *J. D. Farley*: Poset exponentiation and a counterexample Birkhoff said in 1942 he did not have. *Order* *39* (2022), 243–250. [zbl](#) [MR](#) [doi](#)
- [12] *B. Jónsson, R. McKenzie*: Powers of partially ordered sets: Cancellation and refinement properties. *Math. Scand.* *51* (1982), 87–120. [zbl](#) [MR](#) [doi](#)
- [13] *L. Lovász*: Operations with structures. *Acta Math. Acad. Sci. Hung.* *18* (1967), 321–328. [zbl](#) [MR](#) [doi](#)
- [14] *R. McKenzie*: Cardinal multiplication of structures with a reflexive relation. *Fundam. Math.* *70* (1971), 59–101. [zbl](#) [MR](#) [doi](#)
- [15] *R. McKenzie*: Arithmetic of finite ordered sets: Cancellation of exponents, II. *Order* *17* (2000), 309–332. [zbl](#) [MR](#) [doi](#)
- [16] *R. McKenzie*: The zig-zag property and exponential cancellation of ordered sets. *Order* *20* (2003), 185–221. [zbl](#) [MR](#) [doi](#)
- [17] *B. Schröder*: *Ordered Sets: An Introduction with Connections from Combinatorics to Topology*. Birkhäuser, Basel, 2016. [zbl](#) [MR](#) [doi](#)

*Author’s address:* Jonathan David Farley, Department of Mathematics, Morgan State University, 1700 E. Cold Spring Lane, Baltimore, MD 21251, United States of America, e-mail: [lattice.theory@gmail.com](mailto:lattice.theory@gmail.com).