

Władysław Kulpa

Some remarks on paracompactness of GO-spaces

In: Zdeněk Frolík (ed.): Abstracta. 7th Winter School on Abstract Analysis.  
Czechoslovak Academy of Sciences, Praha, 1979. pp. 40--41.

Persistent URL: <http://dml.cz/dmlcz/701144>

## Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic,  
1979

Institute of Mathematics of the Academy of Sciences of the Czech Republic  
provides access to digitized documents strictly for personal use. Each copy of any  
part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery  
and stamped with digital signature within the project *DML-CZ*:  
*The Czech Digital Mathematics Library* <http://project.dml.cz>

## Some remarks on paracompactness of GO-spaces

by

W. Kulpa (Katowice).

Let  $X = (X, \leq)$  be a linearly ordered set. A subset  $C \subseteq X$  is said to be convex, whenever  $a, b \in C$  and  $a \leq b$  imply that  $[a, b] \subseteq C$ . A convex set  $C \subseteq A$ ,  $A \subseteq X$ , is called a convexity-component of  $A$ , whenever  $C' \cap C \neq \emptyset$  implies  $C' \subseteq C$  for each convex set  $C' \subseteq A$ .

A linearly ordered topological space  $X$  is a triple  $X = (X, \leq, \lambda(\leq))$  where  $(X, \leq)$  is a linearly ordered set on which a topology is defined by the subbase of all sets  $(\leftarrow, a), (b, \rightarrow)$  with  $a, b \in X$ .

A generalized ordered space  $X$  (abbreviated GO-space) is a triple  $X = (X, \leq, \tau)$ , where  $\tau \supset \lambda(\leq)$  is a topology with a base consisting of convex subsets of  $X$ .

A continuous map  $f: X \rightarrow M$  from a GO-space  $X$  is said to be convexity-paracompact (convexity-zero-dimensional) iff each convexity component of  $f^{-1}(m)$ ,  $m \in M$ , is paracompact (is a one-point set).

Theorem 1. If a GO-space  $X$  has a convexity-paracompact map into a metric space, then  $X$  is paracompact.

Theorem 2. If a GO-space  $X$  has a perfect map onto a Dieudonné complete space, then  $X$  is paracompact.

Theorem 1 strengthens a result of Faber [1] who has proved that each GO-space which has a convexity-zero-dimensional map into a metric space must be paracompact. Theorem 2 is a strengthening of a result of Lutzer [4] who has proved that each Dieudonné complete GO-space is paracompact.

The above theorems one can obtain from the Pressing-Down Lemma and from the following results:

Factorization Theorem [3]. Let  $f: X \rightarrow M$  be a continuous map from a GO-space  $X = (X, \langle, \tau)$  into a metric space  $M$ . Then there exists a metric GO-space  $Z = (Z, \langle, \mathcal{T})$  and continuous maps  $g: X \xrightarrow{\text{onto}} Z$ ,  $h: Z \rightarrow M$  such that  $f = h \circ g$  and  $g(x) \neq g(y)$  whenever  $x \leq y$ , for each  $x, y \in X$ .

Lemma [3]. If a GO-space  $X$  has a continuous map  $f: X \rightarrow M$  into a space  $M$  with a  $\mathcal{G}_\delta$ -diagonal such that for each  $m \in M$ ,  $f^{-1}(m)$  is paracompact, then  $X$  paracompact.

Lemma [3]. If  $f: S \rightarrow M$  is a continuous map from a stationary set  $S \subset \kappa$ ,  $\kappa = \text{cf} \kappa > \omega$ , into a space  $M$  with a  $\mathcal{G}_\delta$ -diagonal, then there is an  $\alpha < \kappa$  such that  $f|_{S \cap [\alpha, \kappa)}$  is constant.

The last Lemma generalizes a result of Lutzer [4] who has proved that if  $f: S \rightarrow M$  is a continuous map of a stationary set  $S \subset \kappa$ ,  $\kappa = \text{cf} \kappa > \omega$ , into a metric space  $M$ , then  $\text{card}(f[S]) < \kappa$ .

Theorem [1]. A GO-space  $X$  is not paracompact iff some closed subspace of  $X$  is homeomorphic to a stationary set  $S \subset \kappa$ ,  $\kappa = \text{cf} \kappa > \omega$ .

#### References.

1. R. Engelking, D. Lutzer, Paracompactness in ordered spaces, *Fund. Math.* XCIV (1977), 49 - 58.
2. H.J. Faber, Metrizable in generalized ordered spaces, *Mathematisch Centrum, Amsterdam* 1974.
3. W. Kubiś, A factorization theorem for GO-spaces, preprint.
4. D.J. Lutzer, Stationary sets and paracompactness in ordered spaces: a survey, *Seminar Uniform Spaces* (directed by E. Enohik) (1976 - 1977), 1 - 23.