

Aurelian Cernea

On the Solution Set of a Nonconvex Nonclosed Second Order Inclusion

In: Karol Mikula (ed.): Proceedings of Equadiff 14, Conference on Differential Equations and Their Applications, Bratislava, July 24-28, 2017. Slovak University of Technology in Bratislava, SPEKTRUM STU Publishing, Bratislava, 2017. pp. 21–28.

Persistent URL: <http://dml.cz/dmlcz/703028>

Terms of use:

© Slovak University of Technology in Bratislava, 2017

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON THE SOLUTION SET OF A NONCONVEX NONCLOSED SECOND-ORDER EVOLUTION INCLUSION

AURELIAN CERNEA*

Abstract. We consider a nonconvex and nonclosed second-order evolution inclusion and we prove the arcwise connectedness of the set of its mild solutions.

Key words. set-valued contraction, fixed point, solution set

AMS subject classifications. 34A60

1. Introduction. This paper is concerned with the following problem

$$(1.1) \quad x'' \in A(t)x + F(t, x, H(t, x)), \quad x(0) = x_0, \quad x'(0) = y_0,$$

where X is a real separable Banach space, $\mathcal{P}(X)$ is the family of all subsets of X , $I = [0, T]$, $F(., ., .) : I \times X^2 \rightarrow \mathcal{P}(X)$, $H(., .) : I \times X \rightarrow \mathcal{P}(X)$ and $\{A(t)\}_{t \geq 0}$ is a family of linear closed operators from X into X that generates an evolution system of operators $\{\mathcal{U}(t, s)\}_{t, s \in [0, T]}$. The general framework of evolution operators $\{A(t)\}_{t \geq 0}$ that define problem (1.1) has been developed by Kozak ([10]) and improved by Henriquez ([8]).

When F does not depend on the last variable (1.1) reduces to

$$(1.2) \quad x'' \in A(t)x + F(t, x), \quad x(0) = x_0, \quad x'(0) = y_0.$$

Existence results and qualitative properties of the solutions of problem (1.2) may be found in [1, 2, 3, 4, 8, 9] etc. In all the papers concerned with the set-valued framework, the set-valued map F is assumed to be at least closed-valued. Such an assumption is quite natural in order to obtain good properties of the solution set, but it is interesting to investigate the problem when the right-hand side of the multivalued equation may have nonclosed values.

Following the approach in [12] we consider the problem (1.1), where F and H are closed-valued multifunctions Lipschitzian with respect to the second variable and F is contractive in the third variable. Obviously, the right-hand side of the differential inclusion in (1.1) is in general neither convex nor closed. We prove the arcwise connectedness of the solution set of problem (1.1). The main tool is a result ([11, 12]) concerning the arcwise connectedness of the fixed point set of a class of nonconvex nonclosed set-valued contractions.

We note that similar results for other classes of differential inclusions may be found in our previous papers [5, 6, 7].

The paper is organized as follows: in Section 2 we recall some preliminary results that we use in the sequel and in Section 3 we prove our main result.

2. Preliminaries. Let Z be a metric space with the distance d_Z and let 2^Z be the family of all nonempty closed subsets of Z . For $a \in Z$ and $A, B \in 2^Z$ set

*Faculty of Mathematics and Computer Science, University of Bucharest, Academiei 14, 010014 Bucharest, Romania (acernea@fmi.unibuc.ro).

$d_Z(a, B) = \inf_{b \in B} d_Z(a, b)$ and $d_Z^*(A, B) = \sup_{a \in A} d_Z(a, B)$. Denote by D_Z the Pompeiu-Hausdorff generalized metric on 2^Z defined by

$$D_Z(A, B) = \max\{d_Z^*(A, B), d_Z^*(B, A)\}, \quad A, B \in 2^Z.$$

In what follows, when the product $Z = Z_1 \times Z_2$ of metric spaces $Z_i, i = 1, 2$, is considered, it is assumed that Z is equipped with the distance $d_Z((z_1, z_2), (z'_1, z'_2)) = \sum_{i=1}^2 d_{Z_i}(z_i, z'_i)$.

Let X be a nonempty set and let $F : X \rightarrow 2^Z$ be a set-valued map from X to Z . The range of F is the set $F(X) = \cup_{x \in X} F(x)$. Let (X, \mathcal{F}) be a measurable space. The multifunction $F : X \rightarrow 2^Z$ is called measurable if $F^{-1}(\Omega) \in \mathcal{F}$ for any open set $\Omega \subset Z$, where $F^{-1}(\Omega) = \{x \in X; F(x) \cap \Omega \neq \emptyset\}$. Let (X, d_X) be a metric space. The multifunction F is called Hausdorff continuous if for any $x_0 \in X$ and every $\epsilon > 0$ there exists $\delta > 0$ such that $x \in X, d_X(x, x_0) < \delta$ implies $D_Z(F(x), F(x_0)) < \epsilon$.

Let (T, \mathcal{F}, μ) be a finite, positive, nonatomic measure space and let $(X, |\cdot|_X)$ be a Banach space. We denote by $L^1(T, X)$ the Banach space of all (equivalence classes of) Bochner integrable functions $u : T \rightarrow X$ endowed with the norm

$$|u|_{L^1(T, X)} = \int_T |u(t)|_X d\mu$$

A nonempty set $K \subset L^1(T, X)$ is called decomposable if, for every $u, v \in K$ and every $A \in \mathcal{F}$, one has

$$\chi_A \cdot u + \chi_{T \setminus A} \cdot v \in K$$

where $\chi_B, B \in \mathcal{F}$ indicates the characteristic function of B .

A metric space Z is called an absolute retract if, for any metric space X and any nonempty closed set $X_0 \subset X$, every continuous function $g : X_0 \rightarrow Z$ has a continuous extension $g : X \rightarrow Z$ over X . It is obvious that every continuous image of an absolute retract is an arcwise connected space.

In what follows we recall some preliminary results that are the main tools in the proof of our result.

Let (T, \mathcal{F}, μ) be a finite, positive, nonatomic measure space, S a separable Banach space and let $(X, |\cdot|_X)$ be a real Banach space. To simplify the notation we write E in place of $L^1(T, X)$. The proofs of the next two lemmas may be found in [11].

LEMMA 2.1. *Assume that $\phi : S \times E \rightarrow 2^E$ and $\psi : S \times E \times E \rightarrow 2^E$ are Hausdorff continuous multifunctions with nonempty, closed, decomposable values, satisfying the following conditions*

a) *There exists $L \in [0, 1)$ such that, for every $s \in S$ and every $u, u' \in E$,*

$$D_E(\phi(s, u), \phi(s, u')) \leq L|u - u'|_E.$$

b) *There exists $M \in [0, 1)$ such that $L + M < 1$ and for every $s \in S$ and every $(u, v), (u', v') \in E \times E$,*

$$D_E(\psi(s, u, v), \psi(s, u', v')) \leq M(|u - u'|_E + |v - v'|_E).$$

Set $\text{Fix}(\Gamma(s, \cdot)) = \{u \in E; u \in \Gamma(s, u)\}$, where $\Gamma(s, u) = \psi(s, u, \phi(s, u)), (s, u) \in S \times E$. Then

1) *For every $s \in S$ the set $\text{Fix}(\Gamma(s, \cdot))$ is nonempty and arcwise connected.*

2) For any $s_i \in S$, and any $u_i \in \text{Fix}(\Gamma(s, \cdot))$, $i = 1, \dots, p$ there exists a continuous function $\gamma : S \rightarrow E$ such that $\gamma(s) \in \text{Fix}(\Gamma(s, \cdot))$ for all $s \in S$ and $\gamma(s_i) = u_i$, $i = 1, \dots, p$.

LEMMA 2.2. Let $U : T \rightarrow 2^X$ and $V : T \times X \rightarrow 2^X$ be two nonempty closed-valued multifunctions satisfying the following conditions

a) U is measurable and there exists $r \in L^1(T)$ such that $D_X(U(t), \{0\}) \leq r(t)$ for almost all $t \in T$.

b) The multifunction $t \rightarrow V(t, x)$ is measurable for every $x \in X$.

c) The multifunction $x \rightarrow V(t, x)$ is Hausdorff continuous for all $t \in T$.

Let $v : T \rightarrow X$ be a measurable selection from $t \rightarrow V(t, U(t))$.

Then there exists a selection $u \in L^1(T, X)$ such that $v(t) \in V(t, u(t))$, $t \in T$.

In what follows $\{A(t)\}_{t \geq 0}$ is a family of linear closed operators from X into X that generates an evolution system of operators $\{\mathcal{U}(t, s)\}_{t, s \in I}$. By hypothesis the domain of $A(t)$, $D(A(t))$ is dense in X and is independent of t . The following definition is taken from [8, 10].

DEFINITION 2.3. A family of bounded linear operators $\mathcal{U}(t, s) : X \rightarrow X$, $(t, s) \in \Delta := \{(t, s) \in I \times I; s \leq t\}$ is called an evolution operator of the equation

$$(2.1) \quad x''(t) = A(t)x(t)$$

if

i) For any $x \in X$, the map $(t, s) \rightarrow \mathcal{U}(t, s)x$ is continuously differentiable and

a) $\mathcal{U}(t, t) = 0$, $t \in I$.

b) If $t \in I$, $x \in X$ then $\frac{\partial}{\partial t} \mathcal{U}(t, s)x|_{t=s} = x$ and $\frac{\partial}{\partial s} \mathcal{U}(t, s)x|_{t=s} = -x$.

ii) If $(t, s) \in \Delta$, then $\frac{\partial}{\partial s} \mathcal{U}(t, s)x \in D(A(t))$, the map $(t, s) \rightarrow \mathcal{U}(t, s)x$ is of class C^2 and

a) $\frac{\partial^2}{\partial t^2} \mathcal{U}(t, s)x \equiv A(t)\mathcal{U}(t, s)x$.

b) $\frac{\partial^2}{\partial s^2} \mathcal{U}(t, s)x \equiv \mathcal{U}(t, s)A(t)x$.

c) $\frac{\partial^2}{\partial s \partial t} \mathcal{U}(t, s)x|_{t=s} = 0$.

iii) If $(t, s) \in \Delta$, then there exist $\frac{\partial^3}{\partial t^2 \partial s} \mathcal{U}(t, s)x$, $\frac{\partial^3}{\partial s^2 \partial t} \mathcal{U}(t, s)x$ and

a) $\frac{\partial^3}{\partial t^2 \partial s} \mathcal{U}(t, s)x \equiv A(t) \frac{\partial}{\partial s} \mathcal{U}(t, s)x$ and the map $(t, s) \rightarrow A(t) \frac{\partial}{\partial s} \mathcal{U}(t, s)x$ is continuous.

b) $\frac{\partial^3}{\partial s^2 \partial t} \mathcal{U}(t, s)x \equiv \frac{\partial}{\partial t} \mathcal{U}(t, s)A(s)x$.

As an example for equation (2.1) one may consider the problem (e.g., [8])

$$\frac{\partial^2 z}{\partial t^2}(t, \tau) = \frac{\partial^2 z}{\partial \tau^2}(t, \tau) + a(t) \frac{\partial z}{\partial t}(t, \tau), \quad t \in [0, T], \tau \in [0, 2\pi],$$

$$z(t, 0) = z(t, \pi) = 0, \quad \frac{\partial z}{\partial \tau}(t, 0) = \frac{\partial z}{\partial \tau}(t, 2\pi), \quad t \in [0, T],$$

where $a(\cdot) : I \rightarrow \mathbf{R}$ is a continuous function. This problem is modeled in the space $X = L^2(\mathbf{R}, \mathbf{C})$ of 2π -periodic 2-integrable functions from \mathbf{R} to \mathbf{C} , $A_1 z = \frac{d^2 z(\tau)}{d\tau^2}$ with domain $H^2(\mathbf{R}, \mathbf{C})$ the Sobolev space of 2π -periodic functions whose derivatives belong to $L^2(\mathbf{R}, \mathbf{C})$. It is well known that A_1 is the infinitesimal generator of strongly continuous cosine functions $C(t)$ on X . Moreover, A_1 has discrete spectrum; namely the spectrum of A_1 consists of eigenvalues $-n^2$, $n \in \mathbf{Z}$ with associated eigenvectors $z_n(\tau) = \frac{1}{\sqrt{2\pi}} e^{in\tau}$, $n \in \mathbf{N}$. The set $\{z_n\}$, $n \in \mathbf{N}$ is an orthonormal basis of X .

In particular, $A_1 z = \sum_{n \in \mathbf{Z}} -n^2 \langle z, z_n \rangle z_n$, $z \in D(A_1)$. The cosine function is given by $C(t)z = \sum_{n \in \mathbf{Z}} \cos(nt) \langle z, z_n \rangle z_n$ with the associated sine function $S(t)z = t \langle z, z_0 \rangle z_0 + \sum_{n \in \mathbf{Z} \setminus \{0\}} \frac{\sin(nt)}{n} \langle z, z_n \rangle z_n$.

For $t \in I$ define the operator $A_2(t)z = a(t) \frac{dz(\tau)}{d\tau}$ with domain $D(A_2(t)) = H^1(\mathbf{R}, \mathbf{C})$. Set $A(t) = A_1 + A_2(t)$. It has been proved in [10] that this family generates an evolution operator as in Definition 2.3.

DEFINITION 2.4. A continuous mapping $x(\cdot) \in C(I, X)$ is called a mild solution of problem (1.1) if there exists a (Bochner) integrable function $f(\cdot) \in L^1(I, X)$ such that

$$(2.2) \quad f(t) \in F(t, x(t)) \quad a.e. (I),$$

$$(2.3) \quad x(t) = -\frac{\partial}{\partial s} \mathcal{U}(t, 0)x_0 + \mathcal{U}(t, 0)y_0 + \int_0^t \mathcal{U}(t, s)f(s)ds, \quad t \in I.$$

We shall call $(x(\cdot), f(\cdot))$ a trajectory-selection pair of (1.1) if $f(\cdot)$ verifies (2.2) and $x(\cdot)$ is defined by (2.3).

We shall use the following notations for the solution sets of (1.1).

$$(2.4) \quad \mathcal{S}(x_0, y_0) = \{x(\cdot); \quad x(\cdot) \text{ is a mild solution of (1.1)}\}.$$

In order to study problem (1.1) we introduce the following hypothesis.

HYPOTHESIS 2.5. i) There exists an evolution operator $\{\mathcal{U}(t, s)\}_{t, s \in I}$ associated to the family $\{A(t)\}_{t \geq 0}$.

ii) There exist $M, M_0 \geq 0$ such that $|\mathcal{U}(t, s)|_{B(X)} \leq M$, $|\frac{\partial}{\partial s} \mathcal{U}(t, s)| \leq M_0$, for all $(t, s) \in \Delta$.

$F : I \times X \times X \rightarrow \mathcal{P}(X)$ and $H : I \times X \rightarrow \mathcal{P}(X)$ are two set-valued maps with nonempty closed values, satisfying

iii) The set-valued maps $t \rightarrow F(t, u, v)$ and $t \rightarrow H(t, u)$ are measurable for all $u, v \in X$.

iv) There exist $l(\cdot) \in L^1(I, \mathbf{R})$ such that, for every $u, u' \in X$,

$$D(H(t, u), H(t, u')) \leq l(t)|u - u'| \quad a.e. (I).$$

v) There exist $m(\cdot) \in L^1(I, \mathbf{R})$ and $\theta \in [0, 1)$ such that, for every $u, v, u', v' \in X$,

$$D(F(t, u, v), F(t, u', v')) \leq m(t)|u - u'| + \theta|v - v'| \quad a.e. (I).$$

vi) There exist $f, g \in L^1(I, \mathbf{R})$ such that

$$d(0, F(t, 0, 0)) \leq f(t), \quad d(0, H(t, 0)) \leq g(t) \quad a.e. (I).$$

In what follows $N(t) = \max\{l(t), m(t)\}$, $t \in I$, $N^*(t) = \int_0^t N(s)ds$.

Given $\alpha \in \mathbf{R}$ we denote by L^1 the Banach space of all (equivalence classes of) Lebesgue measurable functions $\sigma : I \rightarrow X$ endowed with the norm

$$|\sigma|_1 = \int_0^T e^{-\alpha N^*(t)} |\sigma(t)| dt.$$

3. Main result. Even if the multifunction from the right-hand side of (1.1) has, in general, nonclosed nonconvex values, its solution set $\mathcal{S}(x_0, y_0)$ defined in (2.4) has some meaningful properties, stated in theorem below.

THEOREM 3.1. *Assume that Hypothesis 2.5 is satisfied and let $\alpha > \frac{2M}{1-\theta}$. Then*

1) *For every $(x_0, y_0) \in X \times X$, the solution set $\mathcal{S}(x_0, y_0)$ is nonempty and arcwise connected in the space $C(I, X)$.*

2) *For any $(\xi_i, \mu_i) \in X \times X$ and any $x_i \in \mathcal{S}(\xi_i, \mu_i)$, $i = 1, \dots, p$, there exists a continuous function $s : X \times X \rightarrow C(I, X)$ such that $s(\xi, \mu) \in \mathcal{S}(\xi, \mu)$ for any $(\xi, \mu) \in X \times X$ and $s(\xi_i, \mu_i) = x_i$, $i = 1, \dots, p$.*

3) *The set $\mathcal{S} = \cup_{(\xi, \mu) \in X \times X} \mathcal{S}(\xi, \mu)$ is arcwise connected in $C(I, X)$.*

Proof. 1) For $(\xi, \mu) \in X \times X$ and $f \in L^1$, set

$$(3.1) \quad x_{\xi, \mu}(t) = -\frac{\partial}{\partial s} \mathcal{U}(t, 0)\xi + \mathcal{U}(t, 0)\mu + \int_0^t \mathcal{U}(t, s)f(s)ds$$

and consider $\lambda : X \times X \rightarrow C(I, X)$ defined by $\lambda(\xi, \mu)(t) = -\frac{\partial}{\partial s} \mathcal{U}(t, 0)\xi + \mathcal{U}(t, 0)\mu$.

We prove that the multifunctions $\phi : X \times X \times L^1 \rightarrow 2^{L^1}$ and $\psi : X \times X \times L^1 \times L^1 \rightarrow 2^{L^1}$ given by

$$\phi((\xi, \mu), u) = \{v \in L^1; \quad v(t) \in H(t, x_{\xi, \mu}(t)) \quad a.e.(I)\},$$

$$\psi((\xi, \mu), u, v) = \{w \in L^1; \quad w(t) \in F(t, x_{\xi, \mu}(t), v(t)) \quad a.e.(I)\},$$

$(\xi, \mu) \in X \times X$, $u, v \in L^1$ satisfy the hypotheses of Lemma 2.1.

Since $x_{\xi, \mu}(\cdot)$ is measurable and H satisfies Hypothesis 2.5 iii) and iv), the multifunction $t \rightarrow H(t, x_{\xi, \mu}(t))$ is measurable and nonempty closed-valued, it has a measurable selection. Therefore due to Hypothesis 2.5 vi), the set $\phi((\xi, \mu), u)$ is nonempty. The fact that the set $\phi((\xi, \mu), u)$ is closed and decomposable follows by a simple computation. In the same way we obtain that $\psi((\xi, \mu), u, v)$ is a nonempty closed decomposable set.

Pick $((\xi, \mu), u), ((\xi_1, \mu_1), u_1) \in X \times X \times L^1$ and choose $v \in \phi((\xi, \mu), u)$. For each $\varepsilon > 0$ there exists $v_1 \in \phi((\xi_1, \mu_1), u_1)$ such that, for every $t \in I$, one has

$$\begin{aligned} |v(t) - v_1(t)| &\leq D(H(t, x_{\xi, \mu}(t)), H(t, x_{\xi_1, \mu_1}(t))) + \varepsilon \leq \\ l(t)[M_0|\xi - \xi_1| + M|\mu - \mu_1| + M \int_0^t |u(s) - u_1(s)|ds] + \varepsilon. \end{aligned}$$

Hence

$$\begin{aligned} |v - v_1|_1 &\leq [M_0|\xi - \xi_1| + M|\mu - \mu_1|] \int_0^T e^{-\alpha N^*(t)} l(t)dt + M \int_0^T e^{-\alpha N^*(t)} \\ l(t)(\int_0^t |u(s) - u_1(s)|ds)dt + \varepsilon T &\leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + \frac{M}{\alpha}|u - u_1|_1 + \varepsilon T \end{aligned}$$

for any $\varepsilon > 0$.

This implies

$$d_{L^1}(v, \phi((\xi_1, \mu_1), u_1)) \leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + \frac{M}{\alpha}|u - u_1|_1$$

for all $v \in \phi((\xi, \mu), u)$. Therefore,

$$d_{L^1}^*(\phi((\xi, \mu), u), \phi((\xi_1, \mu_1), u_1)) \leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + \frac{M}{\alpha}|u - u_1|_1$$

Consequently,

$$D_{L^1}(\phi((\xi, \mu), u), \phi((\xi_1, \mu_1), u_1)) \leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + \frac{M}{\alpha}|u - u_1|_1$$

which shows that ϕ is Hausdorff continuous and satisfies the assumptions of Lemma 2.1.

Pick $((\xi, \mu), u, v), ((\xi_1, \mu_1), u_1, v_1) \in X \times X \times L^1 \times L^1$ and choose $w \in \psi((\xi, \mu), u, v)$. Then, as before, for each $\varepsilon > 0$ there exists $w_1 \in \psi((\xi_1, \mu_1), u_1, v_1)$ such that for every $t \in I$

$$|w(t) - w_1(t)| \leq D(F(t, x_{\xi, \mu}(t), v(t)), F(t, x_{\xi_1, \mu_1}(t), v_1(t))) + \varepsilon \leq m(t)[M_0|\xi - \xi_1| + M|\mu - \mu_1| + M \int_0^t |u(s) - u_1(s)| ds] + \theta|v(t) - v_1(t)| + \varepsilon.$$

Hence

$$\begin{aligned} |w - w_1|_1 &\leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + \frac{M}{\alpha}|u - u_1|_1 + \theta|v - v_1|_1 + \varepsilon T \\ &\leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + (\frac{M}{\alpha} + \theta)(|u - u_1|_1 + |v - v_1|_1) + \varepsilon T \\ &\leq \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + (\frac{M}{\alpha} + \theta)d_{L^1 \times L^1}((u, v), (u_1, v_1)) + \varepsilon T. \end{aligned}$$

As above, we deduce that

$$\begin{aligned} D_{L^1}(\psi((\xi, \mu), u, v), \psi((\xi_1, \mu_1), u_1, v_1)) &\leq \\ \frac{1}{\alpha}[M_0|\xi - \xi_1| + M|\mu - \mu_1|] + (\frac{M}{\alpha} + \theta)d_{L^1 \times L^1}((u, v), (u_1, v_1)). \end{aligned}$$

namely, the multifunction ψ is Hausdorff continuous and satisfies the hypothesis of Lemma 2.1.

Define $\Gamma((\xi, \mu), u) = \psi((\xi, \mu), u, \phi((\xi, \mu), u)), ((\xi, \mu), u) \in X^2 \times L^1$. According to Lemma 2.1, the set $Fix(\Gamma((\xi, \mu), \cdot)) = \{u \in L^1; u \in \Gamma((\xi, \mu), u)\}$ is nonempty and arcwise connected in $L^1(I, X)$. Moreover, for fixed $(\xi_i, \mu_i) \in X^2$ and $u_i \in Fix(\Gamma((\xi_i, \mu_i), \cdot)), i = 1, \dots, p$, there exists a continuous function $\gamma : X^2 \rightarrow L^1$ such that

$$(3.2) \quad \gamma((\xi, \mu)) \in Fix(\Gamma((\xi, \mu), \cdot)), \quad \forall (\xi, \mu) \in X^2,$$

$$(3.3) \quad \gamma((\xi_i, \mu_i)) = u_i, \quad i = 1, \dots, p.$$

We shall prove that

$$(3.4) \quad Fix(\Gamma((\xi, \mu), \cdot)) = \{u \in L^1; u(t) \in F(t, x_{\xi, \mu}(t), H(t, x_{\xi, \mu}(t))) \text{ a.e. } (I)\}.$$

Denote by $A(\xi, \mu)$ the right-hand side of (3.4). If $u \in Fix(\Gamma((\xi, \mu), \cdot))$ then there is $v \in \phi((\xi, \mu), v)$ such that $u \in \psi((\xi, \mu), u, v)$. Therefore, $v(t) \in H(t, x_{\xi, \mu}(t))$ and

$$u(t) \in F(t, x_{\xi, \mu}(t), v(t)) \subset F(t, x_{\xi, \mu}(t), H(t, x_{\xi, \mu}(t))) \text{ a.e. } (I),$$

so that $Fix(\Gamma((\xi, \mu), \cdot)) \subset A(\xi, \mu)$.

Let now $u \in A(\xi, \mu)$. By Lemma 2.2, there exists a selection $v \in L^1$ of the multifunction $t \rightarrow H(t, x_{\xi, \mu}(t))$ satisfying

$$u(t) \in F(t, x_{\xi, \mu}(t), v(t)) \text{ a.e. } (I).$$

Hence, $v \in \phi((\xi, \mu), v)$, $u \in \psi((\xi, \mu), u, v)$ and thus $u \in \Gamma((\xi, \mu), u)$, which completes the proof of (3.4).

We next note that the function $T : L^1 \rightarrow C(I, X)$,

$$T(u)(t) := \int_0^t \mathcal{U}(t, s)u(s)ds$$

is continuous and one has

$$(3.5) \quad \mathcal{S}(\xi, \mu) = \lambda(\xi, \mu) + T(\text{Fix}(\Gamma((\xi, \mu), \cdot))), \quad (\xi, \mu) \in X^2.$$

Since $\text{Fix}(\Gamma((\xi, \mu), \cdot))$ is nonempty and arcwise connected in L^1 , the set $\mathcal{S}(\xi, \mu)$ has the same properties in $C(I, X)$.

2) Let $(\xi_i, \mu_i) \in X^2$ and let $x_i \in \mathcal{S}(\xi_i, \mu_i)$, $i = 1, \dots, p$ be fixed. By (3.5) there exists $v_i \in \text{Fix}(\Gamma((\xi_i, \mu_i), \cdot))$ such that

$$x_i = \lambda(\xi_i, \mu_i) + T(v_i), \quad i = 1, \dots, p.$$

If $\gamma : X^2 \rightarrow L^1$ is a continuous function satisfying (3.2) and (3.3) we define, for every $(\xi, \mu) \in X^2$,

$$s(\xi, \mu) = \lambda(\xi, \mu) + T(\gamma(\xi, \mu)).$$

Obviously, the function $s : X \rightarrow C(I, X)$ is continuous, $s(\xi, \mu) \in \mathcal{S}(\xi, \mu)$ for all $(\xi, \mu) \in X^2$ and

$$s(\xi_i, \mu_i) = \lambda(\xi_i, \mu_i) + T(\gamma(\xi_i, \mu_i)) = \lambda(\xi_i, \mu_i) + T(v_i) = x_i, \quad i = 1, \dots, p.$$

3) Let $x_1, x_2 \in \mathcal{S} = \cup_{(\xi, \mu) \in X^2} \mathcal{S}(\xi, \mu)$ and choose $(\xi_i, \mu_i) \in X^2$, $i = 1, 2$ such that $x_i \in \mathcal{S}(\xi_i, \mu_i)$, $i = 1, 2$. From the conclusion of 2) we deduce the existence of a continuous function $s : X^2 \rightarrow C(I, X)$ satisfying $s(\xi_i, \mu_i) = x_i$, $i = 1, 2$ and $s(\xi, \mu) \in \mathcal{S}(\xi, \mu)$, $(\xi, \mu) \in X^2$. Let $h : [0, 1] \rightarrow X^2$ be a continuous mapping such that $h(0) = (\xi_1, \mu_1)$ and $h(1) = (\xi_2, \mu_2)$. Then the function $s \circ h : [0, 1] \rightarrow C(I, X)$ is continuous and verifies

$$s \circ h(0) = x_1, \quad s \circ h(1) = x_2, \quad s \circ h(\tau) \in \mathcal{S}(h(\tau)) \subset \mathcal{S}, \quad \tau \in [0, 1],$$

which completes the proof. \square

REFERENCES

- [1] A. BALIKI, M. BENCHOHRA, AND J.R. GRAEF, *Global existence and stability of second order functional evolution equations with infinite delay*, Electronic J. Qual. Theory Differ. Equations, 2016, no. 23, (2016), pp. 1–10.
- [2] A. BALIKI, M. BENCHOHRA, AND J.J. NIETO, *Qualitative analysis of second-order functional evolution equations*, Dynamic Syst. Appl., 24 (2015), pp. 559–572.
- [3] M. BENCHOHRA, AND I. MEDJADJ, *Global existence results for second order neutral functional differential equations with state-dependent delay*, Comment. Math. Univ. Carolin., 57 (2016), pp. 169–183.
- [4] M. BENCHOHRA, AND N. REZZOUG, *Measure of noncompactness and second-order evolution equations*, Gulf J. Math., 4 (2016), pp. 71–79.
- [5] A. CERNEA, *On the solution set of some classes of nonconvex nonclosed differential inclusions*, Portugaliae Math., 65 (2008), pp. 485–496.
- [6] A. CERNEA, *On the solution set of a nonconvex nonclosed Sturm-Liouville type differential inclusion*, Comment. Math., 49 (2009), pp. 139–146.
- [7] A. CERNEA, *On the solution set of a nonconvex nonclosed hyperbolic differential inclusion of third order*, J. Nonlinear Convex Anal., 17 (2016), pp. 1171–1179.

- [8] H.R. HENRIQUEZ, *Existence of solutions of nonautonomous second order functional differential equations with infinite delay*, *Nonlinear Anal.*, 74 (2011), pp. 3333–3352.
- [9] H.R. HENRIQUEZ, V. POBLETE, AND J.C. POZO, *Mild solutions of non-autonomous second order problems with nonlocal initial conditions*, *J. Math. Anal. Appl.*, 412 (2014), pp. 1064–1083.
- [10] M. KOZAK, *A fundamental solution of a second-order differential equation in a Banach space*, *Univ. Iagel. Acta. Math.*, 32 (1995), pp. 275–289.
- [11] S. MARANO, *Fixed points of multivalued contractions with nonclosed, nonconvex values*, *Atti. Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.*, 5 (1994), pp. 203–212.
- [12] S. MARANO, AND V. STAIUCU, *On the set of solutions to a class of nonconvex nonclosed differential inclusions*, *Acta Math. Hungar.*, 76 (1997), pp. 287–301.